

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 January 2004 (08.01.2004)

PCT

(10) International Publication Number
WO 2004/003176 A2

(51) International Patent Classification⁷: C12N

(21) International Application Number:
PCT/US2003/020964

(22) International Filing Date: 1 July 2003 (01.07.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/392,455 1 July 2002 (01.07.2002) US
60/393,423 3 July 2002 (03.07.2002) US

(71) Applicants: **THE KENNETH S. WARREN INSTITUTE, INC.** [US/US]; 712 Kitchawan Road, Ossining, NY 10562 (US). **H. LUNDBECK A/S** [DK/DK]; Ottiliavej 9, DK-2500 Valby-Copenhagen (DK).

(72) Inventors: **NIELSEN, Jacob**; Dannebrogsgade 17, 1.th., DK-1660 Copenhagen (DK). **PEDERSEN, Jan, Torleif**; Holcks Plads 14, 1, DK-2700 Bronshøj (DK). **GERWIEN, Jens**; Haraldsgade 52, 1.tv, DK-2100 Copenhagen (DK). **BAY, Katrine**; Allersgade 1, 2.th, DK-2100 Copenhagen (DK). **PEDERSEN, Lars, Ostergaard**; Ved Vigen 14, DK-2400 Copenhagen (DK). **LEIST, Marcel**; Kloverbladsgade 14, 1, DK-2500 Valby (DK). **GEIST, Marie, Aavang**; Saxtorphsveg 44, DK-2500 Valby (DK). **KALLUNKI, Pekka**; Classengade 17, 3.th, DK-2100 Copenhagen (DK). **CHRISTENSEN, Soren**; Lystbaadevej 32, DK-4040 Jyllinge (DK). **SAGER, Thomas**; Stormosevej 35, DK-2465 Smorum (DK).

BRINES, Michael; 1 Wepawaug Road, Woodbridge, CT 06525 (US). **CERAMI, Anthony**; 49 Bramblebush Drive, Croton-on-Hudson, NY 10520 (US). **CERAMI, Carla**; 121 Farrington Avenue, Sleepy Hollow, NY 10591 (US).

(74) Agents: **CORUZZI, Laura, A.** et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: RECOMBINANT TISSUE PROTECTIVE CYTOKINES AND ENCODING NUCLEIC ACIDS THEREOF FOR PROTECTION, RESTORATION, AND ENHANCEMENT OF RESPONSIVE CELLS, TISSUES, AND ORGANS

(57) Abstract: Methods and compositions are provided for protecting or enhancing a responsive cell, tissue, organ or body part function or viability *in vivo*, *in situ* or *ex vivo* in mammals, including human beings, by systemic or local administration of an erythropoietin receptor activity modulator, such as a recombinant tissue protective cytokine.



WO 2004/003176 A2

**RECOMBINANT TISSUE PROTECTIVE CYTOKINES AND ENCODING
NUCLEIC ACIDS THEREOF FOR PROTECTION, RESTORATION, AND
ENHANCEMENT OF RESPONSIVE CELLS, TISSUES, AND ORGANS**

This application claims priority to U.S. provisional patent Application No.
5 60/392,455 filed July 1, 2002, and U.S. provisional patent Application No. 60/393,423 filed
July 3, 2002, the entire contents of each of which is incorporated herein by reference in its
entirety.

1. INTRODUCTION

The present invention is directed to mutein recombinant tissue protective cytokines
10 having one or more amino acid substitutions, pharmaceutical compositions comprising such
cytokines for protecting, maintaining, enhancing, or restoring the function or viability of
responsive mammalian cells and their associated cells, tissues, and organs. This includes
the protection of excitable tissue, such as neuronal and cardiac tissue, from neurotoxins,
hypoxia, and other adverse stimuli, and the enhancement of excitable tissue function, such
15 as for facilitating learning and memory. The present invention is further drawn to
compositions for transporting or facilitating transport of a molecule via transcytosis across
an endothelial cell barrier using mutein recombinant tissue protective cytokines.

2. BACKGROUND OF THE INVENTION

For many years, the only clear physiological role of erythropoietin had been its
20 control of the production of red blood cells. Recently, several lines of evidence suggest
that erythropoietin, a member of the cytokine superfamily, performs other important
physiologic functions which may be mediated through interaction with the erythropoietin
receptor (erythropoietin-R). These actions include mitogenesis, modulation of calcium
influx into smooth muscle cells and neural cells, vasoactive action, i.e.,
25 vasoconstriction/vasodilatation, hyperactivation of platelets and effects on intermediary
metabolism. It is believed that erythropoietin provides compensatory responses that serve
to improve hypoxic cellular microenvironments as well as modulate programmed cell death
caused by metabolic stress. Although studies have established that erythropoietin injected

intracranially protects neurons against hypoxic neuronal injury, intracranial administration is an impractical and by and large unacceptable route of administration for therapeutic use, particularly for normal individuals. Furthermore, previous studies of anemic patients given erythropoietin have concluded that peripherally-administered erythropoietin is not
5 transported into the brain (Marti *et al.*, 1997, *Kidney Int.* 51:416-8; Juul *et al.*, 1999, *Pediatr. Res.* 46:543-547; Buemi *et al.*, 2000, *Nephrol. Dial. Transplant.* 15:422-433.).

Various modified forms of erythropoietin have been described with activities directed towards improving the erythropoietic activity of the molecule, such as those having altered amino acids at the carboxy terminus described in U.S. Patent 5,457,089 and in U.S.
10 Patent 4,835,260; erythropoietin isoforms with various numbers of sialic acid residues per molecule, such as those described in U.S. Patent 5,856,298; polypeptides described in U.S. Patent 4,703,008; agonists described in U.S. Patent 5,767,078; peptides which bind to the erythropoietin receptor as described in U.S. Patents 5,773,569 and 5,830,851; and small-molecule mimetics as described in U.S. Patent 5,835,382.

15 It is towards the use of a recombinant tissue protective cytokine for protecting, maintaining, enhancing, or restoring responsive cells and associated cells, tissues, and, organs *in situ* as well as *ex vivo*, and to delivery of a recombinant tissue protective cytokine across an endothelial cell barrier for the purpose of protecting and enhancing responsive cells and associated cells, tissues, and organs distal to the vasculature, or to carry associated
20 molecules, that the present invention is directed.

3. BRIEF SUMMARY OF THE INVENTION

In one aspect, the present invention is directed to the use of various forms of recombinant tissue protective cytokines for the preparation of pharmaceutical compositions for protecting, maintaining, enhancing, or restoring the function or viability of responsive
25 mammalian cells and their associated cells, tissues, and organs. In one particular aspect, the responsive mammalian cells and their associated cells, tissues, or organs are distal to the vasculature by virtue of a tight endothelial cell barrier. In another particular aspect, the cells, tissues, organs or other bodily parts are isolated from a mammalian body, such as those intended for transplant. By way of non-limiting examples, a responsive cell or tissue
30 may be neuronal, retinal, muscle, heart, lung, liver, kidney, small intestine, adrenal cortex, adrenal medulla, capillary endothelial, testes, ovary, pancreas, bone, skin, or endometrial

cells or tissue. Further, non-limiting examples of responsive cells include photoreceptor (rods and cones), ganglion, bipolar, horizontal, amacrine, Müller, Purkinje, myocardium, pace maker, sinoatrial node, sinus node, junction tissue, atrioventricular node, bundle of His, hepatocytes, stellate, Kupffer, mesangial, renal epithelial, tubular interstitial, goblet, intestinal gland (crypts), enteral endocrine, glomerulosa, fasciculate, reticularis, chromaffin, pericyte, Leydig, Sertoli, sperm, Graffian follicle, primordial follicle, islets of Langerhans, α -cells, β -cells, γ -cells, F-cells, osteoprogenitor, osteoclasts, osteoblasts, endometrial stroma, endometrial, stem and endothelial cells. These examples of responsive cells are merely illustrative. In one aspect, the responsive cell or its associated cells, tissues, or organs are not excitable cells, tissues, or organs, nor do they predominantly comprise excitable cells or tissues. In a particular embodiment, the mammalian cell, tissue, or organ for which an aforementioned recombinant tissue protective cytokine is used are those that have expended or will expend a period of time under at least one condition adverse to the viability of the cell, tissue, or organ. In a particular embodiment, the mammalian cell, tissue, or organ for which an aforementioned recombinant tissue protective cytokine is used express the EPO receptor. Such conditions include traumatic *in situ* hypoxia or metabolic dysfunction, surgically-induced *in situ* hypoxia or metabolic dysfunction, or *in situ* toxin exposure, the latter may be associated with chemotherapy or radiation therapy. In one embodiment, the adverse conditions are a result of cardio-pulmonary bypass (heart-lung machine), as is used for certain surgical procedures.

The recombinant tissue protective cytokines of the invention are useful for the therapeutic or prophylactic treatment of human diseases of the central nervous system (CNS) or peripheral nervous system which have primarily neurological or psychiatric symptoms, as well as ophthalmic diseases, cardiovascular diseases, cardiopulmonary diseases, respiratory diseases, kidney, urinary and reproductive diseases, gastrointestinal diseases and endocrine and metabolic abnormalities.

The invention is also directed to pharmaceutical compositions comprising particular aforementioned recombinant tissue protective cytokines for administration to a mammalian animal, preferably a human being. Such pharmaceutical compositions may be formulated for oral, intranasal, or parenteral administration, or in the form of a perfusate solution for maintaining the viability of cells, tissues, or organs *ex vivo*.

Recombinant tissue protective cytokines useful for the aforementioned purposes may be a mutein, or genetically-modified erythropoietin, that is, an erythropoietin for which

at least one modification of the amino acid backbone of the native molecule exists. "Mutant protein," "variant protein" or "mutein" mean a protein comprising a mutant amino acid sequence and includes polypeptides which differ from the amino acid sequence of native erythropoietin due to amino acid deletions, substitutions, or both. "Native sequence" refers to an amino acid or nucleic acid sequence which is identical to a wild-type or native form of a gene or protein. Furthermore, in one embodiment, the recombinant tissue protective cytokines of the invention have cellular protective activity, but also have one or more of erythropoietin's effects upon the bone marrow, i.e., increased hematocrit (erythropoiesis), vasoactive action (vasoconstriction/vasodilation), hyperactivation of platelets, increased production of thrombocytes, and pro-coagulant activities. In another embodiment, the recombinant tissue protective cytokines of the invention have cellular protective activity, but does not have one or more of erythropoietin's effects upon the bone marrow, i.e., increased hematocrit (erythropoiesis), vasoactive action (vasoconstriction/vasodilation), hyperactivation of platelets, increased production of thrombocytes, and pro-coagulant activities. Preferably, a cellular protective recombinant tissue protective cytokine of the invention lacks at least one of erythropoietin's effects on the bone marrow; more preferably the recombinant tissue protective cytokine would lack erythropoietic activity; and most preferably the recombinant tissue protective cytokine lacks all of erythropoietin's effects on the bone marrow.

By way of non-limiting examples, changes in one or more amino acids may be made, or deletions or additions provided, to a native erythropoietin molecule. In a preferred embodiment, the recombinant tissue protective cytokine has one or more modifications in one or more of the following regions: VLQRY (amino acids 11-15 of native, human erythropoietin; SEQ ID NO:1) and/or TKVNFYAW (amino acids 44-51 of native, human erythropoietin; SEQ ID NO:2) and/or SGLRSLTTL (amino acids 100-108 of native, human erythropoietin; SEQ ID NO:3) and/or SNFLRG (amino acids 146-151 of native, human erythropoietin; SEQ ID NO:4). Other mutations may be provided at amino acids 7, 20, 21, 29, 33, 38, 42, 59, 63, 67, 70, 83, 96, 126, 142, 143, 152, 153, 155, 156, and 161 of SEQ ID NO:10. These other mutations may be alone or in addition to at least one mutation in at least one of the regions mentioned above. In certain embodiments, changes in one or more amino acids of TKVNFYAW (amino acids 44-51 of native, human erythropoietin; SEQ ID NO:2) results in a modified erythropoietin molecule with partial function, i.e., having less erythropoietic activity than rhu-EPO. In other embodiments, changes in one or more amino acids of SGLRSLTTL (amino acids 100-108 of native, human erythropoietin; SEQ ID

NO:3) results in a recombinant tissue protective cytokine with partial function, *i.e.*, having less erythropoietic activity than rhu-EPO. The above described recombinant tissue protective cytokines exhibit tissue protective or cellular protective activity. With respect to erythropoietic activities, the above described recombinant tissue protective cytokines lack or exhibit a decrease in one or more erythropoietic activities. Examples of erythropoietic activity include increasing hematocrit, vasoconstriction, hyperactivating platelets, pro-coagulant activities and increasing production of thrombocytes. Erythropoietic activities can be measured by techniques standard in the art. For example, hematocrit can be measured using the UT-7 cell assays described in Section 6.17, or using the techniques described in the Physicians' Desk Reference (Medical Economics Company, Inc., Montvale, NJ, 2000,) which is incorporated by reference herein in its entirety. In particular, pages 519-525 and 2125-2131 disclose methods which can be employed in measuring hematocrit levels and different hematocrit ranges are disclosed that can be used as targets to avoid toxicity. For example, in patients with chronic renal failure, the PDR recommends dosing erythropoietin to achieve non-toxic target hematocrits ranging from 30% to 36% in a patient (e.g., see PDR, p. 523, col. 1, ll. 17-96 and p. 2129, col. 1, ll. 8-93, and accompanying table in cols. 2 and 3). The PDR notes that toxicity in the form of polycythemia (a condition marked by an abnormal increase in the number of circulating red blood cells) can be avoided by carefully monitoring the hematocrit and adjusting doses of EPO, withholding erythropoietin if the hematocrit approaches the high-end of the target range (36% for this patient population) or increases by more than 4 points in any 2-week period, until the hematocrit returns to the suggested target range (30% to 36% for this patient population; see PDR, p. 523, col. 1, and p. 2129, col. 1, under "Dose Adjustment"). In contrast, for cancer patients on chemotherapy, the PDR teaches to adjust the dosage at a different hematocrit level, *i.e.*, if the hematocrit exceeds 40% (see p. 2129, col. 2, under "Dose Adjustment"). In one embodiment, the recombinant tissue protective cytokine has one or more erythropoietic activities, but at levels that are not sufficient to cause adverse effects, *i.e.* effects that outweigh the therapeutic benefit of the cellular protective activity of a recombinant tissue protective cytokine. In one embodiment, the recombinant tissue protective cytokines that possess one or more erythropoietic activities can still be used in the methods of the invention, provided the levels of erythropoietic activity are measured. In those embodiments where the recombinant tissue protective cytokine possesses one or more erythropoietic activities, the erythropoietic activities can be measured and the dose amount and/or dose regimen of the cytokine can be adjusted to ensure the recombinant tissue protective cytokine is not toxic. In those embodiments where the recombinant tissue

protective cytokine possesses one or more erythropoietic activities, the erythropoietic activities can be measured and the dose amount and/or dose regimen of the cytokine can be adjusted to ensure the recombinant tissue protective cytokine has low toxicity. In one embodiment, the recombinant tissue protective cytokine exhibits a decrease in one or more erythropoietic activities by about 1%, 2%, 4%, 6%, 8%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% in comparison to recombinant Epo.

The invention provides for a recombinant tissue protective cytokine lacking at least one activity selected from the group consisting of increasing hematocrit, vasoconstriction, hyperactivating platelets, pro-coagulant activities and increasing production of thrombocytes. The cytokine comprises at least one responsive cellular protective activity selected from the group consisting of protecting, maintaining, enhancing or restoring the function or viability of a responsive mammalian cell, tissue, or organ.

In one embodiment of the invention, the recombinant tissue protective cytokine comprises one or more altered amino acid residue between position 11 to 15 of SEQ ID NO:10 [SEQ ID NO:1], position 44 to 51 of SEQ ID NO 10 [SEQ ID NO:2], position 100-108 of SEQ ID NO [SEQ ID NO:3], or position 146-151 of SEQ ID NO 10 [SEQ ID NO:4].

In another embodiment, the recombinant tissue protective cytokine comprises an altered amino acid residue at one or more of the following positions of SEQ ID NO: 10: 7, 20, 21, 29, 33, 38, 42, 59, 63, 67, 70, 83, 96, 126, 142, 143, 152, 153, 155, 156, or 161.

In yet another embodiment, the recombinant tissue protective cytokine comprises the amino acid sequence of SEQ ID NO: 10 with one or more of the following changes (each altered sequence has been assigned a separate sequence identification number): an alanine at residue 6 of SEQ ID NO: 10 (SEQ ID NO: 15); an alanine at residue 7 of SEQ ID NO: 10 (SEQ ID NO: 16); a serine at residue 7 of SEQ ID NO: 10 (SEQ ID NO: 17); an isoleucine at residue 10 of SEQ ID NO: 10 (SEQ ID NO: 18); a serine at residue 11 of SEQ ID NO: 10 (SEQ ID NO: 19); an alanine at residue 12 of SEQ ID NO: 10 (SEQ ID NO: 20); an alanine at residue 13 of SEQ ID NO: 10 (SEQ ID NO: 21); an alanine residue 14 of SEQ ID NO: 10 (SEQ ID NO: 22); a glutamic acid at residue 14 of SEQ ID NO: 10 (SEQ ID NO: 23); a glutamine at residue 14 of SEQ ID NO: 10 (SEQ ID NO: 24); an alanine at residue 15 of SEQ ID NO: 10 (SEQ ID NO: 25); a phenylalanine at residue 15 of SEQ ID NO: 10 (SEQ ID NO: 26); an isoleucine at residue 15 of SEQ ID NO: 10 (SEQ ID NO: 27); a

glutamic acid at residue 20 of SEQ ID NO: 10 (SEQ ID NO: 28); an alanine at residue 20 of
SEQ ID NO: 10 (SEQ ID NO: 29); an alanine at residue 21 of SEQ ID NO: 10 (SEQ ID
NO: 30); a lysine at residue 24 of SEQ ID NO: 10 (SEQ ID NO: 31); a serine at residue 29
of SEQ ID NO: 10 (SEQ ID NO: 32); a tyrosine at residue 29 of SEQ ID NO: 10 (SEQ ID
5 NO: 33); an asparagine at residue 30 of SEQ ID NO: 10 (SEQ ID NO: 34); a threonine at
residue 32 of SEQ ID NO: 10 (SEQ ID NO: 35); a serine at residue 33 of SEQ ID NO: 10
(SEQ ID NO: 36); a tyrosine at residue 33 of SEQ ID NO: 10 (SEQ ID NO: 37); a lysine at
residue 38 of SEQ ID NO: 10 (SEQ ID NO: 38); a lysine at residue 83 of SEQ ID NO: 10
(SEQ ID NO: 39); an asparagine at residue 42 of SEQ ID NO: 10 (SEQ ID NO: 40); an
10 alanine at residue 42 of SEQ ID NO: 10 (SEQ ID NO: 41); an alanine at residue 43 of SEQ
ID NO: 10 (SEQ ID NO: 42); an isoleucine at residue 44 of SEQ ID NO: 10 (SEQ ID NO:
43); an aspartic acid at residue 45 of SEQ ID NO: 10 (SEQ ID NO: 44); an alanine at
residue 45 of SEQ ID NO: 10 (SEQ ID NO: 45); an alanine at residue 46 of SEQ ID NO: 10
(SEQ ID NO: 46); an alanine at residue 47 of SEQ ID NO: 10 (SEQ ID NO: 47); an
15 isoleucine at residue 48 of SEQ ID NO: 10 (SEQ ID NO: 48); an alanine at residue 48 of
SEQ ID NO: 10 (SEQ ID NO: 49); an alanine at residue 49 of SEQ ID NO: 10 (SEQ ID
NO: 50); a serine at residue 49 of SEQ ID NO: 10 (SEQ ID NO: 51); a phenylalanine at
residue 51 of SEQ ID NO: 10 (SEQ ID NO: 52); an asparagine at residue 51 of SEQ ID
NO: 10 (SEQ ID NO: 53); an alanine at residue 52 of SEQ ID NO: 10 (SEQ ID NO: 54); an
20 asparagine at residue 59 of SEQ ID NO: 10 (SEQ ID NO: 55); a threonine at residue 62 of
SEQ ID NO: 10 (SEQ ID NO: 56); a serine at residue 67 of SEQ ID NO: 10 (SEQ ID NO:
57); an alanine at residue 70 of SEQ ID NO: 10 (SEQ ID NO: 58); an arginine at residue 96
of SEQ ID NO: 10 (SEQ ID NO: 59); an alanine at residue 97 of SEQ ID NO: 10 (SEQ ID
NO: 60); an arginine at residue 100 of SEQ ID NO: 10 (SEQ ID NO: 61); a glutamic acid at
25 residue 100 of SEQ ID NO: 10 (SEQ ID NO: 62); an alanine at residue 100 of SEQ ID NO:
10 (SEQ ID NO: 63); a threonine at residue 100 of SEQ ID NO: 10 (SEQ ID NO: 64); an
alanine at residue 101 of SEQ ID NO: 10 (SEQ ID NO: 65); an isoleucine at residue 101 of
SEQ ID NO: 10 (SEQ ID NO: 66); an alanine at residue 102 of SEQ ID NO: 10 (SEQ ID
NO: 67); an alanine at residue 103 of SEQ ID NO: 10 (SEQ ID NO: 68); a glutamic acid at
30 residue 103 of SEQ ID NO: 10 (SEQ ID NO: 69); an alanine at residue 104 of SEQ ID NO:
10 (SEQ ID NO: 70); an isoleucine at residue 104 of SEQ ID NO: 10 (SEQ ID NO: 71); an
alanine at residue 105 of SEQ ID NO: 10 (SEQ ID NO: 72); an alanine at residue 106 of
SEQ ID NO: 10 (SEQ ID NO: 73); an isoleucine at residue 106 of SEQ ID NO: 10 (SEQ ID
NO: 74); an alanine at residue 107 of SEQ ID NO: 10 (SEQ ID NO: 75); a leucine at
35 residue 107 of SEQ ID NO: 10 (SEQ ID NO: 76); a lysine at residue 108 of SEQ ID NO: 10

(SEQ ID NO: 77); an alanine at residue 108 of SEQ ID NO: 10 (SEQ ID NO: 78); a serine at residue 108 of SEQ ID NO: 10 (SEQ ID NO: 79); an alanine at residue 116 of SEQ ID NO: 10 (SEQ ID NO: 80); an alanine at residue 126 of SEQ ID NO: 10 (SEQ ID NO: 81); an alanine at residue 132 of SEQ ID NO: 10 (SEQ ID NO: 82); an alanine at residue 133 of SEQ ID NO: 10 (SEQ ID NO: 83); an alanine at residue 134 of SEQ ID NO: 10 (SEQ ID NO: 84); an alanine at residue 140 of SEQ ID NO: 10 (SEQ ID NO: 85); an isoleucine at residue 142 of SEQ ID NO: 10 (SEQ ID NO: 86); an alanine at residue 143 of SEQ ID NO: 10 (SEQ ID NO: 87); an alanine at residue 146 of SEQ ID NO: 10 (SEQ ID NO: 88); a lysine at residue 147 of SEQ ID NO: 10 (SEQ ID NO: 89); an alanine at residue 147 of SEQ ID NO: 10 (SEQ ID NO: 90); a tyrosine at residue 148 of SEQ ID NO: 10 (SEQ ID NO: 91); an alanine at residue 148 of SEQ ID NO: 10 (SEQ ID NO: 92); an alanine at residue 149 of SEQ ID NO: 10 (SEQ ID NO: 93); an alanine at residue 150 of SEQ ID NO: 10 (SEQ ID NO: 94); a glutamic acid at residue 150 of SEQ ID NO: 10 (SEQ ID NO: 95); an alanine at residue 151 of SEQ ID NO: 10 (SEQ ID NO: 96); an alanine at residue 152 of SEQ ID NO: 10 (SEQ ID NO: 97); a tryptophan at residue 152 of SEQ ID NO: 10 (SEQ ID NO: 98); an alanine at residue 153 of SEQ ID NO: 10 (SEQ ID NO: 99); an alanine at residue 154 of SEQ ID NO: 10 (SEQ ID NO: 100); an alanine at residue 155 of SEQ ID NO: 10 (SEQ ID NO: 101); an alanine at residue 158 of SEQ ID NO: 10 (SEQ ID NO: 102); a serine at residue 160 of SEQ ID NO: 10 (SEQ ID NO: 103); an alanine at residue 161 of SEQ ID NO: 10 (SEQ ID NO: 104); or an alanine at residue 162 of SEQ ID NO: 10 (SEQ ID NO: 105). In one embodiment, the recombinant tissue protective cytokine comprises the amino acid sequence of SEQ ID NO: 10 with one or more of the amino acid residue substitutions of SEQ ID NOs: 15-105 and 119.

In yet another embodiment, the recombinant tissue protective cytokine comprises the amino acid sequence of SEQ ID NO: 10 with a deletion of amino acid residues 44-49 of SEQ ID NO: 10.

In still another embodiment, the recombinant tissue protective cytokine comprises, the amino acid sequence of SEQ ID NO: 10 with at least one of the following changes (each altered sequence has been assigned a separate sequence identification number): i) an aspartic acid at residue 45, and a glutamic acid at residue 100 of SEQ ID NO: 10 (SEQ ID NO: 106); ii) an asparagine at residue 30, a threonine at residue 32 of SEQ ID NO: 10 (SEQ ID NO: 107); iii) an aspartic acid at residue 45, a glutamic acid at residue 150 SEQ ID NO: 10 (SEQ ID NO: 108); iv) a glutamic acid at residue 103, and a serine at residue 108 of

SEQ ID NO: 10 (SEQ ID NO: 109); v) an alanine at residue 140 and an alanine at residue 52 of SEQ ID NO: 10 (SEQ ID NO: 110); vi) an alanine at residue 140, an alanine at residue 52, an alanine at residue 45 of SEQ ID NO: 10 (SEQ ID NO: 111); vii) an alanine at residue 97, and an alanine at residue 152 of SEQ ID NO: 10 (SEQ ID NO: 112); iix) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45 of SEQ ID NO: 10 (SEQ ID NO: 113); ix) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45, and an alanine at residue 52 of SEQ ID NO: 10 (SEQ ID NO: 114); x) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45, an alanine at residue 52, and an alanine at residue 140 of SEQ ID NO: 10 (SEQ ID NO: 115); xi) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45, an alanine at residue 52, an alanine at residue 140, an alanine at residue 154, a lysine at residue 24, a lysine at residue 38, a lysine at residue 83, a lysine at residue 24 and an alanine at residue 15 of SEQ ID NO: 10 (SEQ ID NO: 116); xii) a lysine at residue 24, a lysine at residue 38, and a lysine at residue 83 SEQ ID NO: 10 (SEQ ID NO: 117); or xiv) a lysine at residue 24 and an alanine at residue 15 SEQ ID NO: 10 (SEQ ID NO: 118). In one embodiment, the recombinant tissue protective cytokine comprises, the amino acid sequence of SEQ ID NO: 10 with at least one of the following amino acid residue substitutions of SEQ ID NOs: 106-118.

One embodiment of the invention is directed to the recombinant tissue protective cytokine as described herein above, further comprising a chemical modification of one or more amino acids. In another embodiment the chemical modification comprises altering the charge of the recombinant tissue protective cytokine. In yet another embodiment, a positive or negative charge is chemically added to an amino acid residue where a charged amino acid residue is modified to an uncharged residue.

Moreover, such aforementioned recombinant tissue protective cytokines may be further modified by having a chemical modification of one or more amino acids, such as described in the following co-pending applications: PCT application serial no. PCT/US01/49479, filed December 28, 2001, U.S. Patent Application Serial No. 09/753,132 filed December 29, 2000, and U.S. Patent Application Attorney Docket No. KW00-009C02-US filed July 3, 2002, each of these applications is incorporated herein by reference in their entirety. These further chemical modifications may be used to enhance the tissue protective activities of the recombinant tissue protective cytokines or suppress any effects the recombinant tissue protective cytokines may have on bone marrow. In a further embodiment, the additional chemical modification is provided to restore solubility of the

molecule that may be reduced as a result of the aforementioned genetic modification, such as chemically adding a positive or negative charge to the molecule if a charged amino acid residue is changed to an uncharged residue.

By way of non-limiting examples, recombinant tissue protective cytokines of the invention include human erythropoietin mutein S100E (SEQ ID NO:5), human erythropoietin mutein K45D (SEQ ID NO:6), and any of the nonerythropoietic yet cellular protective recombinant tissue protective cytokines or those able to benefit a responsive cell, tissue or organ, that are described in Elliott *et al.*, 1997, Blood 89:493-502; Boissel *et al.*, Journal of Biological Chemistry, vol. 268, No. 21, pp. 15983-15993 (1993); Wen *et al.*, Journal of Biological Chemistry, vol. 269, No. 36, pp. 22839-22846 (1994); and Syed *et al.*, Nature, vol. 395, pp. 511-516 (1998), which are incorporated herein by reference in their entireties. The present invention is directed to methods for the use of any of the aforementioned recombinant tissue protective cytokines for the protection, restoration, and enhancement of responsive cells, tissues, and organs.

Other recombinant tissue protective cytokines of the invention include an aforementioned erythropoietin comprising at least one genetically altered amino acid with at least one additional modification which may be another modification of at least one additional amino acid of the erythropoietin molecule, or a modification of at least one carbohydrate of the erythropoietin molecule. The genetically altered amino acid(s) may be the one or among those further modified. Of course, recombinant tissue protective cytokine molecules useful for the purposes herein may have a plurality of modifications as compared to the native erythropoietin molecule, such as multiple modifications of the amino acid portion of the molecule, multiple modifications of the carbohydrate portion of the molecule, or at least a second modification of the amino acid portion of the molecule and at least one modification of the carbohydrate portion of the molecule. The recombinant tissue protective cytokine molecule retains its ability of protecting, maintaining, enhancing or restoring the function or viability of responsive mammalian cells, yet other properties of the recombinant tissue protective cytokine unrelated to the aforementioned, desirable feature may be absent as compared to the native molecule. In a preferred embodiment, the recombinant tissue protective cytokine is non-erythropoietic.

In another embodiment, the recombinant tissue protective cytokines can be modified by fucosylation to alter glycosylation patterns on a glycoprotein.

One embodiment of the invention is directed to the recombinant tissue protective cytokine as described herein above is a human erythropoietin mutein. In another embodiment of the invention the recombinant tissue protective cytokine is a human phenylglyoxal erythropoietin mutein. In another embodiment of the invention, the recombinant tissue protective cytokine is a human asialoerythropoietin mutein.

In one embodiment, as described herein above, the recombinant tissue protective cytokine comprises at least one responsive cellular protective activity selected from the group consisting of protecting, maintaining, enhancing or restoring the function or viability of a responsive mammalian cell, tissue, or organ. In such an embodiment, the responsive mammalian cell comprises a neuronal, muscle, heart, lung, liver, kidney, small intestine, adrenal cortex, adrenal medulla, capillary, endothelial, testes, ovary, endometrial, or stem cell. In other embodiments, the cell comprises a photoreceptor, ganglion, bipolar, horizontal, amacrine, Müller, myocardium, pace maker, sinoatrial node, sinus node, atrioventricular node, bundle of His, hepatocyte, stellate, Kupffer, mesangial, goblet, intestinal gland, enteral endocrine, glomerulosa, fasciculate, reticularis, chromaffin, pericyte, Leydig, Sertoli, sperm, Graffian follicles, primordial follicles, endometrial stroma cells, or endometrial cell.

According to another aspect of the invention, the recombinant tissue protective cytokine, as described herein above, is capable of traversing an endothelial cell barrier. In a related embodiment, the endothelial cell barrier comprises the blood-brain barrier, the blood-eye barrier, the blood testis barrier, the blood-ovary barrier, blood-placenta, blood-heart, blood-kidney, and the blood-uterus barrier.

In another embodiment of the invention, the recombinant tissue protective cytokine as described herein above is further modified. In one embodiment, the recombinant tissue protective cytokine is selected from the group consisting of: i) a cytokine having a reduced number or no sialic acid moieties; ii) a cytokine having a reduced number or no N-linked or O-linked carbohydrates; iii) a cytokine having at least a reduced carbohydrate content by virtue of treatment of native cytokine with at least one glycosidase; iv) a cytokine having at least one or more oxidized carbohydrates; v) a cytokine having at least one or more oxidized carbohydrates and is chemically reduced; vi) a cytokine having at least one or more modified arginine residues; vii) a cytokine having at least one or more modified lysine residues or a modification of the N-terminal amino group of a cytokine molecule; viii) a cytokine having at least a modified tyrosine residue; ix) a cytokine having at least a

modified aspartic acid or glutamic acid residue; x) a cytokine having at a modified tryptophan residue; xi) a cytokine having at least one amino acid group removed; xii) a cytokine having at least one opening of at least one of the cystine linkages in the cytokine molecule; xiii) a truncated cytokine; xiv) a cytokine having at least one polyethylene glycol molecule attached; xv) a cytokine having at least one fatty acid attached; xvi) a cytokine having a non-mammalian glycosylation pattern by virtue of the expression of a recombinant cytokine in non-mammalian cells; and xvi) a cytokine having at least one histidine tagged amino acid to facilitate purification.

In one embodiment, the recombinant tissue protective cytokine of the invention has a reduced number of sialic acid moieties, or no sialic acid moieties. In a preferred embodiment, the recombinant tissue protective cytokine is the asialo form of an erythropoietin (i.e. has no sialic acid moieties), and most preferably, a human asialoerythropoietin. In another embodiment, the recombinant tissue protective cytokine has 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, or 13 sialic acid moieties. The number of available sites for sialylation may be altered by the presence of one or more altered or modified amino acids in the recombinant tissue protective cytokine. Therefore, the present invention covers embodiments wherein the recombinant tissue protective cytokine is either hyposialylated or hypersialylated. In a preferred aspect, the erythropoietin mutein has more than the 14 sialic acid moieties present in native erythropoietin.

In one embodiment, the recombinant tissue protective cytokine is an erythropoietin with no N-linked carbohydrates. In another embodiment, the recombinant tissue protective cytokine is an erythropoietin with a reduced number of N-linked carbohydrates. In one embodiment, the recombinant tissue protective cytokine is an erythropoietin with no O-linked carbohydrates. In another embodiment, the recombinant tissue protective cytokine is an erythropoietin with a reduced number of O-linked carbohydrates.

In another embodiment, the recombinant tissue protective cytokines can be modified by fucosylation to alter glycosylation patterns on a glycoprotein.

In yet another embodiment, the recombinant tissue protective cytokine is treated with at least one glycosidase. In another embodiment, the recombinant tissue protective cytokine has at least a reduced carbohydrate content by virtue of treatment of the recombinant tissue protective cytokine with at least one glycosidase.

In yet another embodiment, the carbohydrate portion of the recombinant tissue protective cytokine has at least a non-mammalian glycosylation pattern by virtue of the expression of a recombinant erythropoietin in non-mammalian cells. In preferred embodiments, the recombinant tissue protective cytokines are expressed in insect cells,
5 plant cells, bacteria cells, or yeast cells.

In yet another embodiment, the recombinant tissue protective cytokine further has at least one or more oxidized carbohydrates which also may be chemically reduced. In a preferred embodiment, the recombinant tissue protective cytokine is periodate-oxidized erythropoietin. In certain embodiments, the periodate-oxidized erythropoietin is
10 chemically reduced with sodium cyanoborohydride.

In yet another embodiment, the recombinant tissue protective cytokine for the aforementioned uses has at least one or more modified arginine residues. In one embodiment, the recombinant tissue protective cytokine comprises an R-glyoxal moiety on the one or more arginine residues, wherein R is aryl or alkyl moiety. In yet another
15 embodiment, the recombinant tissue protective cytokine is phenylglyoxal-erythropoietin. In yet another embodiment, the recombinant tissue protective cytokine is an erythropoietin in which an arginine residue is modified by reaction with a vicinal diketone, such as but not limited to, 2,3-butanedione and cyclohexanedione. In yet another embodiment, the recombinant tissue protective cytokine is an erythropoietin in which an arginine residue is
20 reacted with 3-deoxyglucosone.

In yet another embodiment, the recombinant tissue protective cytokine comprises at least one or more modified lysine residues or a modification of the N-terminal amino group of the erythropoietin molecule, such modifications as those resulting from reaction of the lysine residue or N-terminal amino group with an amino-group-modifying agent. The
25 modified lysine residue further may be chemically reduced. In one preferred embodiment, a recombinant tissue protective cytokine is biotinylated or carbamylated or acylated, such as acetylated, via one or more lysine groups. In another preferred embodiment, the lysine is reacted with an aldehyde or reducing sugar to form an imine, which may be stabilized by reduction as with sodium cyanoborohydride to form an N-alkylated lysine such as glucitolyl
30 lysine, or which in the case of reducing sugars may be stabilized by Amadori or Heyns rearrangement to form an alpha-deoxy alpha-amino sugar such as alpha-deoxy-alpha-fructosyllysine. In another preferred embodiment, the lysine group is carbamylated (carbamoylated), such as by virtue of reaction with cyanate ion, alkyl-carbamylated, aryl-

carbamylated, or aryl-thiocarbamylated with an alkyl-isocyanate, aryl-isocyanate, or aryl isothiocyanate, respectively, or it may be acylated by a reactive alkylcarboxylic or arylcarboxylic acid derivative, such as by reaction with acetic anhydride, succinic anhydride or phthalic anhydride. At least one lysine group may also be trinitrophenyl, modified by

5 reaction with a trinitrobenzenesulfonic acid, or preferably its salts. In another embodiment, lysine residues may be modified by reaction with a glyoxal derivative, such as reaction with glyoxal, methylglyoxal or 3-deoxyglucosone, to form the corresponding alpha-carboxyalkyl derivatives. In a related embodiment, the carbamylated cytokine is comprised of alpha-N-carbamoylerythropoietin; N-epsilon-carbamoylerythropoietin; alpha-N-carbamoyl, N-

10 epsilon-carbamoylerythropoietin; alpha-N-carbamoylasialoerythropoietin; N-epsilon-carbamoylasialoerythropoietin; alpha-N-carbamoyl, N-epsilon-carbamoylasialoerythropoietin; alpha-N-carbamoylhyposialoerythropoietin; N-epsilon-carbamoylhyposialoerythropoietin; and alpha-N-carbamoyl, N-epsilon-carbamoylhyposialoerythropoietin. In yet another embodiment, the recombinant tissue

15 protective cytokine comprises at least one acylated lysine residue. In yet another embodiment, the recombinant tissue protective cytokine comprises at least one acylated lysine residue. In yet another embodiment, the recombinant tissue protective cytokine comprises at least one acylated lysine residue. In a related embodiment, the acetylated cytokine is comprised of alpha-N-acetylerythropoietin; N-epsilon-acetylerythropoietin;

20 alpha-N-acetyl, N-epsilon-acetylerythropoietin; alpha-N-acetylasialoerythropoietin; N-epsilon-acetylasialoerythropoietin; alpha-N-acetyl, N-epsilon-acetylasialoerythropoietin; alpha-N-acetylhyposialoerythropoietin; N-epsilon-acetylhyposialoerythropoietin; alpha-N-acetyl, N-epsilon-acetylhyposialoerythropoietin; alpha-N-acetylhypersialoerythropoietin; N-epsilon-acetylhypersialoerythropoietin; alpha-N-acetyl, and N-epsilon-

25 acetylhypersialoerythropoietin.

In yet another embodiment, the recombinant tissue protective cytokine has a lysine residue that is succinylated. In a related embodiment, the succinylated cytokine is comprised of alpha-N-succinylerythropoietin; N-epsilon-succinylerythropoietin; alpha-N-succinyl, N-epsilon-succinylerythropoietin; alpha-N-succinylasialoerythropoietin; N-

30 epsilon-succinylasialoerythropoietin; alpha-N-succinyl, N-epsilon-succinylasialoerythropoietin; alpha-N-succinylhyposialoerythropoietin; N-epsilon-succinylhyposialoerythropoietin; alpha-N-succinyl, N-epsilon-succinylhyposialoerythropoietin; alpha-N-succinylhypersialoerythropoietin; N-epsilon-succinylhypersialoerythropoietin; and N-epsilon-succinylhypersialoerythropoietin.

In one embodiment, at least one tyrosine residue of a recombinant tissue protective cytokine may be modified in an aromatic ring position by an electrophilic reagent, such as by nitration or iodination. In a related embodiment, the recombinant tissue protective cytokine as described herein above comprises at least one lysine residue modified by 2, 4, 6 trinitrobenzenesulfonate sodium or another salt thereof.

In another embodiment, the recombinant tissue protective cytokine comprises at least one nitrated or iodinated tyrosine residue.

In another embodiment, the recombinant tissue protective cytokine comprises an aspartic acid or glutamic acid residue that is reacted with a carbodiimide followed by reaction with an amine. In a related embodiment, the amine is glycine.

In one embodiment, at least a tryptophan residue of a recombinant tissue protective cytokine is modified, such as by reaction with n-bromosuccinimide or n-chlorosuccinimide.

In another embodiment, a recombinant tissue protective cytokine is provided having at least one erythropoietin amino group removed, such as by reaction with ninhydrin followed by reduction of the resulting carbonyl group by reaction with borohydride.

In yet another embodiment, a recombinant tissue protective cytokine is provided having at least an opening of at least one of the cystine linkages in the molecule by reaction with a reducing agent such as dithiothreitol, followed by reaction of the subsequent sulfhydryls with iodoacetamide, iodoacetic acid or another electrophile to prevent reformation of the disulfide linkages.

In yet another embodiment, a recombinant tissue protective cytokine is subjected to a limited chemical proteolysis that targets specific residues, for example, to cleave after tryptophan residues. Such resulting recombinant tissue protective cytokine fragments are embraced herein.

As noted above, a recombinant tissue protective cytokine useful for the purposes herein optionally may have at least one of the aforementioned chemical modifications in addition to the genetically altered amino acid(s), but may have more than one of the above modifications. By way of example of a recombinant tissue protective cytokine with one modification to the carbohydrate portion of the molecule and one modification to the amino acid portion, a recombinant tissue protective cytokine is an asialoerythropoietin that has its lysine residues biotinylated, acylated (such as acetylated) or carbamylated. The

recombinant tissue protective cytokines can also be modified by the addition of fatty acid chains. In another embodiment, a recombinant tissue protective cytokine is modified by pegylation, to create pegylated tissue protective cytokines by the addition of polyethylene glycol (PEG).

5 According to one aspect of the invention, there is provided an isolated nucleic acid molecule that comprises a nucleotide sequence which encodes a polypeptide comprising the recombinant tissue protective cytokine as described herein above. In one embodiment, the isolated nucleic acid molecule comprises the nucleotide sequence of nucleotide residues 5461 through 6041 of the vector construct of SEQ ID NO: 208, nucleotide residues 5461
10 through 6041 of SEQ ID NO: 209, nucleotide residues 5461 through 6041 of SEQ ID NO: 210, nucleotide residues 5461 through 6041 of SEQ ID NO: 211, or nucleotide residues 5461 through 6041 of SEQ ID NO: 212.

 In one embodiment of the invention, there is provided an isolated nucleic acid molecule that comprises a nucleotide sequence (i.e., a cDNA, a nucleotide sequence
15 interrupted by introns, or uninterrupted by introns), which encodes a polypeptide comprising or consisting of the recombinant tissue protective cytokine as described herein above with the proviso that the nucleic acid molecule does not encode a recombinant tissue protective cytokine that comprises one or more of the following amino acid substitutions: I6A, C7A, K20A, P42A, D43A, K45D, K45A, F48A, Y49A, K52A, K49A, S100E, R103A,
20 K116A, T132A, I133A, K140A, N147K, N147A, R150A, R150E, G151A, K152A, K154A, G158A, C161A, or R162A. In a related embodiment, there is provided an isolated nucleic acid molecule that comprises a nucleotide sequence which encodes a polypeptide comprising the recombinant tissue protective cytokine as described herein above with the proviso that the nucleic acid molecule does not encode a recombinant tissue protective
25 cytokine that comprises any of the following combinations of substitutions: N24K/N38K/N83K or A30N/H32T. In one embodiment, the a nucleotide sequence, encoding the recombinant tissue protective cytokine, is synthesized using preferred codons that facilitate optimal expression in a particular host cell. Such preferred codons can be optimal for expression in cells of a species of plant, bacteria, yeast, mammal, fungi, or
30 insect.

 The invention also provides for a vector comprising the nucleic acid molecule. The invention also provides for an expression vector comprising the nucleic acid molecule and at least one regulatory region operably linked to the nucleic acid molecule. In one

embodiment, the vector is a pCiNeo vector. In another embodiment, the invention provides for a cell comprising the expression vector. In yet another embodiment, there is provided a genetically-engineered cell which comprises the nucleic acid molecule.

In another embodiment, the present invention also embraces compositions, including
5 pharmaceutical compositions, comprising one or more of the aforementioned recombinant tissue protective cytokines.

According to another aspect of the invention, there is provided a pharmaceutical composition comprising a recombinant tissue protective cytokine as described herein above, lacking at least one erythropoietic activity selected from the group consisting of increasing
10 hematocrit, vasoconstriction, hyperactivating platelets, pro-coagulant activities and increasing production of thrombocytes. According to another aspect of the invention, there is provided a pharmaceutical composition comprising a recombinant tissue protective cytokine as described herein above, but the cytokines do not lack at least one erythropoietic activity selected from the group consisting of increasing hematocrit, vasoconstriction,
15 hyperactivating platelets, pro-coagulant activities and increasing production of thrombocytes. The cytokine comprises at least one responsive cellular protective activity selected from the group consisting of protecting, maintaining, enhancing or restoring the function or viability of a responsive mammalian cell, tissue or organ. The recombinant tissue protective cytokine of the pharmaceutical composition may comprise the amino acid
20 sequence of SEQ ID NO: 10 with at least one of the following changes, i.e. substitutions, (each change or combination of changes listed has been assigned a separate sequence identification number): i) an aspartic acid at residue 45, and a glutamic acid at residue 100 of SEQ ID NO: 10 (SEQ ID NO: 106); ii) an asparagine at residue 30, a threonine at residue 32 of SEQ ID NO: 10 (SEQ ID NO: 107); iii) an aspartic acid at residue 45, a glutamic acid
25 at residue 150 SEQ ID NO: 10 (SEQ ID NO: 108); iv) a glutamic acid at residue 103, and a serine at residue 108 of SEQ ID NO: 10 (SEQ ID NO: 109); v) an alanine at residue 140 and an alanine at residue 52 of SEQ ID NO: 10 (SEQ ID NO: 110); vi) an alanine at residue 140, an alanine at residue 52, an alanine at residue 45 of SEQ ID NO: 10 (SEQ ID NO: 111); vii) an alanine at residue 97, and an alanine at residue 152 of SEQ ID NO: 10 (SEQ
30 ID NO: 112); iix) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45 of SEQ ID NO: 10 (SEQ ID NO: 113); ix) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45, and an alanine at residue 52 of SEQ ID NO: 10 (SEQ ID NO: 114); x) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45, an

alanine at residue 52, and an alanine at residue 140 of SEQ ID NO: 10 (SEQ ID NO: 115);
xi) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45, an alanine at
residue 52, an alanine at residue 140, an alanine at residue 154, a lysine at residue 24, a
lysine at residue 38, a lysine at residue 83, a lysine at residue 24 and an alanine at residue 15
5 of SEQ ID NO: 10 (SEQ ID NO: 116); xii) a lysine at residue 24, a lysine at residue 38, and
a lysine at residue 83 SEQ ID NO: 10 (SEQ ID NO: 117); or xiv) a lysine at residue 24 and
an alanine at residue 15 SEQ ID NO: 10 (SEQ ID NO: 118).

According to another aspect of the invention, there is provided a pharmaceutical
composition for protecting, maintaining, enhancing, or restoring the function or viability of
10 responsive mammalian cells and their associated cells, tissues, and organs, comprising a
therapeutically effective amount of a recombinant tissue protective cytokine, comprising at
least one of the following amino acid residue substitutions: (each change or combination of
changes listed has been assigned a separate sequence identification number): a tryptophan at
residue 152 of SEQ ID NO: 10 (SEQ ID NO: 98); an alanine at residue 14 and an alanine at
15 residue 15 of SEQ ID NO: 10 (SEQ ID NO: 119); an alanine at residue 6 of SEQ ID NO: 10
(SEQ ID NO: 15); an alanine at residue 7 of SEQ ID NO: 10 (SEQ ID NO: 16); an alanine
at residue 43 of SEQ ID NO: 10 (SEQ ID NO: 42); an alanine at residue 42 of SEQ ID NO:
10 (SEQ ID NO: 41); an alanine at residue 48 of SEQ ID NO: 10 (SEQ ID NO: 49); an
alanine at residue 49 of SEQ ID NO: 10 (SEQ ID NO: 50); an threonine at residue 32 of
20 SEQ ID NO: 10 (SEQ ID NO: 35); an alanine at residue 133 of SEQ ID NO: 10 (SEQ ID
NO: 83); an alanine at residue 134 of SEQ ID NO: 10 (SEQ ID NO: 84); an alanine at
residue 147 of SEQ ID NO: 10 (SEQ ID NO: 90); an alanine at residue 148 of SEQ ID NO:
10 (SEQ ID NO: 92); an alanine at residue 150 of SEQ ID NO: 10 (SEQ ID NO: 94); an
alanine at residue 151 of SEQ ID NO: 10 (SEQ ID NO: 96); an alanine at residue 158 of
25 SEQ ID NO: 10 (SEQ ID NO: 102); an alanine at residue 161 of SEQ ID NO: 10 (SEQ ID
NO: 104); or an alanine at residue 162 of SEQ ID NO: 10 (SEQ ID NO: 105).

In one embodiment, the pharmaceutical composition described above herein is
formulated for oral, intranasal, or parenteral administration. In another embodiment, the
pharmaceutical composition is formulated as a perfusate solution.

30 In certain embodiments, the pharmaceutical compositions of the invention for
protecting, maintaining, enhancing, or restoring the function or viability of responsive
mammalian cells and their associated cells, tissues, and organs, comprise a therapeutically

effective amount of a recombinant tissue protective cytokine, comprising at least one substitution of amino acid residues of native, human erythropoietin amino acid sequence.

5 In other embodiments, a pharmaceutical composition of the invention for protecting, maintaining, enhancing, or restoring the function or viability of responsive mammalian cells and their associated cells, tissues, and organs, comprises a therapeutically effective amount of a recombinant tissue protective cytokine, comprising cellular protective activity may lack one or more erythropoietic activities or effects such as increasing hematocrit, vasoactive action (vasoconstriction/vasodilatation), hyperactivating platelets, pro-coagulant activities and increasing production of thrombocytes.

10 In other embodiments, a pharmaceutical composition of the invention for protecting, maintaining, enhancing, or restoring the function or viability of responsive mammalian cells and their associated cells, tissues, and organs, comprises a therapeutically effective amount of a recombinant tissue protective cytokine, comprising cellular protective activity also has one or more erythropoietic activities or effects such as increasing hematocrit, vasoactive
15 action (vasoconstriction/vasodilatation), hyperactivating platelets, pro-coagulant activities and increasing production of thrombocytes.

According to one aspect of the invention, there is provided a method for protecting, maintaining or enhancing the viability of a cell, tissue, or organ isolated from a mammalian body comprising exposing said cell, tissue, or organ to a pharmaceutical composition
20 comprising a recombinant tissue protective cytokine comprised of an erythropoietin that lacks at least one erythropoietic activity selected from the group consisting of increasing hematocrit, vasoactive action (vasoconstriction/vasodilatation), hyperactivating platelets, pro-coagulant activity and increasing production of thrombocytes. In certain embodiments, the protection does not effect bone marrow.

25 The invention also provides for a method for protecting, maintaining or enhancing the viability of a cell, tissue, or organ isolated from a mammalian body comprising exposing said cell, tissue, or organ to a pharmaceutical composition comprising a recombinant tissue protective cytokine comprised, as described herein above, that lacks at least one erythropoietic activity selected from the group consisting of increasing hematocrit,
30 vasoactive action (vasoconstriction/vasodilatation), hyperactivating platelets, pro-coagulant activity and increasing production of thrombocytes.

The invention further provides for the use of a recombinant tissue protective cytokine as described herein above, that lacks at least one erythropoietic activity selected from the group consisting of increasing hematocrit, vasoactive action (vasoconstriction/vasodilatation), hyperactivating platelets, pro-coagulant activity and increasing production of thrombocytes, for the preparation of a pharmaceutical composition for the protection against and prevention of a tissue injury as well as the restoration of and rejuvenation of tissue and tissue function in a mammal. In one embodiment, the injury is caused by a seizure disorder, multiple sclerosis, stroke, hypotension, cardiac arrest, ischemia, myocardial infarction, inflammation, age-related loss of cognitive function, radiation damage, cerebral palsy, neurodegenerative disease, Alzheimer's disease, Parkinson's disease, Leigh's disease, AIDS dementia, memory loss, amyotrophic lateral sclerosis, alcoholism, mood disorder, anxiety disorder, attention deficit disorder, hyperactivity, autism, Creutzfeld-Jakob disease, brain or spinal cord trauma or ischemia, heart-lung bypass, chronic heart failure, macular degeneration, diabetic neuropathy, diabetic retinopathy, glaucoma, retinal ischemia, or retinal trauma.

According to another aspect of the invention, there is provided a method for facilitating the transcytosis of a molecule across an endothelial cell barrier in a mammal comprising administration to said mammal a composition comprising said molecule in association with a recombinant tissue protective cytokine as described herein above, lacking at least one activity selected from the group consisting of increasing hematocrit, increasing blood pressure, hyperactivating platelets, and increasing production of thrombocytes. In one embodiment, the association is a labile covalent bond, a stable covalent bond, or a non-covalent association with a binding site for said molecule. According to another aspect of the invention, there is provided a method for facilitating the transcytosis of a molecule across an endothelial cell barrier in a mammal comprising administration to said mammal a composition comprising said molecule in association with a recombinant tissue protective cytokine as described herein above, and having activity selected from the group consisting of increasing hematocrit, increasing blood pressure, hyperactivating platelets, and increasing production of thrombocytes. In one embodiment, the association is a labile covalent bond, a stable covalent bond, or a non-covalent association with a binding site for said molecule. In another embodiment, the endothelial cell barrier is selected from the group consisting of the blood-brain barrier, the blood-eye barrier, the blood-testis barrier, the blood-ovary barrier, the blood-heart, the blood kidney, and the blood-placenta barrier. In yet another embodiment, the molecule is a receptor agonist or antagonist hormone, a

neurotrophic factor, an antimicrobial agent, an antiviral agent, a radiopharmaceutical, an antisense oligonucleotide, an antibody, an immunosuppressant, a dye, a marker, or an anti-cancer drug.

According to another aspect of the invention, there is provided a composition for
5 transporting a molecule via transcytosis across an endothelial cell barrier comprising said molecule in association with a recombinant tissue protective cytokine, as described herein above, lacking at least one erythropoietic activity selected from the group consisting of increasing hematocrit, vasoactive action (vasoconstriction/vasodilatation), hyperactivating platelets, pro-coagulant activity and increasing production of thrombocytes. According to
10 another aspect of the invention, there is provided a composition for transporting a molecule via transcytosis across an endothelial cell barrier comprising said molecule in association with a recombinant tissue protective cytokine, as described herein above, and having at least one erythropoietic activity selected from the group consisting of increasing hematocrit, vasoactive action (vasoconstriction/vasodilatation), hyperactivating platelets, pro-coagulant
15 activity and increasing production of thrombocytes. In one embodiment, the association is a labile covalent bond, a stable covalent bond, or a non-covalent association with a binding site for said molecule. In another embodiment, the molecule is a receptor agonist or antagonist hormone, a neurotrophic factor, an antimicrobial agent, a radiopharmaceutical, an antisense oligonucleotide, an antibody, an immunosuppressant, a dye, a marker, or an
20 anti-cancer drug.

The invention also provides for the use of an recombinant tissue protective cytokine as described herein above, lacking at least one erythropoietic activity selected from the group consisting of increasing hematocrit, vasoactive action
(vasoconstriction/vasodilatation), hyperactivating platelets, pro-coagulant activities and
25 increasing production of thrombocytes. In one embodiment, the association is a labile covalent bond, a stable covalent bond, or a non-covalent association with a binding site for said molecule. In another embodiment, the molecule is a receptor agonist or antagonist hormone, a neurotrophic factor, an antimicrobial agent, a radiopharmaceutical, an antisense oligonucleotide, an antibody, an immunosuppressant, a dye, or a marker, or an anti-cancer
30 drug.

Thus, the invention is directed to a cellular protective use of any recombinant tissue protective cytokine with an alteration in at least one amino acid of the native erythropoietin counterpart, wherein the recombinant tissue protective cytokine has cellular protective

activity as described herein. Such cellular protective activity includes, but is not limited to, neuroprotective activity. The invention is further directed to a use of any of the aforementioned recombinant tissue protective cytokines in the treatment of a responsive cell, tissue or organ, in particular for treatment of a condition or disease involving such a responsive cell, tissue or organ. In one such embodiment, the recombinant tissue protective cytokines have at least one erythropoietic activity selected from the group consisting of increasing hematocrit, vasoactive action (vasoconstriction/vasodilatation), hyperactivating platelets, pro-coagulant activities and increasing production of thrombocytes. A recombinant tissue protective cytokine of the invention preferably maintains the three-dimensional conformation of native erythropoietin. The recombinant tissue protective cytokine may or may not have erythropoietic activity.

In one embodiment of the invention, the recombinant tissue protective cytokine is created as a recombinant protein with N terminal fusion of HisTag (6xHis residues). In certain embodiments, additional amino acid sequences may be added as a spacer. In a specific embodiment, the histidine-tagged recombinant tissue protective cytokine muteins of the invention, include, but are not limited to, K45D-6xHis and S100E-6xHis.

In another aspect of the invention, any of the foregoing recombinant tissue protective cytokines can be used in the preparation of a pharmaceutical composition for *ex vivo* treatment of cells, tissues, and organs for the purpose of protecting, maintaining, enhancing, or restoring the function or viability of responsive mammalian cells and their associated cells, tissues, and organs. Such *ex vivo* treatment is useful, for example, for the preservation of cells, tissues, or organs for transplant, whether autotransplant or xenotransplant. The cells, tissue or organ may be bathed in a solution comprising erythropoietin muteins or recombinant tissue protective cytokines, or the perfusate instilled into the organ through the vasculature or other means, to maintain cellular functioning during the period wherein the cells, tissue or organ is not integrated with the vasculature of the donor or recipient. Administration of the perfusate may be made to a donor prior to organ harvesting, as well as to the harvested organ and to the recipient. Moreover, the aforementioned use of any recombinant tissue protective cytokine is useful whenever a cell, tissue or organ is isolated from the vasculature of the individual and thus essentially existing *ex vivo* for a period of time, the term isolated referring to restricting or clamping the vasculature of or to the cell, tissue, organ or bodily part, such as may be performed during surgery, including, in particular, cardio pulmonary bypass surgery; bypassing the

vasculature of the cell, tissue, organ or bodily part; removing the cell, tissue, organ or bodily part from the mammalian body, such may be done in advance of xenotransplantation or prior to and during autotransplantation; or traumatic amputation of a cell, tissue, organ or bodily part. Thus, this aspect of the invention pertains both to the perfusion with an
5 erythropoietin mutein *in situ* and *ex vivo*. *Ex vivo*, the recombinant tissue protective cytokine may be provided in a cell, tissue or organ preservation solution. For either aspect, the exposing may be by way of continuous perfusion, pulsatile perfusion, infusion, bathing, injection, or catheterization.

In yet a further aspect, the invention is directed to a method for protecting,
10 maintaining, enhancing, or restoring the viability of a mammalian cell, tissue, organ or bodily part which includes a responsive cell or tissue, in which the cell, tissue, organ or bodily part is isolated from the mammalian body. The method includes at least exposing the isolated mammalian cell, tissue, organ or bodily part to an amount of an erythropoietin mutein or recombinant tissue protective cytokine for a duration which is effective to protect,
15 maintain, enhance, or restore the aforesaid viability. In non-limiting examples, isolated refers to restricting or clamping the vasculature of or to the cell, tissue, organ or bodily part, such as may be performed during surgery, in particular, cardio pulmonary bypass surgery; bypassing the vasculature of the cell, tissue, organ or bodily part; removing the cell, tissue, organ or bodily part from the mammalian body, such may be done in advance of
20 xenotransplantation or prior to and during autotransplantation; or traumatic amputation of a cell, tissue, organ or bodily part. Thus, this aspect of the invention pertains both to the perfusion with an erythropoietin mutein or recombinant tissue protective cytokine *in situ* and *ex vivo*. *Ex vivo*, the recombinant tissue protective cytokine may be provided in a cell, tissue or organ preservation solution. For either aspect, the exposing may be by way of
25 continuous perfusion, pulsatile perfusion, infusion, bathing, injection, or catheterization.

By way of non-limiting examples, the aforementioned *ex vivo* responsive cell or tissue may be or comprise neuronal, retinal, muscle, heart, lung, liver, kidney, small intestine, adrenal cortex, adrenal medulla, capillary endothelial, testis, ovary, pancreas, bone, bone marrow, skin, umbilical chord blood, or endometrial cells or tissue. These
30 examples of responsive cells are merely illustrative.

All of the foregoing methods and uses are preferably applicable to human beings, but are useful as well for any mammal, such as, but not limited to, companion animals, domesticated animals, livestock and zoo animals. Routes of administration of the

aforementioned pharmaceutical compositions include oral, intravenous, intranasal, topical, intraluminal, inhalation or parenteral administration, the latter including intravenous, intraarterial, subcutaneous, intramuscular, intraperitoneal, submucosal or intradermal. For *ex vivo* use, a perfusate or bath solution is preferred. This includes perfusing an isolated
5 portion of the vasculature *in situ*.

In yet another aspect of the invention, any of the aforementioned recombinant tissue protective cytokines are useful in preparing a pharmaceutical composition for restoring a dysfunctional cell, tissue or organ when administered after the onset of the disease or condition responsible for the dysfunction. By way of non-limiting example, administration
10 of a pharmaceutical composition comprising a recombinant tissue protective cytokine restores cognitive function in animals previously having brain trauma, even when administered long after (*e.g.*, one day, three days, five days, a week, a month, or longer) the initial trauma. The present invention encompasses pharmaceutical compositions for the treatment (*i.e.* ameliorating or reversing the symptoms or effects of) and prevention, (*i.e.*
15 delaying the onset of, inhibiting, or stopping) of subsequent damage to cells and tissues that cascades from initial trauma. Recombinant tissue protective cytokines useful for such applications include any of the particular aforementioned recombinant tissue protective cytokines. Any form of a recombinant tissue protective cytokine capable of benefiting responsive cells is embraced in this aspect of the invention.

In yet another embodiment, the invention provides methods for the use of the
20 aforementioned recombinant tissue protective cytokine for restoring a dysfunctional cell, tissue or organ when administered after the onset of the disease or condition responsible for the dysfunction. By way of non-limiting example, methods for administration of a pharmaceutical composition comprising a recombinant tissue protective cytokine restores
25 cognitive function in animals previously having brain trauma, even when administered long after (*e.g.*, three days, five days, a week, a month, or longer) the trauma has subsided. Recombinant tissue protective cytokines and further modifications thereof are as herein above described. Any form of a recombinant tissue protective cytokine capable of benefiting responsive cells is embraced in this aspect of the invention.

In still yet a further aspect of the present invention, methods are provided for
30 facilitating the transcytosis of a molecule across an endothelial cell barrier in a mammal by administration of a composition of a molecule in association with an erythropoietin mutein or a recombinant tissue protective cytokine as herein before described.

The association between the molecule to be transported and the recombinant tissue protective cytokine may be, for example, a labile covalent bond, a stable covalent bond, or a noncovalent association with a binding site for the molecule. The recombinant tissue protective cytokine and a protein to be transported may be expressed as a fusion
5 polypeptide. Endothelial cell barriers may be the blood-brain barrier, the blood-heart barrier, the blood-kidney barrier, the blood-eye barrier, the blood-testis barrier, the blood-ovary barrier and the blood-placenta barrier. Suitable molecules for transport by the method of the present invention include hormones, such as growth hormone, antibiotics, and anti-cancer agents.

10 It is a further aspect of the present invention to provide a composition for facilitating the transcytosis of a molecule across an endothelial cell barrier in a mammal, said composition comprising said molecule in association with a recombinant tissue protective cytokine such as is described above.

In a still further aspect of the present invention, any of the aforementioned
15 recombinant tissue protective cytokines are useful in preparing a pharmaceutical composition for facilitating the transcytosis of a molecule across an endothelial cell barrier in a mammal, said composition comprising said molecule in association with a recombinant tissue protective cytokine as described herein above.

The association may be, for example, a labile covalent bond, a fusion polypeptide, a
20 stable covalent bond, or a noncovalent association with a binding site for the molecule. Endothelial cell barriers may be the blood-brain barrier, the blood-eye barrier, the blood-testes barrier, the blood-ovary barrier, and the blood-placenta barrier. Suitable molecules for transport by the method of the present invention include, for example, hormones, such as growth hormone, neurotrophic factors, antibiotics, antivirals, or antifungals such as those
25 normally excluded from the brain and other barriered organs, peptide radiopharmaceuticals, antisense drugs, antibodies against biologically-active agents, pharmaceuticals, dyes, markers, and anti-cancer agents

These and other aspects of the present invention will be better appreciated by reference to the following Figures and Detailed Description.

Figure 1 shows the distribution of erythropoietin receptor in a normal human brain, in thin sections stained with an anti-erythropoietin antibody.

Figure 2 is a higher power magnification of the image in Figure 1.

Figure 3 shows, using gold-labeled secondary antibodies, the ultramicroscopic
5 distribution of erythropoietin receptors.

Figure 4, prepared similarly to Figure 3, shows high density erythropoietin receptors at the luminal and anti-luminal surfaces of human brain capillaries.

Figure 5 depicts the translocation of parenterally-administered erythropoietin into the cerebrospinal fluid.

10 **Figure 6A and 6B** indicates the results of the SK-N-SH neuroblastoma cell neuroprotection assay (against rotenone) for erythropoietin as well as the recombinant tissue protective cytokines with the K45D and S100E recombinant tissue protective cytokines. The y-axis on the graph indicates the absorbance readings, and the data are means \pm range of duplicate determinations. The graph within **Figure 6A** clearly indicates that the viability
15 of the cells within the K45D and S100E samples maintained their viability demonstrating their cellular protective effect. **Figure 6B** shows the plasmid map of hEPO-6xHisTag-PCiNeo.

Figure 7 compares the *in vitro* efficacy of erythropoietin and asialoerythropoietin on the viability of serum-starved P19 cells.

20 **Figure 8** is another experiment which compares the *in vitro* efficacy of erythropoietin and asialoerythropoietin on the viability of serum-starved P19 cells.

Figure 9 shows protection of erythropoietin and asialoerythropoietin in a rat focal cerebral ischemia model.

Figure 10 shows a dose response comparing the efficacy of human erythropoietin
25 and human asialoerythropoietin in middle cerebral artery occlusion in a model of ischemic stroke.

Figure 11 shows the activity of iodinated erythropoietin in the P19 assay.

5 **Figure 12** shows the effect of biotinylated erythropoietin and asialoerythropoietin in the P19 assay.

Figure 13 compares the *in vitro* efficacy of erythropoietin with phenylglyoxal-modified erythropoietin on the viability of serum-starved P19 cells.

10 **Figure 14** shows the effect of tissue protective cytokines in the water intoxication assay.

Figure 15 shows the maintenance of the function of a heart prepared for transplantation by an erythropoietin.

Figure 16 shows the protection of the myocardium from ischemic damage by erythropoietin after temporary vascular occlusion.

15 **Figure 17A, 17B, 17C, and 17D** depicts the effects of a erythropoietin treatment in a rat glaucoma model.

Figure 18 shows the extent of preservation of retinal function by an erythropoietin in the rat glaucoma model.

20 **Figure 19** depicts the restoration of cognitive function following brain trauma by administration of an erythropoietin starting five days after trauma.

Figure 20 depicts the restoration of cognitive function following brain trauma by administration of an erythropoietin starting 30 days after trauma.

Figure 21 depicts the efficacy of human asialoerythropoietin in a kainate model of cerebral toxicity.

25 **Figure 22** depicts the efficacy of tissue protective cytokines in a rat spinal cord injury model.

Figure 23 shows the efficacy of tissue protective cytokines within a rabbit spinal cord injury model.

30 **Figure 24A, 24B, and 24C** shows a coronal section of the brain cortical layer stained by hematoxylin and eosin.

Figure 25A, 25B, and 25C shows coronal sections of frontal cortex adjacent to the region of infarction stained by GFAP antibody.

Figure 26A and 26B shows coronal sections of brain cortical layer stained by OX-42 antibody.

35 **Figure 27A and 27B** shows coronal sections of brain cortical layer adjacent to the region of infarction stained by OX-42 antibody.

Figure 28 shows the efficacy of an erythropoietin against inflammation in an EAE model.

5 **Figure 29** compares the affects of dexamethasone and an erythropoietin on inflammation in the EAE model.

Figure 30A and 30B shows that erythropoietin suppresses inflammation associated with neuronal death.

Figure 31 shows that human erythropoietin and recombinant tissue protective
10 cytokines R130E and R150E effectively reduce cell death induced by NMDA when added to the primary hippocampal neuron cell cultures prior to NMDA treatment. Cells treated with R103E (5 nM) exhibited significantly less cell death in comparison to vehicle control cells ($p=0.01$). Cells treated with R103E (5 nM) exhibited significantly less cell death in
15 comparison to vehicle control cells ($p=0.01$). Cells treated with R150E (5 nM) exhibited approximately a 20% decrease in cell death in comparison to solvent control cells ($p=0.001$). Statistics: ANOVA plus Tukey's post-hoc test.

Figure 32 shows neuronal protection from serum withdrawal in P19 cells. The percent of apoptotic cells decreased for cells pretreated with Epo, EpoWT, and recombinant tissue protective cytokine S100E. Cells treated with Epo exhibited approximately a 20%
20 decrease in apoptotic cell death in comparison to untreated control cells. Cells treated with EpoWT and S100E both exhibited approximately a 10% decrease in apoptotic cell death in comparison to untreated control cells.

Figure 33A and 33B Show the effect of pre-incubation with S100E in differentiated PC12 cells submitted to NGF withdrawal in two independent experiments. Differentiated
25 PC12 cells were pre-treated with S100E at the indicated concentrations for 24 h, Figure 33A (3 pM) Figure 33B (0.00003 pM-3pM). Viability was measured in the MTT assay. NGF (100 ng/ml) was used as a positive control and NGF-free medium (-NGF) as a negative control. Data presented in Figure 33 are presented as % viability of positive control (+NGF) ($n=8$ in both experiments). There is a statistically significant increase in viability of S100E
30 treated cells compared to negative control cells (-NGF) by use of one-way ANOVA and Bonferroni post-hoc test. $***p<0.001$, $*p<0.05$. The effects observed with S100E were similar to those of Epo in this test system with respect to potency and efficacy.

Figure 34A, 34B, and 34C Shows the effect of pre-incubation with Epo in differentiated PC12 cells submitted to NGF withdrawal. Differentiated PC12 cells were pre-
35 treated with Epo, S100E, or carbamylated Epo (30 pM-30 nM) for 24 h. The chemically modified Epo molecule, AA24496, has a 10000 times lower activity than EPO in the UT-7 cell assay. Viability was measured in the MTT assay. NGF (100 ng/ml) was used as a positive control and NGF-free medium (-NGF) as a negative control.

5 **Figure 35** shows concentration-response curves of Epo, K45D and S100E in UT-7 cells. Different concentrations of Epo, EpoWT, K45D and S100E were added to UT-7 cells. Viability was measured 48 h later in the WST-1 assay. Data are mean \pm SD of three different experiments each performed in duplicate. The curve is a non-linear regression curve fit.

Figure 36 shows dose response curves of Epo, R103E and R150E in UT-7 cells.
10 Different concentrations of Epo, EpoWT, R103E and R150E were added to UT-7 cells. Viability was measured 48 h later in the WST-1 assay. Data are mean \pm SD of three different experiments each performed in duplicate. The curve is a non-linear regression curve fit.

Figure 37 is a graph demonstrating the locomotor ratings of the rats recovering from the spinal cord trauma over a period of forty-two days. As can be seen from the graph, the
15 rats that were given S100E recovered from the injury more readily and demonstrated better overall recovery from the injury than the control rats and rats administered methylprednisolone.

Figure 38 shows the ratio of the latency of the injured eye over the latency the normal eye for the various treatment regimens. The rat treated with EPO exhibited a latency of 1.2,
20 which is better than the rat treated with saline. Each of the four recombinant tissue protective cytokines resulted in latency results equal to or better than EPO with R103E, R150E, and S100E showing a statistical improvement over EPO.

The present invention relates to mutein recombinant tissue protective cytokines. In particular, the present invention provides compositions comprising isolated nucleic acid molecules encoding recombinant tissue protective cytokine muteins, as well as isolated and/or recombinant cells and vectors comprising the nucleic acid molecules. The invention further encompasses isolated polypeptides of mutein recombinant tissue protective cytokine, lacking at least one erythropoietic activity selected from the group consisting of increasing hematocrit, vasoactive action (vasoconstriction/vasodilatation), hyperactivating platelets, pro-coagulant activities and increasing production of thrombocytes, the cytokine having at least one responsive cellular protective activity selected from the group consisting of protecting, maintaining, enhancing or restoring the function or viability of a responsive mammalian cell, tissue, or organ. The invention also encompasses methods for protecting, maintaining or enhancing the viability of a cell, tissue, or organ isolated from a mammalian body using the recombinant tissue protective cytokine muteins of the invention, and use of such muteins in treatment and prevention of diseases and conditions.

“Responsive cell” refers to a mammalian cell whose function or viability may be maintained, promoted, enhanced, regenerated, or in any other way benefited, by exposure to an erythropoietin. Non-limiting examples of such cells include neuronal, retinal, muscle, heart, lung, liver, kidney, small intestine, adrenal cortex, adrenal medulla, capillary endothelial, testes, ovary, pancreas, bone, skin, and endometrial cells. In particular, responsive cells would include, without limitation, neuronal cells; Purkinje cells; retinal cells: photoreceptor (rods and cones), ganglion, bipolar, horizontal, amacrine, and Müller cells; muscle cells; heart cells: myocardium, pace maker, sinoatrial node, sinus node, and junction tissue cells (atrioventricular node and bundle of his); lung cells; liver cells: hepatocytes, stellate, and Kupffer cells; kidney cells: mesangial, renal epithelial, and tubular interstitial cells; small intestine cells: goblet, intestinal gland (crypts) and enteral endocrine cells; adrenal cortex cells: glomerulosa, fasciculate, and reticularis cells; adrenal medulla cells: chromaffin cells; capillary cells: pericyte cells; testes cells: Leydig, Sertoli, and sperm cells and their precursors; ovary cells: Graffian follicle and primordial follicle cells; pancreas cells: islets of Langerhans, α -cells, β -cells, γ -cells, and F-cells; bone cells: osteoprogenitors, osteoclasts, and osteoblasts; skin cells; endometrial cells: endometrial stroma and endometrial cells; as well as the stem and endothelial cells present in the above listed organs. Moreover, such responsive cells and the benefits provided thereto by a recombinant tissue protective cytokine may be extended to provide protection or enhancement indirectly to other cells that are not directly responsive, or of tissues or organs

which contain such non-responsive cells. These other cells, tissues, or organs which benefit indirectly from the enhancement of responsive cells present as part of the cells, tissue or organ as "associated" cells, tissues, and organs. Thus, benefits of a recombinant tissue protective cytokine as described herein may be provided as a result of the presence of a small number or proportion of responsive cells in a tissue or organ, for example, excitable or neuronal tissue present in such tissue, or the Leydig cells of the testis, which make testosterone. In one aspect, the responsive cell or its associated cells, tissues, or organs are not excitable cells, tissues, or organs, or do not predominantly comprise excitable cells or tissues.

10 The methods of the invention provide for the local or systemic protection or enhancement of cells, tissues, and organs within a mammalian body, under a wide variety of normal and adverse conditions, or protection of those which are destined for relocation to another mammalian body. In addition, restoration or regeneration of dysfunction is also provided. As mentioned above, the ability of an erythropoietin mutein or a recombinant tissue protective cytokine to cross a tight endothelial cell barrier and exert its positive effects on responsive cells (as well as other types of cells) distal to the vasculature offers the potential to prevent as well as treat a wide variety of conditions and diseases which otherwise cause significant cellular and tissue damage in an animal, including human beings, and moreover, permit success of heretofore untenable surgical procedures for which risk traditionally outweighed the benefits. The duration and degree of purposeful adverse conditions induced for ultimate benefit, such as high-dose chemotherapy, radiation therapy, prolonged *ex vivo* transplant survival, and prolonged periods of surgically-induced ischemia, may be carried out by taking advantage of the invention herein. However, the invention is not so limited, but includes as one aspect, methods or compositions wherein the target responsive cells are distal to the vasculature by virtue of an endothelial-cell barrier or endothelial tight junctions. In general, the invention is directed to any responsive cells and associated cells, tissues, and organs which may benefit from exposure to a recombinant tissue protective cytokine. Furthermore, cellular, tissue or organ dysfunction may be restored or regenerated after an acute adverse event (such as trauma) by exposure to a recombinant tissue protective cytokine.

The invention is therefore directed generally to the use of recombinant tissue protective cytokines for the preparation of pharmaceutical compositions for the aforementioned purposes in which cellular function is maintained, promoted, enhanced,

regenerated, or in any other way benefited. The invention is also directed to methods for maintaining, enhancing, promoting, or regenerating cellular function by administering to a mammal an effective amount of a recombinant tissue protective cytokine as described herein. The invention is further directed to methods for maintaining, promoting, enhancing,
5 or regenerating cellular function *ex vivo* by exposing a cell, a tissue or an organ to a recombinant tissue protective cytokine. The invention is also directed to a perfusate composition comprising a recombinant tissue protective cytokine for use in organ or tissue preservation.

The various methods of the invention utilize a pharmaceutical composition which at
10 least includes a recombinant tissue protective cytokine at an effective amount for the particular route and duration of exposure to exert positive effects or benefits on responsive cells within or removed from a mammalian body. Where the target cell, tissues, or organs of the intended therapy require the recombinant tissue protective cytokine to cross an endothelial cell barrier, the pharmaceutical composition includes the recombinant tissue
15 protective cytokine at a concentration which is capable, after crossing the endothelial cell barrier, of exerting its desirable effects upon the responsive cells. Molecules capable of interacting with an erythropoietin receptor, and modulating cellular protective activity within the cell are useful in the context of the present invention.

5.1. NUCLEIC ACIDS OF THE INVENTION

20 A recombinant tissue protective cytokine comprising a nucleic acid molecule of the invention includes nucleic acids encoding tissue protective cytokines comprising an erythropoietin mutein lacking or exhibiting a decrease in at least one erythropoietic activity selected from the group consisting of increasing hematocrit, vasoactive action (vasoconstriction/vasodilatation), hyperactivating platelets, pro-coagulant activities and
25 increasing production of thrombocytes, the cytokine having at least one responsive cellular protective activity selected from the group consisting of protecting, maintaining, enhancing or restoring the function or viability of a responsive mammalian cell, tissue or organ. A tissue protective cytokine comprising a nucleic acid molecule of the invention includes nucleic acids encoding the erythropoietin mutein, with the activity described above,
30 comprising one or more altered amino acid residue between position 11-15 of SEQ ID NO:10 [SEQ ID NO:1], position 44-51 of SEQ ID NO 10 [SEQ ID NO:2], position 100-108 of SEQ ID NO [SEQ ID NO:3], or position 146-151 of SEQ ID NO 10 [SEQ ID NO:4]. A tissue protective cytokine comprising a nucleic acid molecule of the invention includes

nucleic acids encoding the erythropoietin mutein, with the activity described above, comprising an altered amino acid residue at one or more of the following positions of SEQ ID NO: 10: 7, 20, 21, 29, 33, 38, 42, 59, 63, 67, 70, 83, 96, 126, 142, 143, 152, 153, 155, 156, or 161. A tissue protective cytokine comprising a nucleic acid molecule of the invention includes nucleic acids encoding the erythropoietin mutein, with the activity described above, comprising the amino acid sequence of SEQ ID NO: 10 with one or more of the following changes: an alanine at residue 6 of SEQ ID NO: 10, an alanine at residue 7 of SEQ ID NO: 10, a serine at residue 7 of SEQ ID NO: 10, an isoleucine at residue 10 of SEQ ID NO: 10, a serine at residue 11 of SEQ ID NO: 10, an alanine at residue 12 of SEQ ID NO: 10, an alanine at residue 13 of SEQ ID NO: 10, an alanine residue 14 of SEQ ID NO: 10, a glutamic acid at residue 14 of SEQ ID NO: 10, a glutamine at residue 14 of SEQ ID NO: 10, an alanine at residue 15 of SEQ ID NO: 10, a phenylalanine at residue 15 of SEQ ID NO: 10, an isoleucine at residue 15 of SEQ ID NO: 10, a glutamic acid at residue 20 of SEQ ID NO: 10, an alanine at residue 20 of SEQ ID NO: 10, an alanine at residue 21 of SEQ ID NO: 10, a lysine at residue 24 of SEQ ID NO: 10, a serine at residue 29 of SEQ ID NO: 10; a tyrosine at residue 29 of SEQ ID NO: 10, an asparagine at residue 30 of SEQ ID NO: 10, a threonine at residue 32 of SEQ ID NO: 10, a serine at residue 33 of SEQ ID NO: 10, a tyrosine at residue 33 of SEQ ID NO: 10, a lysine at residue 38 of SEQ ID NO: 10, a lysine at residue 83 of SEQ ID NO: 10, an asparagine at residue 42 of SEQ ID NO: 10, an alanine at residue 42 of SEQ ID NO: 10, an alanine at residue 43, an isoleucine at residue 44 of SEQ ID NO: 10, an aspartic acid at residue 45 of SEQ ID NO: 10, an alanine at residue 45 of SEQ ID NO: 10, an alanine at residue 46 of SEQ ID NO: 10, an alanine at residue 47 of SEQ ID NO: 10, an isoleucine at residue 48 of SEQ ID NO: 10, an alanine at residue 48 of SEQ ID NO: 10, an alanine at residue 49 of SEQ ID NO: 10, a serine at residue 49 of SEQ ID NO: 10, a phenylalanine at residue 51 of SEQ ID NO: 10, an asparagine at residue 51 of SEQ ID NO: 10, an alanine at residue 52 of SEQ ID NO: 10, an asparagine at residue 59 of SEQ ID NO: 10, a threonine at residue 62 of SEQ ID NO: 10, a serine at residue 67 of SEQ ID NO: 10, an alanine at residue 70 of SEQ ID NO: 10, an arginine at residue 96 of SEQ ID NO: 10, an alanine at residue 97 of SEQ ID NO: 10, an arginine at residue 100 of SEQ ID NO: 10, a glutamic acid at residue 100 of SEQ ID NO: 10 of SEQ ID NO: 10, an alanine at residue 100, a threonine at residue 100 of SEQ ID NO: 10, an alanine at residue 101 of SEQ ID NO: 10, an isoleucine at residue 101 of SEQ ID NO: 10, an alanine at residue 102 of SEQ ID NO: 10, an alanine at residue 103 of SEQ ID NO: 10, a glutamic acid at residue 103 of SEQ ID NO: 10, an alanine at residue 104 of SEQ ID NO: 10, an isoleucine at residue 104 of SEQ ID NO: 10, an alanine at residue 105 of

SEQ ID NO: 10, an alanine at residue 106 of SEQ ID NO: 10, an isoleucine at residue 106 of SEQ ID NO: 10, an alanine at residue 107 of SEQ ID NO: 10, a leucine at residue 107 of SEQ ID NO: 10, a lysine at residue 108 of SEQ ID NO: 10, an alanine at residue 108 of SEQ ID NO: 10, a serine at residue 108 of SEQ ID NO: 10, an alanine at residue 116 of
 5 SEQ ID NO: 10, an alanine at residue 126 of SEQ ID NO: 10, an alanine at residue 132 of SEQ ID NO: 10, an alanine at residue 133 of SEQ ID NO: 10, an alanine at residue 134 of SEQ ID NO: 10, an alanine at residue 140 of SEQ ID NO: 10, an isoleucine at residue 142 of SEQ ID NO: 10, an alanine at residue 143 of SEQ ID NO: 10, an alanine at residue 146 of SEQ ID NO: 10, a lysine at residue 147 of SEQ ID NO: 10, an alanine at residue 147 of
 10 SEQ ID NO: 10, a tyrosine at residue 148 of SEQ ID NO: 10, an alanine at residue 148 of SEQ ID NO: 10, an alanine at residue 149 of SEQ ID NO: 10, an alanine at residue 150 of SEQ ID NO: 10, a glutamic acid at residue 150 of SEQ ID NO: 10, an alanine at residue 151 of SEQ ID NO: 10, an alanine at residue 152 of SEQ ID NO: 10, a tryptophan at residue 152 of SEQ ID NO: 10, an alanine at residue 153 of SEQ ID NO: 10, an alanine at
 15 residue 154 of SEQ ID NO: 10, an alanine at residue 155 of SEQ ID NO: 10, an alanine at residue 158 of SEQ ID NO: 10, a serine at residue 160 of SEQ ID NO: 10, an alanine at residue 161 of SEQ ID NO: 10, or an alanine at residue 162 of SEQ ID NO: 10.

The nucleic acid molecules of the invention further include nucleotide sequences that encode recombinant erythropoietin muteins having at least 30%, 35%, 40%, 45%, 50%,
 20 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, or higher amino acid sequence identity to one of the erythropoietin muteins described above. To determine the percent identity of two amino acid sequences or of two nucleic acids encoding erythropoietin muteins, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal
 25 alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a
 30 function of the number of identical positions shared by the sequences (*i.e.*, % identity = # of identical overlapping positions/total # of overlapping positions x 100%). In one embodiment, the two sequences are the same length.

The nucleic acid molecules of the invention further include nucleotide sequences that encode recombinant erythropoietin muteins wherein the erythropoietin encoding nucleic acid sequence that is altered by one or more of the substitutions, deletions, or modifications described above comprises at least 30%, 35%, 40%, 45%, 50%, 55%, 60%, 5 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 98% sequence identity to SEQ ID NO: 7. The nucleic acid molecules of the invention also include nucleotide sequences that encode recombinant erythropoietin muteins wherein the erythropoietin encoding nucleic acid sequence that is altered by one or more of the substitutions, deletions, or modifications described above is a non-human erythropoietin encoding nucleic acid.

10 The determination of percent identity between two sequences can also be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, Proc. Natl. Acad. Sci. USA 87:2264-2268, modified as in Karlin and Altschul, 1993, Proc. Natl. Acad. Sci. USA 90:5873-5877. Such an algorithm is 15 incorporated into the NBLAST and XBLAST programs of Altschul, et al., 1990, J. Mol. Biol. 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to a nucleic acid molecule of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous 20 to a protein molecule of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., 1997, *Nucleic Acids Res.* 25:3389-3402. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules (Altschul et al., 1997, *supra*). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the 25 respective programs (e.g., XBLAST and NBLAST) can be used (see <http://www.ncbi.nlm.nih.gov>). Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, 1988, *CABIOS* 4:11-17. Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When 30 utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent identity, typically only exact matches are counted.

The nucleic acid molecules of the invention further include: (a) any nucleotide
5 sequence that hybridizes to an erythropoietin mutein or a recombinant tissue protective
cytokine encoding nucleic acid molecule of the invention described above, under stringent
conditions, *e.g.*, hybridization to filter-bound DNA in 6x sodium chloride/sodium citrate
(SSC) at about 45°C followed by one or more washes in 0.2xSSC/0.1% SDS at about 50-
65°C, or (b) under highly stringent conditions, *e.g.*, hybridization to filter-bound nucleic
10 acid in 6xSSC at about 45°C followed by one or more washes in 0.1x SSC/0.2% SDS at
about 68°C, or under other hybridization conditions which are apparent to those of skill in
the art (see, for example, Ausubel F.M. et al., eds., 1989, Current Protocols in Molecular
Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York,
at pp. 6.3.1-6.3.6 and 2.10.3). Preferably the encoding erythropoietin mutein nucleic acid
15 molecule that hybridizes under conditions described under (a) and (b), above, is one that
comprises the complement of a nucleic acid molecule that encodes a erythropoietin mutein.
In a preferred embodiment, nucleic acid molecules that hybridize under conditions (a) and
(b), above, encode protein products, *e.g.*, protein products functionally equivalent, *i.e.*
having one or more of the activities of erythropoietin described above, to an erythropoietin
20 mutein. Preferably, the nucleic acids of the invention are human.

The nucleic acid molecules of the invention further include the above nucleotide
sequences that hybridize to a erythropoietin mutein or a recombinant tissue protective
cytokine as described above and further lack or exhibit a decrease in at least one
erythropoietic activity selected from the group consisting of increasing hematocrit,
25 vasoactive action (vasoconstriction/vasodilatation), hyperactivating platelets, pro-coagulant
activities and increasing production of thrombocytes, the cytokine or mutein comprising at
least one responsive cellular protective activity selected from the group consisting of
protecting, maintaining, enhancing or restoring the function or viability of a responsive
mammalian cell, tissue or organ. The decrease may be a slight diminishment or near lack of
30 one of the erythropoietic activities. Such decreases can be measured by standard techniques
known in the art (Gruber et al., 2002, J. Biol Chem. 277(81):27581-27584; Page et al.,
1996, Cytokine 8(1):66-69; Park et al., 1997, Mol. Cells 7(6):699-704; Wolf et al., 1997,
Thromb Haemost 78:1505-1509; and Dale et al., 2002, Nature 415:175-179. The UT-7 cell

assays described in Section 6.17 are one, non-limiting, example of a technique to measure decreased or diminished erythropoietic activity.

The nucleic acid molecules of the invention further comprise the complements of the nucleic acids described above.

5 Fragments of the erythropoietin mutein nucleic acid molecules refer to erythropoietin mutein nucleic acid sequences described above that can be at least 10, 12, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1050, or more contiguous nucleotides in length. Alternatively, the fragments can comprise sequences that encode at least 10, 20, 30, 40, 50, 60, 70, 80 or more contiguous amino acid
10 residues of the erythropoietin mutein. In one embodiment, the erythropoietin mutein nucleic acid molecule encodes a gene product exhibiting at least one biological activity of a corresponding erythropoietin mutein. Fragments of the erythropoietin mutein nucleic acid molecules can also refer to portions of erythropoietin mutein coding regions that encode domains of, or mature erythropoietin mutein.

15 Erythropoietin derived from other organisms may be used to create the erythropoietin muteins of the invention. With respect to the cloning of variants of the erythropoietin mutein or recombinant tissue protective cytokine nucleic acids and homologous and orthologs from other species, the isolated erythropoietin nucleic acid sequences disclosed herein may be labeled and used to screen a cDNA library constructed
20 from mRNA obtained from appropriate cells or tissues derived from the organism of interest. The hybridization conditions used should generally be of a lower stringency when the cDNA library is derived from an organism different from the type of organism from which the labeled sequence was derived, and can routinely be determined based on, *e.g.*, relative relatedness of the target and reference organisms.

25 Alternatively, the labeled fragment may be used to screen a genomic library derived from the organism of interest, again, using appropriately stringent conditions. Appropriate stringency conditions are well known to those of skill in the art as discussed above, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example,
30 Sambrook, *et al.*, 1989, Molecular Cloning, A Laboratory Manual, Second Edition, Cold Spring Harbor Press, N.Y.; and Ausubel, *et al.*, 1989-1999, Current Protocols in Molecular

Biology, Green Publishing Associates and Wiley Interscience, N.Y., both of which are incorporated herein by reference in their entirety.

In a preferred embodiment, to make a recombinant tissue protective cytokine DNA can be amplified from genomic or cDNA (i.e. SEQ ID NO:7) by polymerase chain reaction (PCR) amplification using primers designed from the known sequence of a related or homologous recombinant tissue protective cytokine. PCR is used to amplify the desired sequence in DNA clone or a genomic or cDNA library, prior to selection. PCR can be carried out, e.g., by use of a thermal cycler and Taq polymerase (Gene Amp®). The polymerase chain reaction (PCR) is commonly used for obtaining genes or gene fragments of interest. For example, a nucleotide sequence encoding a recombinant tissue protective cytokine of any desired length can be generated using PCR primers that flank the nucleotide sequence encoding open reading frame. Alternatively, a recombinant tissue protective cytokine gene sequence can be cleaved at appropriate sites with restriction endonuclease(s) if such sites are available, releasing a fragment of DNA encoding the recombinant tissue protective cytokine gene. If convenient restriction sites are not available, they may be created in the appropriate positions by site-directed mutagenesis and/or DNA amplification methods known in the art (see, for example, Shankarappa et al., 1992, PCR Method Appl. 1: 277-278). The DNA fragment that encodes the recombinant tissue protective cytokine is then isolated, and ligated into an appropriate expression vector, care being taken to ensure that the proper translation reading frame is maintained.

Any technique for mutagenesis known in the art can be used to modify individual nucleotides in a DNA sequence, for purpose of making amino acid substitution(s) in the expressed peptide sequence, or for creating/deleting restriction sites to facilitate further manipulations. Such techniques include but are not limited to, chemical mutagenesis, *in vitro* site-directed mutagenesis (Hutchinson et al., 1978, J. Biol. Chem. 253: 6551), oligonucleotide-directed mutagenesis (Smith, 1985, Ann. Rev. Genet. 19: 423-463; Hill et al., 1987, Methods Enzymol. 155: 558-568) and as described in section 6.3, PCR-based overlap extension (Ho et al., 1989, Gene 77: 51-59), PCR-based megaprimer mutagenesis (Sarkar et al., 1990, Biotechniques 8: 404-407), etc. Modifications can be confirmed, e.g., by double-stranded dideoxynucleotide DNA sequencing.

The invention also includes nucleic acid molecules, preferably DNA molecules, that are the complements of the nucleotide sequences of the preceding paragraphs.

In certain embodiments, the nucleic acid molecules of the invention are present as part of nucleic acid molecules comprising nucleic acid sequences that contain or encode heterologous (*e.g.*, vector, expression vector, or fusion protein) sequences.

5.2. RECOMBINANT TISSUE PROTECTIVE CYTOKINES OF THE INVENTION

Recombinant tissue protective cytokines of the invention include erythropoietin muteins, that maintain partial or full erythropoietic activity. Erythropoietin is a glycoprotein hormone which in humans has a molecular weight of about 34 kDa. The mature protein comprises 165 amino acids, and the glycosyl residues comprise about 40% of the weight of the molecule. The forms of recombinant tissue protective cytokine useful in the practice of the present invention encompass at least a single amino acid change in naturally-occurring, synthetic and recombinant forms of the following human and other mammalian erythropoietin-related molecules: erythropoietin, asialoerythropoietin, deglycosylated erythropoietin, erythropoietin analogs, erythropoietin mimetics, erythropoietin fragments, hybrid erythropoietin molecules, erythropoietin receptor-binding molecules, erythropoietin agonists, renal erythropoietin, brain erythropoietin, oligomers and multimers thereof, and congeners thereof. Such equivalent recombinant tissue protective cytokines include mutant erythropoietins, which may contain substitutions, deletions, including internal deletions, additions, including additions yielding fusion proteins, or conservative substitutions of amino acid residues within and/or adjacent to the amino acid sequence, but that result in a "silent" change, in that the change produces a functionally equivalent erythropoietin mutein or recombinant tissue protective cytokine. In a preferred embodiment, the recombinant tissue protective cytokine is nonerythropoietic, *i.e.* lacking or exhibiting diminished erythropoietic activity. Conservative amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Alternatively, non-conservative amino acid changes, and larger insertions and deletions may be used to create functionally altered recombinant tissue protective cytokines. Such mutants can be used to alter erythropoietin

properties in desirable ways. For example, in one embodiment, an erythropoietin useful for the practice of the invention can be a recombinant tissue protective cytokine altered in one or more amino acids within the four functional domains of erythropoietin which affect receptor binding: VLQRY (SEQ ID NO:1) and/or TKVNFYAW (SEQ ID NO:2) and/or
5 SGLRSLTTL (SEQ ID NO:3) and/or SNFLRG (SEQ ID NO:4). In another embodiment, erythropoietins containing mutations in the surrounding areas of the molecule which affect the kinetics or receptor-binding properties of the molecule can be used. Determining which alterations, or which positions in the domains will effect binding can be accomplished using standard methods. For example, the domains may be altered by pair-wise alanine mutations
10 (ala-scanning mutagenesis) followed by measurement of binding kinetics of mutants to examine the effect on binding to a receptor (Bernat *et al.*, 2003, PNAS 100:952-957; Wells *et al.*, 1989, Science 244:1081-1085).

The term "recombinant tissue protective cytokine" as well as "a recombinant tissue protective cytokine" may be used interchangeably or conjunctively, to encompass the
15 recombinant tissue protective cytokines of the invention and further modifications thereof, such as deglycosylated, asialylated, and other partially glycosylated forms of the recombinant tissue protective cytokine, or chemical modifications of the amino acids. Non-limiting examples of such variants are described in Tsuda *et al.*, 1990, Eur. J. Biochem. 188:405-411, incorporated herein by reference. Cytokines are highly flexible, and, in the
20 case of human growth hormone it is known that flexibility is required for activation (Wells *et al.*, 1989, Science 244:1081-1085). Thus, mutations that stabilize the three dimensional structure of a cytokine, preventing normal activation of the erythropoietin receptor are encompassed by the instant invention. In addition, a variety of host systems may be used for expression and production of recombinant tissue protective cytokines, including, but not
25 limited to, bacteria, yeast, insect, plant, and mammalian, including human, cell systems. For example, recombinant erythropoietin produced in bacteria, which do not glycosylate, asialylate, or partially glycosylate the product, could be used to produce non-glycosylated forms of the recombinant tissue protective cytokine or may be further glycosylated using known methods in the art, such as, but not limited to, those techniques disclosed in U.S.
30 Patent Application Nos: US 2003/0040037 A1 and US 2003/0003529 for use of fucosylation to adjust glycosylation of proteins. Alternatively, recombinant tissue protective cytokine can be produced in other systems capable of glycosylating expressed proteins, *e.g.*, plants, and including human cells.

As noted above, the invention herein embraces any and all erythropoietin receptor activity modulator molecules capable of exerting positive activity on responsive cells, regardless of any structural relationship of the molecule with erythropoietin.

5 In addition, the recombinant tissue protective cytokine may be modified to tailor its activities for a specific tissue or tissues. Several non-limiting strategies which may be carried out to achieve this desired tissue specificity include modifications that shorten circulating half-life and thus reduce the time the recombinant tissue protective cytokine can interact with erythroid precursors, or modification of the primary structure of the erythropoietin mutein or recombinant tissue protective cytokine molecule. One approach to
10 reducing circulating half life is to remove or modify the glycosylation moieties, of which erythropoietin has three N-linked and one O-linked. Such variants of a glycosylated recombinant tissue protective cytokine can be produced in a number of ways. For example, techniques to modify the primary structure of erythropoietin to generate the tissue protective cytokines of the present invention are myriad and include substitution of one or more
15 specific amino acids, i.e., by mutating the amino acids at the N-linked or O-linked glycosylation sites and/or, chemical modification of one or more amino acids, or addition of other structures which interfere with the interaction of erythropoietin with any of its receptors. Use of such forms of recombinant tissue protective cytokines is fully embraced herein. The sialic acids which terminate the end of the sugar chains can be removed by
20 specific sialidases depending on the chemical linkage connecting the sialic acid to the sugar chain. Alternatively, the glycosylated structure can be dismantled in different ways by using other enzymes that cleave at specific linkages. In a preferred embodiment, the half-life of the non-erythropoietic recombinant tissue protective cytokine of the invention is reduced by about 90% from that of native erythropoietin.

25 Some of these recombinant tissue protective cytokine molecules will nevertheless mimic the actions of erythropoietin itself in other tissues or organs. For example, a 17-mer containing the amino-acid sequence of 31-47 of native erythropoietin is inactive for erythropoiesis but fully active for neural cells *in vitro* (Campana & O'Brien, 1998: Int. J. Mol. Med. 1:235-41).

30 Furthermore, derivative recombinant tissue protective cytokine molecules desirable for the uses described herein may be generated by guanidination, amidination, carbamylation (carbamoylation), trinitrophenylation, acylation such as acetylation or succinylation, nitration, or modification of arginine, aspartic acid, glutamic acid, lysine,

tyrosine, tryptophan, or cysteine residues or carboxyl groups, among other procedures, such as limited proteolysis, removal of amino groups, and/or mutational substitution of arginine, lysine, tyrosine, tryptophan, or cysteine residues by molecular biological techniques to produce erythropoietin muteins or recombinant tissue protective cytokines which maintain an adequate level of activities for specific organs and tissues but not for others, such as erythrocytes (*e.g.*, Satake *et al*; 1990, *Biochim. Biophys. Acta* 1038:125-9; incorporated herein by reference in its entirety). One non-limiting example as described hereinbelow is the modification of erythropoietin arginine residues by reaction with a glyoxal such as phenylglyoxal (according to the protocol of Takahashi, 1977, *J. Biochem.* 81:395-402). As will be seen below, such a recombinant tissue protective cytokine molecule fully retains the neurotrophic effect of erythropoietin. Such recombinant tissue protective cytokine molecules are fully embraced for the various uses and compositions described herein. In addition, these chemical modifications may be further used to enhance the protective effects of the recombinant tissue protective cytokines or neutralize any changes in the charge of the molecule resulting from the amino acid mutation of the native erythropoietin. Such modifications are described in co-pending applications; serial no. PCT/US01/49479, filed December 28, 2001; serial no. 09/753,132, filed December 29, 2000 and Attorney's Docket No. KW00-009C02-US, filed July 3, 2002, all of which are incorporated herein in their entireties.

Synthetic and recombinant molecules, such as brain erythropoietin and renal erythropoietin, recombinant mammalian forms of erythropoietin, as well as its naturally-occurring, tumor-derived, and recombinant isoforms, such as recombinantly-expressed molecules and those prepared by homologous recombination are provided herein. Furthermore, the present invention includes molecules including peptides which bind the erythropoietin receptor, as well as recombinant constructs or other molecules which possess part or all of the structural and/or biological properties of erythropoietin, including fragments and multimers of erythropoietin or its fragments. Erythropoietin muteins or other recombinant tissue protective cytokines which have additional or reduced numbers of glycosylation sites are included herein. As noted above, the terms "erythropoietin" and "mimetics" as well as the other terms are used interchangeably herein to refer to the responsive cell protective and enhancing molecules related to erythropoietin as well as the molecules which are capable of crossing endothelial cell barriers. Furthermore, molecules produced by transgenic animals are also encompassed here. It should be noted that erythropoietin molecules as embraced herein do not necessarily resemble erythropoietin

structurally or in any other manner, except for ability to interact with the erythropoietin receptor or modulate erythropoietin receptor activity or activate erythropoietin-activated signaling cascades, as described herein.

By way of non-limiting examples, forms of recombinant tissue protective cytokines
5 useful for the practice of the present invention include recombinant tissue protective
cytokines, such as those with altered amino acids at the carboxy terminus described in U.S.
Patent 5,457,089 and in U.S. Patent 4,835,260; asialoerythropoietin and erythropoietin
isoforms with various numbers of sialic acid residues per molecule, such as described in
10 U.S. Patent 5,856,298; polypeptides described in U.S. Patent 4,703,008; agonists described
in U.S. Patent 5,767,078; peptides which bind to the erythropoietin receptor as described in
U.S. Patents 5,773,569 and 5,830,851; small-molecule mimetics which activate the
erythropoietin receptor, as described in U.S. Patent 5,835,382; and erythropoietin analogs
described in WO 9505465, WO 9718318, and WO 9818926. All of the aforementioned
15 citations are incorporated herein to the extent that such disclosures refer to the various
alternate forms or processes for preparing such forms of the recombinant tissue protective
cytokines of the present invention.

Erythropoietin can be obtained commercially, for example, under the trademarks of
PROCRT, available from Ortho Biotech Inc., Raritan, NJ, and EPOGEN, available from
Amgen, Inc., Thousand Oaks, CA.

20 The activity (in units) of erythropoietin (EPO) and erythropoietin-like molecules is
traditionally defined based on its effectiveness in stimulating red cell production in rodent
models (and as derived by international standards of erythropoietin). One unit (U) of
regular erythropoietin (MW of ~ 30,000 to 34,000) is ~ 8 ng of protein (1 mg protein is
approximately 125,000 U). However, as the effect on erythropoiesis is incidental to the
25 desired activities herein and may not necessarily be a detectable property of certain of the
recombinant tissue protective cytokines of the invention, the definition of activity based on
erythropoiesis is inappropriate. Thus, as used herein, the activity unit of erythropoietin or
erythropoietin-related molecules is defined as the amount of protein required to elicit the
same activity in neural or other responsive cellular systems as is elicited by WHO
30 international standard erythropoietin in the same system. The skilled artisan will readily
determine the units of a non-erythropoietic recombinant tissue protective cytokine or related
molecule following the guidance herein.

The recombinant tissue protective cytokine muteins include, but are not limited to, those proteins and polypeptides encoded by the erythropoietin nucleic acid sequences described in Section 6.3. The invention encompasses muteins that are functionally equivalent to the erythropoietin gene product described in Section 6.3. Such erythropoietin gene products may contain one or more deletions, additions or substitutions of erythropoietin amino acid residues within the amino acid sequence encoded by an erythropoietin nucleic acid sequence, but which result in a silent change, thus producing a functionally equivalent erythropoietin gene product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved.

The recombinant tissue protective cytokine muteins of the invention can be generated by mutagenesis, e.g., discrete point mutation or truncation. A recombinant tissue protective cytokine mutein of the invention retains the cellular protective biological activities of the naturally occurring form, but may lack one or more of the erythropoietic activities of the naturally occurring form of the protein. Thus, specific biological effects can be elicited by addition of a mutein of limited function.

Modification of the structure of the recombinant tissue protective cytokine muteins can be for such purposes as enhancing efficacy, stability, or post-translational modifications (e.g., to alter the phosphorylation pattern of the muteins). Such modified recombinant tissue protective cytokine muteins, when designed to retain at least one cellular protective activity of the naturally-occurring form of the protein, or to produce specific antagonists thereof, are considered functional equivalents of the recombinant tissue protective cytokine muteins. Such modified recombinant tissue protective cytokine muteins can be produced, for instance, by amino acid substitution, deletion, or addition.

For example, it is reasonable to expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid (i.e. isosteric and/or isoelectric mutations) will not have a major effect on the biological activity of the resulting molecule.

Whether a change in the amino acid sequence of a recombinant tissue protective cytokine mutein results in a functional homolog, or non-functional homolog (i.e. lacking one or more of the activities of the non-mutated cytokine), can be readily determined by

assessing the ability of the variant mutein to produce a response in cells in a fashion similar to the wild-type cytokine, or competitively inhibit such a response. Recombinant tissue protective cytokine muteins in which more than one replacement has taken place can readily be tested in the same manner.

5 Muteins of the invention exhibiting altered function can be identified by screening combinatorial libraries of mutants, e.g., truncation mutants, of the recombinant tissue protective cytokine of the invention for desired activity or lack thereof. In one embodiment, a variegated library of variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of variants can
10 be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into nucleic acid sequences such that a degenerate set of potential protein sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (e.g., for phage display). There are a variety of methods which can be used to produce libraries of potential variants of the recombinant tissue protective cytokines of the invention
15 from a degenerate oligonucleotide sequence. Methods for synthesizing degenerate oligonucleotides are known in the art (see, e.g., Narang, 1983, Tetrahedron 39:3; Itakura et al., 1984, Annu. Rev. Biochem. 53:323; Itakura et al., 1984, Science 198:1056; Ike et al., 1983, Nucleic Acid Res. 11:477).

 In addition, libraries of fragments of the coding sequence of a recombinant tissue
20 protective cytokines of the invention can be used to generate a variegated population of recombinant tissue protective cytokines for screening and subsequent selection of muteins. For example, a library of coding sequence fragments can be generated by treating a double stranded PCR fragment of the coding sequence of interest with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA,
25 renaturing the DNA to form double stranded DNA which can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal and internal fragments of various sizes of the recombinant tissue protective cytokine muteins of
30 interest.

 Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. The most widely used techniques, which are amenable

to high through put analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify muteins of a recombinant tissue protective cytokine of the invention (Arkin and Yourvan, 1992, Proc. Natl. Acad. Sci. USA 89:7811-7815; Delgrave et al., 1993, Protein Engineering 6(3):327-331).

10 An isolated nucleic acid molecule encoding a mutein can be created by introducing one or more nucleotide substitutions, additions or deletions into the erythropoietin nucleotide sequence, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded recombinant tissue protective cytokine. Mutations can be introduced by standard techniques, such as site-directed mutagenesis and PCR-mediated
15 mutagenesis. Briefly, PCR primers are designed that delete the trinucleotide codon of the amino acid to be changed and replace it with the trinucleotide codon of the amino acid to be included. This primer is used in the PCR amplification of DNA encoding the recombinant tissue protective cytokine of interest. This fragment is then isolated and inserted into the full length cDNA encoding the tissue protective cytokine of interest and expressed
20 recombinantly. The resulting recombinant tissue protective cytokine now includes the amino acid replacement.

Either conservative or non-conservative amino acid substitutions can be made at one or more amino acid residues. Both conservative and non-conservative substitutions can be made. Conservative replacements are those that take place within a family of amino acids
25 that are related in their side chains. Genetically encoded amino acids can be divided into four families: (1) acidic = aspartate, glutamate; (2) basic = lysine, arginine, histidine; (3) nonpolar = alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan; and (4) uncharged polar = glycine, asparagine, glutamine, cysteine, serine, threonine, tyrosine. In similar fashion, the amino acid repertoire can be grouped as (1)
30 acidic = aspartate, glutamate; (2) basic = lysine, arginine histidine, (3) aliphatic = glycine, alanine, valine, leucine, isoleucine, serine, threonine, with serine and threonine optionally be grouped separately as aliphatic-hydroxyl; (4) aromatic = phenylalanine, tyrosine, tryptophan; (5) amide = asparagine, glutamine; and (6) sulfur -containing = cysteine and

methionine. (See, for example, Biochemistry, 4th ed., Ed. by L. Stryer, WH Freeman and Co.: 1995).

Alternatively, mutations can be introduced randomly along all or part of the coding sequence of a recombinant tissue protective cytokine, such as by saturation mutagenesis, and the resultant mutants can be screened for biological activity to identify mutants that retain activity. Following mutagenesis, the encoded protein can be expressed recombinantly and the activity of the recombinant tissue protective cytokine can be determined.

Further to the above-mentioned erythropoietin modifications useful herein, the following discussion expands on the various recombinant tissue protective cytokines of the invention. As described in Elliott *et al.*, Boissel *et al.*, and Wen *et al.*, mentioned above, the following erythropoietin muteins are useful for the purposes described herein, and may be provided in a pharmaceutical composition for the methods herein. In the mutein nomenclature used throughout herein, the changed amino acid is depicted with the native amino acid's one-letter code first, followed by its position in the erythropoietin molecule, followed by the replacement amino acid one-letter code. For example, "human erythropoietin S100E" or "recombinant tissue protective cytokine S100E" refers to a human erythropoietin molecule in which amino acid 100, a serine has been changed to glutamic acid. Such muteins useful for the practice of the present invention include but are not limited to human erythropoietin with at least one of the following amino acid changes:

I6A, C7A, C7S,
R10I, V11S, L12A, E13A, R14A, R14E, R14Q, Y15A, Y15F, Y15I,
K20E, K20A,
E21A,
N24K, C29S, C29Y, A30N, H32T,
C33S, C33Y, N38K, N83K,
P42N,
P42A, D43A, T44I, K45D, K45A, V46A, N47A, F48I, F48A, Y49A, Y49S, 44-49 deletion,
W51F, W51N, K52A,
Q59N,
E62T,
L67S,
L70A,
D96R, K97A

S100R, S100E, S100A, S100T, G101A, G101I, L102A, R103A, R103E, S104A, S104I,
 L105A, T106A, T106I, T107A, T107L, L108K, L108A, L108S,
 K116A,
 S126A,
 5 T132A,
 I133A, T134A,
 K140A,
 F142I,
 R143A,
 10 S146A, N147K, N147A, F148Y, P148A, L149A, R150A, R150E, G151A,
 K152A, K152W,
 L153A,
 K154A,
 L155A, G158A,
 15 C160S, C161A, or R162A.

In preferred embodiments, an erythropoietin mutein or a recombinant tissue
 protective cytokine of the invention comprises one or more of the above substitutions. In
 other embodiments, erythropoietin mutein or another recombinant tissue protective cytokine
 20 of the invention comprises one of the above substitutions or a combination thereof.

In an alternative embodiment, the recombinant tissue protective cytokines,
 pharmaceutical compositions, use, and treatment methods of the invention comprise one or
 more of the above substitutions with the proviso that they do not comprise one or more of
 the following substitutions: I6A, C7A, K20A, P42A, D43A, K45D, K45A, F48A, Y49A,
 25 K52A, K49A, S100E, R103A, K116A, T132A, I133A, K140A, N147K, N147A, R150A,
 R150E, G151A, K152A, K154A, G158A, C161A, or R162A. In a related embodiment of
 the invention, the recombinant tissue protective cytokines, pharmaceutical compositions,
 use, and treatment methods of the invention comprise one or more of the above substitutions
 with the proviso that they do not comprise any of the following combinations of
 30 substitutions: N24K/N38K/N83K or A30N/H32T.

In certain embodiments, more than one of the amino acid changes above can be
 combined to make a mutein. Examples of such combinations include, but are not limited to:
 K45D/S100E, A30N/H32T, K45D/R150E, R103E/L108S, K140A/K52A,

K140A/K52A/K45A, K97A/K152A, K97A/K152A/K45A, K97A/K152A/K45A/K52A, K97A/K152A/K45A/K52A/K140A, K97A/K152A/K45A/K52A/K140A/K154A, N24K/N38K/N83K, and N24K/Y15A. In certain embodiments, the recombinant tissue protective cytokine mutein of the invention does not comprise one or more of the above multiple substitutions. In certain embodiments the pharmaceutical compositions of the invention comprising the recombinant tissue protective cytokine mutein of the invention do not comprise one or more of the above multiple substitutions. In certain embodiments the use and treatment methods of the invention which utilize the recombinant tissue protective cytokine mutein of the invention do not comprise one or more of the above multiple substitutions.

Certain modifications or combinations of modifications can effect the flexibility of a erythropoietin muteins effecting binding to a receptor, such as the erythropoietin receptor or a secondary receptor to which erythropoietin or an erythropoietin mutein binds. Examples of such modifications or combinations thereof useful in the compositions and methods of the invention, include, but are not limited to, K152W, R14A/Y15A, I6A, C7A, D43A, P42A, F48A, Y49A, T132A, I133A, T134A, N147A, P148A, R150A, G151A, G158A, C161A, and R162A. Corresponding mutations are known to be detrimental in human growth hormone (Wells et al.). In certain embodiments, the recombinant tissue protective cytokine mutein of the invention does not comprise one or more of the above substitutions. In certain embodiments the pharmaceutical compositions of the invention comprising the recombinant tissue protective cytokine mutein of the invention do not comprise one or more of the above substitutions. In certain embodiments the use and treatment methods of the invention which utilize the recombinant tissue protective cytokine mutein of the invention do not comprise one or more of the above substitutions.

In addition to one of the foregoing amino acid modifications, a recombinant tissue protective cytokine of the invention may also have at least no sialic acid moieties, referred to as an asialoerythropoietin mutein. Preferably, an asialoerythropoietin mutein of the invention is human asialoerythropoietin. In alternative embodiments, the recombinant tissue protective cytokine of the invention may have at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, or 13 sialic acid residues. It may be prepared by desialylating a recombinant tissue protective cytokine using a sialidase, such as is described in the manufacturer's packaging for Sialydase A from ProZyme Inc., San Leandro, California. Typically, PROZYME® GLYCOPRO® sequencing-grade SIALYDASE A™ (N-acetylneuraminate glycohydrolase,

EC 3.2.1.18) is used to cleave all non-reducing terminal sialic acid residues from complex carbohydrates and glycoproteins such as erythropoietin. It will also cleave branched sialic acids (linked to an internal residue). Sialydase A is isolated from a clone of *Arthrobacter ureafaciens*.

5 A non-limiting example of sialylation of a glycopeptide is found in U.S. Patent Application No. US 2003/0040037, which discloses methods of sialylation using mammalian or bacterial sialyltransferases. Another non-limiting example of methods for sialylation and alteration of sialylation patterns on glycoproteins is found in U. S. Patent Application No. US 2002/0160460 A1 and in US 6,399,336 B1. Therein, *in vitro* methods
10 for sialylating recombinant glycoproteins are disclosed where a sialic acid donor moiety is combined with a glycoprotein having a galactose or N-acetylgalactosamine acceptor moiety. In such methods a sialyltransferase combined with the acceptor and donor attached a sialic acid to a saccharide.

 A recombinant tissue protective cytokine of the invention may have at least a
15 reduced number of N-linked carbohydrates. To remove N-linked carbohydrates, a recombinant tissue protective cytokine may be treated with hydrazine, in accordance, for example, with the methods described by Hermentin *et al.*, 1996, *Glycobiology* 6(2):217-30. As noted above, erythropoietin has three N-linked carbohydrate moieties; the present invention embraces those erythropoietins with two, one, or no N-linked carbohydrate.

20 A recombinant tissue protective cytokine of the invention may have at least a reduced carbohydrate content by virtue of treatment of a recombinant tissue protective cytokine with at least one glycosidase. For example, the procedure of Chen and Evangelista, 1998, *Electrophoresis* 19(15):2639-44, may be followed. Furthermore, removal of the O-linked carbohydrate may be achieved following the methods described in
25 Hokke *et al.*, 1995, *Eur. J. Biochem.* 228(3):981-1008.

 The carbohydrate portion of a recombinant tissue protective cytokine molecule may have at least a non-mammalian glycosylation pattern by virtue of the expression of a recombinant erythropoietin mutein in non-mammalian cells. Preferably, the recombinant tissue protective cytokines of the invention are expressed in insect or plant cells. By way of
30 non-limiting example, expression of a recombinant tissue protective cytokine in insect cells using a baculovirus expression system may be carried out in accordance with Quelle *et al.*, 1989, *Blood* 74(2):652-657. Another method is described in U.S. Patent 5,637,477.

Expression in a plant system may be carried out in accordance with the method of Matsumoto *et al.*, 1993, Biosci. Biotech. Biochem. 57(8):1249-1252. Alternatively, expression in bacteria will result in non-glycosylated forms of a recombinant tissue protective cytokine. These are merely exemplary of methods useful for the production of a recombinant tissue protective cytokine of the invention and are in no way limiting.

A non-limiting example of modification of glycosylation patterns is using fucosylation as disclosed in U.S. Patent Application No. US 2003/0040037 A1 and in U.S. Patent Application No. US2003/0003529 A1. Therein, methods are disclosed for modifying a glycosylation pattern of a glycopeptide by contacting a glycopeptide having an acceptor moiety for a fucosyltransferase with a reaction mixture having a fucose donor moiety to modify the glycosylation pattern of the glycopeptide. Methods are also disclosed for modification of glycosylation patterns using recombinant glycopeptide.

A recombinant tissue protective cytokine of the invention may have at least one or more oxidized carbohydrates that also may be chemically reduced. For example, the recombinant tissue protective cytokine may be a periodate-oxidized erythropoietin mutein; the periodate-oxidized erythropoietin mutein also may be chemically reduced with a borohydride salt such as sodium borohydride or sodium cyanoborohydride. Periodate oxidation of erythropoietin mutein may be carried out, for example, by the methods described by Linsley *et al.*, 1994, Anal. Biochem. 219(2):207-17. Chemical reduction following periodate oxidation may be carried out following the methods of Tonelli and Meints, 1978, J. Supramol. Struct. 8(1):67-78.

It should be noted that certain of the aforementioned and following amino acid modifications to a native erythropoietin may not be possible as the particular target amino acid for chemical modification in the native molecule has been altered to form the recombinant tissue protective cytokine of the invention. Of course, the altered amino acid may be subject to chemical modification in its own right, and the present invention embraces all such molecules. One of skill in the art will readily determine the available amino acid residues of a recombinant tissue protective cytokine of the invention and modification(s) available thereto.

A recombinant tissue protective cytokine for the aforementioned uses may have at least one or more modified arginine residues. For example, the recombinant tissue protective cytokine may comprise a R-glyoxal moiety on the one or more arginine residues,

where R may be an aryl, heteroaryl, lower alkyl, lower alkoxy, or cycloalkyl group, or an alpha-deoxyglycitoyl group. As used herein, the term lower "alkyl" means a straight- or branched-chain saturated aliphatic hydrocarbon group preferably containing 1-6 carbon atoms. Representative of such groups are methyl, ethyl, isopropyl, isobutyl, butyl, pentyl, 5 hexyl and the like. The term "alkoxy" means a lower alkyl group as defined above attached to the remainder of the molecule by oxygen. Examples of alkoxy include methoxy, ethoxy, propoxy, isopropoxy and the like. The term "cycloalkyl" refers to cyclic alkyl groups with three up to about 8 carbons, including for example cyclopropyl, cyclobutyl, cyclohexyl and the like. The term aryl refers to phenyl and naphthyl groups. The term heteroaryl refers to 10 heterocyclic groups containing 4-10 ring members and 1-3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur. Examples include but are not limited to isoxazolyl, phenylisoxazolyl, furyl, pyrimidinyl, quinolyl, tetrahydroquinolyl, pyridyl, imidazolyl, pyrrolidinyl, 1,2,4-triazolyl, thiazolyl, thienyl, and the like. The R group may be substituted, as for example the 2,3,4-trihydroxybutyl group of 3-deoxyglucosone.

15 Typical examples of R-glyoxal compounds are glyoxal, methylglyoxal, 3-deoxyglucosone, and phenylglyoxal. Preferred R-glyoxal compounds are methylglyoxal or phenylglyoxal. An exemplary method for such modification may be found in Werber *et al.*, 1975, *Isr. J. Med. Sci.* 11(11): 1169-70, using phenylglyoxal.

In a further example, at least one arginine residue may be modified by reaction with 20 a vicinal diketone such as 2,3-butanedione or cyclohexanedione, preferably in ca. 50 millimolar borate buffer at pH 8-9. A procedure for the latter modification with 2,3-butanedione may be carried out in accordance with Riordan, 1973, *Biochemistry* 12(20): 3915-3923; and that with cyclohexanone according to Patthy *et al.*, 1975, *J. Biol. Chem* 250(2): 565-9.

25 A recombinant tissue protective cytokine of the invention may comprise at least one or more modified lysine residues or a modification of the N-terminal amino group of the erythropoietin molecule, such modifications as those resulting from reaction of the lysine residue with an amino-group-modifying agent. In another embodiment, lysine residues may be modified by reaction with glyoxal derivatives, such as reaction with glyoxal, 30 methylglyoxal and 3-deoxyglucosone to form alpha-carboxyalkyl derivatives. Examples are reaction with glyoxal to form carboxymethyllysine as in Glomb and Monnier, 1995, *J. Biol. Chem.* 270(17):10017-26, or with methylglyoxal to form (1-carboxyethyl)lysine as in Degenhardt *et al.*, 1998, *Cell. Mol. Biol. (Noisy-le-grand)* 44(7):1139-45. The modified

lysine residue further may be chemically reduced. For example, a recombinant tissue protective cytokine may be biotinylated via lysine groups, in which D-biotinoyl- ϵ -aminocaproic acid-N-hydroxysuccinimide ester was reacted with erythropoietin, followed by removal of unreacted biotin by gel filtration on a Centricon 10 column, as described by
5 Wojchowski and Caslake, 1989, Blood 74(3):952-8. In this paper, the authors use three different methods of biotinylating erythropoietin, any of which may be used for the preparation of the erythropoietins for the uses herein. Biotin may be added to (1) the sialic acid moieties (2) carboxylate groups or (3) amino groups.

In another preferred embodiment, the lysine may be reacted with an aldehyde or
10 reducing sugar to form an imine, which may be stabilized by reduction as with sodium cyanoborohydride to form an N-alkylated lysine such as glucitolyl lysine, or which in the case of reducing sugars may be stabilized by Amadori or Heyns rearrangement to form an alpha-deoxy alpha-amino sugar such as alpha-deoxy-alpha-fructosyllsine. As an example, preparation of a fructosyllsine-modified protein by incubation with 0.5 M glucose in
15 sodium phosphate buffer at pH 7.4 for 60 days is described by Makita *et al.*, 1992, J. Biol. Chem. 267:5133-5138. In another example, the lysine group may be carbamylated, such as by virtue of reaction with cyanate ion, or alkyl- or aryl-carbamylated or -thiocarbamylated with an alkyl- or aryl-isocyanate or -isothiocyanate, or it may be acylated by a reactive alkyl- or arylcarboxylic acid derivative, such as by reaction with acetic anhydride or
20 succinic anhydride or phthalic anhydride. Exemplary are the modification of lysine groups with 4-sulfophenylisothiocyanate or with acetic anhydride, both as described in Gao *et al.*, 1994, Proc Natl Acad Sci USA 91(25):12027-30. Lysine groups may also be trinitrophenyl modified by reaction with trinitrobenzenesulfonic acid or preferably its salts.

At least one tyrosine residue of a recombinant tissue protective cytokine may be
25 modified in an aromatic ring position by an electrophilic reagent, such as by nitration or iodination. By way of non-limiting example, erythropoietin may be reacted with tetranitromethane (Nestler *et al.*, 1985, J. Biol. Chem. 260(12):7316-21; or iodinated as described in Example 4.

At least an aspartic acid or a glutamic acid residue of a recombinant tissue protective
30 cytokine may be modified, such as by reaction with a carbodiimide followed by reaction with an amine such as but not limited to glycnamide.

In another example, a tryptophan residue of a recombinant tissue protective cytokine may be modified, such as by reaction with n-bromosuccinimide or n-chlorosuccinimide, following methods such as described in Josse *et al.*, Chem Biol Interact 1999 May 14;119-120.

5 In yet another example, a recombinant tissue protective cytokine may be prepared by removing at least one amino group, such may be achieved by reaction with ninhydrin followed by reduction of the subsequent carbonyl group by reaction with borohydride.

 In still a further example, a recombinant tissue protective cytokine is provided that has at least an opening of at least one of the cysteine linkages in the erythropoietin molecule
10 by reaction with a reducing agent such as dithiothreitol, followed by reaction of the subsequent sulfhydryls with iodoacetamide, iodoacetic acid or another electrophile to prevent reformation of the disulfide linkages. As noted above, alternatively or in combination, disulfide linkages may be abolished by altering a cysteine molecule that participates in the actual cross-link or at least one other amino acid residue that results in the
15 inability of the erythropoietin mutein to form at least one of the disulfide linkages present in the native molecule.

 A recombinant tissue protective cytokine may be prepared by subjecting an erythropoietin to a limited chemical proteolysis that targets specific residues, for example, to cleave after tryptophan residues. Such resulting recombinant tissue protective cytokine
20 fragments are embraced herein.

 As noted above, a recombinant tissue protective cytokine useful for the purposes herein may have at least one of the aforementioned modifications, but may have more than one of the above modifications. By way of example of a recombinant tissue protective cytokine with one modification to the carbohydrate portion of the molecule and one
25 modification to the amino acid portion, a recombinant tissue protective cytokine may be asialoerythropoietin and have its lysine residue at position 45 changed to aspartic acid.

 Thus, various recombinant tissue protective cytokine molecules and pharmaceutical compositions containing them for the uses described herein are embraced. As mentioned above, such erythropoietin molecules include but are not limited to muteins that are further
30 asialoerythropoietin, N-deglycosylated erythropoietin, O-deglycosylated erythropoietin, erythropoietin with reduced carbohydrate content, erythropoietin with altered glycosylation patterns, erythropoietin with carbohydrates oxidized then reduced, arylglyoxal-modified

erythropoietin, alkylglyoxal-modified erythropoietin, 2,3-butanedione-modified erythropoietin, cyclohexanedione-modified erythropoietin, biotinylated erythropoietin, N-alkylated-lysyl-erythropoietin, glucitolyl lysine erythropoietin, alpha-deoxy-alpha-fructosyllysine-erythropoietin, carbamylated erythropoietin, acetylated erythropoietin, succinylated erythropoietin, alpha-carboxyalkyl erythropoietin, nitrated erythropoietin, iodinated erythropoietin, to name some representative yet non-limiting examples based on the teachings herein. Preferred are the aforementioned modified forms based on human erythropoietin.

Moreover, the invention encompasses the aforementioned recombinant tissue protective cytokines, and pharmaceutical compositions comprising such compounds. By way of non-limiting example, such recombinant tissue protective cytokines include periodate-oxidized erythropoietin mutein, glucitolyl lysine erythropoietin mutein, fructosyl lysine erythropoietin mutein, 3-deoxyglucosone erythropoietin mutein, and carbamylated asialoerythropoietin mutein.

5.3. EXPRESSION SYSTEMS

A variety of host-expression vector systems may be utilized to produce the recombinant tissue protective cytokines, including erythropoietin mutein molecules of the invention. Such host-expression systems represent vehicles by which the recombinant tissue protective cytokines of interest may be produced and subsequently purified, but also represent cells that may, when transformed or transfected with the appropriate nucleotide coding sequences, exhibit the modified erythropoietin gene product *in situ*. These include, but are not limited to, bacteria, insect, plant, mammalian, including human host systems, such as, but not limited to, insect cell systems infected with recombinant virus expression vectors (*e.g.*, baculovirus) containing the recombinant tissue protective cytokine product coding sequences; plant cell systems infected with recombinant virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (*e.g.*, Ti plasmid) containing recombinant tissue protective cytokine coding sequences; or mammalian cell systems, including human cell systems, (*e.g.*, HT1080, COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian viruses (*e.g.*, the adenovirus late promoter; the vaccinia virus 7.5K promoter).

An expression construct, as used herein, refers to a nucleotide sequence encoding a recombinant tissue protective cytokine operably associated with one or more regulatory regions which allows expression of the recombinant tissue protective cytokine in an appropriate host cell. "Operably-associated" refers to an association in which the regulatory regions and the recombinant tissue protective cytokine polypeptide sequence to be expressed are joined and positioned in such a way as to permit transcription, and ultimately, translation of the recombinant tissue protective cytokine sequence. A variety of expression vectors may be used for the expression of recombinant tissue protective cytokine, including, but not limited to, plasmids, cosmids, phage, phagemids, or modified viruses. Examples include bacteriophages such as lambda derivatives, or plasmids such as pBR322 or pUC plasmid derivatives or the Bluescript vector (Stratagene). Typically, such expression vectors comprise a functional origin of replication for propagation of the vector in an appropriate host cell, one or more restriction endonuclease sites for insertion of the recombinant tissue protective cytokine gene sequence, and one or more selection markers.

In preferred embodiments, the pCI-neo vector is used to anneal oligonucleotides to the original human EPO cDNA clone to introduce the mutations as described above. The pCI-neo vector contains the neomycin phosphotransferase gene, a selectable marker for mammalian cells. The pCI-neo Vector can be used for transient expression or for stable expression by selecting transfected cells with the antibiotic G-418. (Brondyk, 1995, New Mammalian Expression Vector with a selectable marker: pCI-neo. Promega Notes 51, 10-14).

For expression of recombinant tissue protective cytokine in mammalian host cells, a variety of regulatory regions can be used, for example, the SV40 early and late promoters, the cytomegalovirus (CMV) immediate early promoter, and the Rous sarcoma virus long terminal repeat (RSV-LTR) promoter. Inducible promoters that may be useful in mammalian cells include, but are not limited to, those associated with the metallothionein II gene, mouse mammary tumor virus glucocorticoid responsive long terminal repeats (MMTV-LTR), and the α -interferon gene (Williams et al., 1989, Cancer Res. 49: 2735-42 ; Taylor et al., 1990, Mol. Cell. Biol. 10: 165-75).

The efficiency of expression of the recombinant tissue protective cytokine in a host cell may be enhanced by the inclusion of appropriate transcription enhancer elements in the expression vector, such as those found in SV40 virus, Hepatitis B virus, cytomegalovirus,

immunoglobulin genes, metallothionein, α -actin (see Bittner et al., 1987, Methods in Enzymol. 153: 516-544; Gorman, 1990, Curr. Op. in Biotechnol. 1: 36-47).

5 The expression vector may also contain sequences that permit maintenance and replication of the vector in more than one type of host cell, or integration of the vector into the host chromosome. Such sequences may include but are not limited to replication origins, autonomously replicating sequences (ARS), centromere DNA, and telomere DNA. It may also be advantageous to use shuttle vectors that can be replicated and maintained in at least two types of host cells.

10 In addition, the expression vector may contain selectable or screenable marker genes for initially isolating or identifying host cells that contain DNA encoding a recombinant tissue protective cytokine. For long term, high yield production of recombinant tissue protective cytokines, stable expression in mammalian, plant, bacterial, or fungal cells can be used. A number of selection systems may be used for mammalian cells, including, but not limited, to the Herpes simplex virus thymidine kinase (Wigler et al., 1977, Cell 11: 223),
15 hypoxanthine-guanine phosphoribosyltransferase (Szybalski and Szybalski, 1962, Proc. Natl. Acad. Sci. U.S.A. 48: 2026), and adenine phosphoribosyltransferase (Lowy et al., 1980, Cell 22: 817) genes can be employed in tk-, hgprt- or aprt- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for dihydrofolate reductase (dhfr), which confers resistance to methotrexate (Wigler et al., 1980, Natl. Acad. Sci. U.S.A. 77: 3567; O'Hare et al., 1981, Proc. Natl. Acad. Sci. U.S.A. 78: 1527); gpt, which
20 confers resistance to mycophenolic acid (Mulligan and Berg, 1981, Proc. Natl. Acad. Sci. U.S.A. 78: 2072); neomycin phosphotransferase (neo), which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., 1981, J. Mol. Biol. 150: 1); and hygromycin phosphotransferase (hyg), which confers resistance to hygromycin (Santerre et al., 1984, Gene 30: 147). Other selectable markers, such as but not limited to histidinol and ZeocinTM can also be used.
25

30 In order to insert the recombinant tissue protective cytokine coding sequence into the cloning site of a vector, DNA sequences with regulatory functions, such as promoters, must be attached to the coding sequences. To do this, linkers or adapters providing the appropriate compatible restriction sites may be ligated to the ends of cDNA or synthetic DNA encoding a recombinant tissue protective cytokine, by techniques well known in the art (Wu et al., 1987, Methods Enzymol. 152: 343-349). Cleavage with a restriction enzyme can be followed by modification to create blunt ends by digesting back or filling in single-

stranded DNA termini before ligation. Alternatively, a desired restriction enzyme site can be introduced into a fragment of DNA by amplification of the DNA by use of PCR with primers containing the desired restriction enzyme site.

5 The expression construct comprising a recombinant tissue protective cytokine-coding sequence operably associated with regulatory regions can be directly introduced into appropriate host cells for expression and production of the recombinant tissue protective cytokines of the invention without further cloning (see e.g., U.S. Patent No. 5,580,859). The expression constructs may also contain DNA sequences that facilitate integration of the coding sequence into the genome of the host cell, e.g., via homologous recombination. In
10 this instance, it is not necessary to employ an expression vector comprising a replication origin suitable for appropriate host cells in order to propagate and express the recombinant tissue protective cytokines in the host cells.

Expression constructs containing cloned recombinant tissue protective cytokines coding sequences can be introduced into the mammalian host cell by a variety of techniques
15 known in the art, including but not limited to calcium phosphate mediated transfection (Wigler et al., 1977, Cell 11: 223-232), liposome-mediated transfection (Schaefer-Ridder et al., 1982, Science 215: 166-168), electroporation (Wolff et al., 1987, Proc. Natl. Acad. Sci. 84: 3344), and microinjection (Cappechi, 1980, Cell 22: 479-488).

In addition, a host cell strain may be chosen that modulates the expression of the
20 inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be
25 chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells that possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells, including human host cells, include but are not limited to HT1080, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, and WI38.

30 For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines that stably express the recombinant tissue protective cytokine-related molecule gene product may be engineered. Rather than using expression

vectors that contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (*e.g.*, promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days
5 in an enriched media, and then they are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to integrate the plasmid into their chromosomes in a stable manner and grow to form foci that in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines that express the recombinant tissue protective cytokine gene
10 product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the recombinant tissue protective cytokine gene product.

Any of the cloning and expression vectors described herein may be synthesized and assembled from known DNA sequences by techniques well known in the art. The
15 regulatory regions and enhancer elements can be of a variety of origins, both natural and synthetic. Some vectors and host cells may be obtained commercially. Non-limiting examples of useful vectors are described in Appendix 5 of Current Protocols in Molecular Biology, 1988, ed. Ausubel et al., Greene Publish. Assoc. & Wiley Interscience, which is incorporated herein by reference; and the catalogs of commercial suppliers such as Clontech
20 Laboratories, Stratagene Inc., and Invitrogen, Inc.

Alternatively, a number of viral-based expression systems may also be utilized with mammalian cells for recombinant expression of tissue protective cytokines. Vectors using DNA virus backbones have been derived from simian virus 40 (SV40) (Hamer et al., 1979, Cell 17: 725), adenovirus (Van Doren et al., 1984, Mol. Cell Biol. 4: 1653), adeno-
25 associated virus (McLaughlin et al., 1988, J. Virol. 62: 1963), and bovine papillomas virus (Zinn et al., 1982, Proc. Natl. Acad. Sci. 79: 4897). In cases where an adenovirus is used as an expression vector, the donor DNA sequence may be ligated to an adenovirus transcription/translation control region, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo*
30 recombination. Insertion in a non-essential region of the viral genome (*e.g.*, region E1 or E3) will result in a recombinant virus that is viable and capable of expressing heterologous products in infected hosts (see, *e.g.*, Logan and Shenk, 1984, Proc. Natl. Acad. Sci. U.S.A. 81: 3655-3659).

Alternatively, the vaccinia 7.5K promoter may be used (see, e.g., Mackett et al., 1982, Proc. Natl. Acad. Sci. U.S.A. 79: 7415-7419; Mackett et al., 1984, J. Virol. 49: 857-864; Panicali et al., 1982, Proc. Natl. Acad. Sci. U.S.A. 79: 4927-4931) In cases where a human host cell is used, vectors based on the Epstein-Barr virus (EBV) origin (OriP) and
5 EBV nuclear antigen 1 (EBNA-1; a trans-acting replication factor) may be used. Such vectors can be used with a broad range of human host cells, e.g., EBO-pCD (Spickofsky et al., 1990, DNA Prot. Eng. Tech. 2: 14-18), pDR2 and α DR2 (available from Clontech Laboratories).

Recombinant tissue protective cytokine expression can also be achieved by a
10 retrovirus-based expression system. In contrast to transfection, retroviruses can efficiently infect and transfer genes to a wide range of cell types including, for example, primary hematopoietic cells. In retroviruses such as Moloney murine leukemia virus, most of the viral gene sequences can be removed and replaced with a recombinant tissue protective cytokine coding sequence, while the missing viral functions can be supplied in trans. The
15 host range for infection by a retroviral vector can also be manipulated by the choice of envelope used for vector packaging.

For example, a retroviral vector can comprise a 5' long terminal repeat (LTR), a 3' LTR, a packaging signal, a bacterial origin of replication, and a selectable marker. The recombinant tissue protective cytokine DNA is inserted into a position between the 5' LTR and 3' LTR, such that transcription from the 5' LTR promoter transcribes the cloned DNA.
20 The 5' LTR comprises a promoter, including but not limited to an LTR promoter, an R region, a U5 region and a primer binding site, in that order. Nucleotide sequences of these LTR elements are well known in the art. A heterologous promoter as well as multiple drug selection markers may also be included in the expression vector to facilitate selection of
25 infected cells (see McLauchlin et al., 1990, Prog. Nucleic Acid Res. and Molec. Biol. 38: 91-135; Morgenstern et al., 1990, Nucleic Acid Res. 18: 3587-3596; Chouluka et al., 1996, J. Virol 70: 1792-1798; Boesen et al., 1994, Biotherapy 6: 291-302; Salmons and Gunzberg, 1993, Human Gene Therapy 4: 129-141; and Grossman and Wilson, 1993, Curr. Opin. in Genetics and Devel. 3: 110-114).

30 In one embodiment of the invention, a recombinant tissue protective cytokine deficient in sialic residues, or completely lacking sialic residues, may be produced in mammalian cell, including a human cell. Such cells may be engineered to be deficient in, or lacking, the enzymes that add sialic acids, i.e., the β -galactoside α 2,3 sialyltransferase ($\text{A}\alpha$

2,3 sialyltransferase@) and the β -galactoside α 2,6 sialyltransferase ($A\alpha$ 2,6 sialyltransferase@) activity). In one embodiment, a mammalian cell is used in which either or both the α 2,3 sialyltransferase gene and/or the α 2,6 sialyltransferase gene, is deleted. Such deletions may be constructed using gene knock-out techniques well known in the art.

5 In another embodiment, dihydrofolate reductase (DHFR) deficient Chinese Hamster Ovary (CHO) cells are used as the host cell for the production of recombinant tissue protective cytokines. CHO cells do not express the enzyme α 2,6 sialyltransferase and therefore do not add sialic acid in the 2,6 linkage to N-linked oligosaccharides of glycoproteins produced in these cells. As a result, recombinant proteins produced in CHO cells lack sialic acid in the

10 2,6 linkage to galactose (Sasaki *et al.* (1987; Takeuchi *et al. supra*; Mutsaers *et al.* Eur. J. Biochem. 156, 651 (1986); Takeuchi *et al.* J. Chromotogr. 400, 207 (1987). In one embodiment, to produce a host cell for the production of asialo-erythropoietin, the gene encoding α 2,3 sialyltransferase in CHO cells is deleted. Such α 2,3 sialyltransferase knock-out CHO cells completely lack sialyltransferase activity, and as a result, are useful for the

15 recombinant expression and production of asialoerythropoietin mutein.

In another embodiment, asialo glycoproteins can be produced by interfering with sialic acid transport into the Golgi apparatus *e.g.*, Eckhardt *et al.*, 1998, J. Biol. Chem. 273:20189-95). Using methods well known to those skilled in the art (*e.g.*, Oelmann *et al.*, 2001, J. Biol. Chem. 276:26291-300), mutagenesis of the nucleotide sugar CMP-sialic acid

20 transporter can be accomplished to produce mutants of Chinese hamster ovary cells. These cells cannot add sialic acid residues to glycoproteins such as a recombinant tissue protective cytokine and produce only asialoerythropoietin mutein.

Transfected mammalian cells producing erythropoietin mutein also produce cytosolic sialidase which if it leaks into the culture medium degrades sialoerythropoietin

25 mutein with high efficiency (*e.g.*, Gramer *et al.*, 1995 Biotechnology 13:692-698). Using methods well known to those knowledgeable in the art (*e.g.*, from information provided in Ferrari *et al.*, 1994, Glycobiology 4:367-373), cell lines can be transfected, mutated or otherwise caused to constitutively produce sialidase. In this manner, asialoerythropoietin mutein can be produced during the manufacture of asialoerythropoietin mutein.

30 The recombinant cells may be cultured under standard conditions of temperature, incubation time, optical density, and media composition. Alternatively, modified culture conditions and media may be used to enhance production of recombinant tissue protective cytokine. For example, recombinant cells may be grown under conditions that promote

inducible recombinant tissue protective cytokine expression. Any technique known in the art may be applied to establish the optimal conditions for producing recombinant tissue protective cytokines. Cellular lysates or extracts comprising recombinant tissue protective cytokines can be further purified to isolate recombinant tissue protective cytokines.

5 To facilitate purification of the recombinant tissue protective cytokines, a marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., 1989, PNAS 86:821, for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags
10 useful for purification include, but are not limited to, the hemagglutinin "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson et al., 1984, Cell 37:767) and the "flag" tag. Any purification method known in the art can be used (see *e.g.*, International Patent Publication WO 93/21232; EP 439,095; Naramura et al., 1994, *Immunol. Lett.* 39:91-99; U.S. Patent 5,474,981; Gillies et al., 1992, *PNAS* 89:1428-
15 1432; and Fell et al., 1991, *J. Immunol.* 146:2446-2452).

5.4. ASSAYS FOR TISSUE PROTECTIVE PROPERTIES OF THE RECOMBINANT TISSUE PROTECTIVE CYTOKINES

Following the manufacture of the recombinant tissue protective cytokines and in some embodiments further chemical modification of such tissue protective cytokines of the
20 present invention, one of ordinary skill in the art can verify the tissue protective attributes of the cytokines and the absence of an effect on the bone marrow using well known assays.

For example, the non-erythropoietic affect of a recombinant tissue protective cytokine can be verified through the use of a TF-1 assay. In this assay, TF-1 cells are grown in a complete RPMI medium supplemented with 5 ng/ml of GM-CSF and 10% FCS
25 for a day at 37 C in a CO2 incubator. The cells are then washed in and suspended at a density of 106 cells/ml for 16 h in starvation medium (5% FCS without GM-CSF). A 96 well plate is prepared by: (1) adding 100 μ l of sterile water to the outer wells to maintain moisture; (2) adding medium (10% FCS without cells or GM-CSF) alone to 5 wells; and (3) seeding 25,000 cells/well with medium containing 10% FCS and the recombinant tissue
30 protective cytokines in the remaining cells (five wells per cytokine being tested). If the cells proliferate, the recombinant tissue protective cytokine may be erythropoietic. The *in vivo* effect of the compound should then be tested on an *in vivo* assay monitoring the

increase of hematocrit due to the recombinant tissue protective cytokine. A negative result – non proliferation of cells in the TF-1 *in vitro* assay and/or no increase in hematocrit within the *in vivo* assay – means that the recombinant tissue protective cytokine is nonerythropoietic.

5 As an alternative to the TF-1 assay described above, one skilled in the art may employ other erythropoietic assays known in the art, including, but not limited to, UT-7 cell assays, such as those described below in the Examples sections.

10 The tissue protective properties of the recombinant tissue protective cytokines may be verified using a P-19 *in vitro* assay or a water intoxication *in vivo* assay in mice, both of which are outlined in further detail below. Alternative assays, include but are not limited to the additional assays outlined in the Examples below, such as the PC-12, and hippocampal slice assays. The above assays are provided merely as examples, and other suitable assays for determining the tissue protective effects and/or bone marrow effects of the recombinant tissue protective cytokines known to those of ordinary skill in the art are contemplated as well.

5.5. PHARMACEUTICAL COMPOSITIONS OF THE INVENTION

15 In the practice of one aspect of the present invention, a pharmaceutical composition as described above containing a recombinant tissue protective cytokine may be administerable to a mammal by any route which provides a sufficient level of a recombinant tissue protective cytokine in the vasculature to permit translocation across an endothelial cell barrier and beneficial effects on responsive cells. When used for the purpose of perfusing a tissue or organ, similar results are desired. In the instance wherein the erythropoietin mutein is used for *ex-vivo* perfusion, the recombinant tissue protective cytokine may be any form of erythropoietin mutein, such as the aforementioned recombinant tissue protective cytokine. In the instance where the cells or tissue is non-vascularized and/or the administration is by bathing the cells or tissue with the composition of the invention, the pharmaceutical composition provides an effective responsive cell-beneficial amount of a recombinant tissue protective cytokine. The endothelial cell barriers across which a recombinant tissue protective cytokine may translocate include tight junctions, perforated junctions, fenestrated junctions, and any other types of endothelial barriers present in a mammal. A preferred barrier is an endothelial cell tight junction, but the invention is not so limiting.

The aforementioned recombinant tissue protective cytokines are useful generally for the therapeutic or prophylactic treatment of human diseases of the central nervous system or peripheral nervous system which have primarily neurological or psychiatric symptoms, ophthalmic diseases, cardiovascular diseases, cardiopulmonary diseases, respiratory
5 diseases, kidney, urinary and reproductive diseases, gastrointestinal diseases and endocrine and metabolic abnormalities. In particular, such conditions and diseases include hypoxic conditions, which adversely affect excitable tissues, such as excitable tissues in the central nervous system tissue, peripheral nervous system tissue, or cardiac or retinal tissue such as, for example, brain, heart, or retina/eye. Therefore, the invention can be used to treat or
10 prevent damage to excitable tissue resulting from hypoxic conditions in a variety of conditions and circumstances. Non-limiting examples of such conditions and circumstances are provided in the table hereinbelow.

In the example of the protection of neuronal tissue pathologies treatable in accordance with the present invention, such pathologies include those which result from
15 reduced oxygenation of neuronal tissues. Any condition which reduces the availability of oxygen to neuronal tissue, resulting in stress, damage, and finally, neuronal cell death, can be treated by the methods of the present invention. Generally referred to as hypoxia and/or ischemia, these conditions arise from or include, but are not limited to, stroke, vascular occlusion, prenatal or postnatal oxygen deprivation, suffocation, choking, near drowning,
20 carbon monoxide poisoning, smoke inhalation, trauma, including surgery and radiotherapy, asphyxia, epilepsy, hypoglycemia, chronic obstructive pulmonary disease, emphysema, adult respiratory distress syndrome, hypotensive shock, septic shock, anaphylactic shock, insulin shock, sickle cell crisis, cardiac arrest, dysrhythmia, nitrogen narcosis, and neurological deficits caused by heart-lung bypass procedures.

25 In one embodiment, for example, the specific recombinant tissue protective cytokine compositions can be administered to prevent injury or tissue damage resulting from risk of injury or tissue damage during surgical procedures, such as, for example, tumor resection or aneurysm repair. Other pathologies caused by or resulting from hypoglycemia which are treatable by the methods described herein include insulin overdose, also referred to as
30 iatrogenic hyperinsulinemia, insulinoma, growth hormone deficiency, hypocortisolism, drug overdose, and certain tumors.

Other pathologies resulting from excitable neuronal tissue damage include seizure disorders, such as epilepsy, convulsions, or chronic seizure disorders. Other treatable

conditions and diseases include, but are not limited to, diseases such as stroke, multiple sclerosis, hypotension, cardiac arrest, Alzheimer's disease, Parkinson's disease, cerebral palsy, brain or spinal cord trauma, AIDS dementia, age-related loss of cognitive function, memory loss, amyotrophic lateral sclerosis, seizure disorders, alcoholism, retinal ischemia, optic nerve damage resulting from glaucoma, and neuronal loss.

The specific composition and methods of the present invention may be used to treat inflammation resulting from disease conditions or various traumas, such as physically or chemically induced inflammation. Such traumas could include angitis, chronic bronchitis, pancreatitis, osteomyelitis, rheumatoid arthritis, glomerulonephritis, optic neuritis, temporal arteritis, encephalitis, meningitis, transverse myelitis, dermatomyositis, polymyositis, necrotizing fasciitis, hepatitis, and necrotizing enterocolitis.

Evidence has demonstrated that activated astrocytes can exert a cytotoxic role towards neurons by producing neurotoxins. Nitric oxide, reactive oxygen species, and cytokines are released from glial cells in response to cerebral ischemia (see Becker, K.J. 2001. Targeting the central nervous system inflammatory response in ischemic stroke. *Curr Opin Neurol* 14:349-353 and Mattson, M.P., Culmsee, C., and Yu, Z.F. 2000. Apoptotic and Antiapoptotic mechanisms in stroke. *Cell Tissue Res* 301:173-187.). Studies have further demonstrated that in models of neurodegeneration, glial activation and subsequent production of inflammatory cytokines depends upon primary neuronal damage (see Viviani, B., Corsini, E., Galli, C.L., Padovani, A., Ciusani, E., and Marinovich, M. 2000. Dying neural cells activate glia through the release of a protease product. *Glia* 32:84-90 and Rabuffetti, M., Scioratti, C., Tarozzo, G., Clementi, E., Manfredi, A.A., and Beltramo, M. 2000. Inhibition of caspase-1-like activity by Ac-Tyr-Val-Ala-Asp-chloromethyl ketone includes long lasting neuroprotection in cerebral ischemia through apoptosis reduction and decrease of proinflammatory cytokines. *J Neurosci* 20:4398-4404). Inflammation and glial activation is common to different forms of neuro degenerative disorders, including cerebral ischemia, brain trauma and experimental allergic encephalomyelitis, disorders in which erythropoietin exerts a cellular protective effect. Inhibition of cytokine production by erythropoietin could, at least in part, mediate its protective effect. However, unlike "classical" anti-inflammatory cytokines such as IL-10 and IL-13, which inhibit tumor necrosis factor production directly, erythropoietin appears to be active only in the presence of neuronal death.

While not wishing to be bound by any particular theory, it appears that this anti-inflammatory activity may be hypothetically explained by several non-limiting theories. First, since erythropoietin prevents apoptosis, inflammatory events triggered by apoptosis would be prevented. Additionally, erythropoietin may prevent the release of molecular
5 signals from dying neurons which stimulate the glia cells or could act directly on the glial cells reducing their reaction to these products. Another possibility is that erythropoietin targets more proximal members of the inflammatory cascade (*e.g.*, caspase 1, reactive oxygen or nitrogen intermediates) that trigger both apoptosis and inflammation.

Furthermore, erythropoietin appears to provide anti-inflammatory protection without
10 the rebound affect typically associated with other anti-inflammatory compounds such as dexamethasone. Once again, not wishing to be bound by any particular theory, it appears as though this may be due to erythropoietin's affect on multipurpose neuro toxins such as nitric oxide (NO). Although activated astrocytes and microglia produce neurotoxic quantities of NO in response to various traumas, NO serves many purposes within the body
15 including the modulation of essential physiological functions. Thus, although the use of an anti-inflammatory may alleviate inflammation by suppressing NO or other neuro toxins, if the anti-inflammatory has too long a half-life it may also interfere with these chemicals' roles in repairing the damage resulting from the trauma that led to the inflammation. It is hypothesized that the recombinant tissue protective cytokines of the present invention are
20 able to alleviate the inflammation without interfering with the restorative capabilities of neurotoxins such as NO.

The specific compositions and methods of the invention may be used to treat conditions of, and damage to, retinal tissue. Such disorders include, but are not limited to retinal ischemia, macular degeneration, retinal detachment, retinitis pigmentosa,
25 arteriosclerotic retinopathy, hypertensive retinopathy, retinal artery blockage, retinal vein blockage, hypotension, and diabetic retinopathy.

In another embodiment, the methods principles of the invention may be used to protect or treat injury resulting from radiation damage to excitable tissue. A further utility of the methods of the present invention is in the treatment of neurotoxin poisoning, such as
30 domoic acid shellfish poisoning, neurolathyrism, and Guam disease, amyotrophic lateral sclerosis, and Parkinson's disease.

As mentioned above, the present invention is also directed to a method for enhancing excitable tissue function in a mammal by peripheral administration of a recombinant tissue protective cytokine as described above. Various diseases and conditions are amenable to treatment using this method, and further, this method is useful for enhancing cognitive function in the absence of any condition or disease. These uses of the present invention are described in further detail below and include enhancement of learning and training in both human and non-human mammals.

Conditions and diseases treatable by the methods of this aspect of the present invention directed to the central nervous system include, but are not limited to, mood disorders, anxiety disorders, depression, autism, attention deficit hyperactivity disorder, and cognitive dysfunction. These conditions benefit from enhancement of neuronal function. Other disorders treatable in accordance with the teachings of the present invention include for example, sleep disruption, sleep apnea, and travel-related disorders; subarachnoid and aneurismal bleeds, hypotensive shock, concussive injury, septic shock, anaphylactic shock, and sequelae of various encephalitides and meningitides, for example, connective tissue disease-related cerebritides such as lupus. Other uses include prevention of or protection from poisoning by neurotoxins, such as domoic acid shellfish poisoning, neurolept anesthesia, and Guam disease, amyotrophic lateral sclerosis, Parkinson's disease; postoperative treatment for embolic or ischemic injury; whole brain irradiation; sickle cell crisis; and eclampsia.

A further group of conditions treatable by the methods of the present invention include mitochondrial dysfunction, of either a hereditary or an acquired nature, which are the cause of a variety of neurological diseases typified by neuronal injury and death. For example, Leigh disease (subacute necrotizing encephalopathy) is characterized by progressive visual loss and encephalopathy, due to neuronal drop out, and myopathy. In these cases, defective mitochondrial metabolism fails to supply enough high energy substrates to fuel the metabolism of excitable cells. An erythropoietin receptor activity modulator optimizes failing function in a variety of mitochondrial diseases. As mentioned above, hypoxic conditions adversely affect excitable tissues. The excitable tissues include, but are not limited to, central nervous system tissue, peripheral nervous system tissue, and heart tissue. In addition to the conditions described above, the methods of the present invention are useful in the treatment of inhalation poisoning, such as carbon monoxide and smoke inhalation, severe asthma, adult respiratory distress syndrome, choking, and near

drowning. Further conditions which create hypoxic conditions or by other means induce excitable tissue damage include hypoglycemia that may occur in inappropriate dosing of insulin, or with insulin-producing neoplasms (insulinoma).

Various neuropsychologic disorders which are believed to originate from excitable tissue damage are treatable by the instant methods. Chronic disorders in which neuronal damage is involved and for which treatment by the present invention is provided include disorders relating to the central nervous system and/or peripheral nervous system including age-related loss of cognitive function and senile dementia, chronic seizure disorders, Alzheimer's disease, Parkinson's disease, dementia, memory loss, amyotrophic lateral sclerosis, multiple sclerosis, tuberous sclerosis, Wilson's Disease cerebral and progressive supranuclear palsy, Guam disease, Lewy body dementia, prion diseases, such as spongiform encephalopathies, *e.g.*, Creutzfeldt-Jakob disease, Huntington's disease, myotonic dystrophy, Freidrich's ataxia and other ataxias, as well as Gilles de la Tourette's syndrome, seizure disorders such as epilepsy and chronic seizure disorder, stroke, brain or spinal cord trauma, AIDS dementia, alcoholism, autism, retinal ischemia, glaucoma, autonomic function disorders such as hypertension and sleep disorders, and neuropsychiatric disorders that include, but are not limited to, schizophrenia, schizoaffective disorder, attention deficit disorder hyperactivity, dysthymic disorder, major depressive disorder, mania, obsessive-compulsive disorder, psychoactive substance use disorders, anxiety, panic disorder, as well as unipolar and bipolar affective disorders. Additional neuropsychiatric and neurodegenerative disorders include, for example, those listed in the American Psychiatric Association's Diagnostic and Statistical Manual of Mental Disorders (DSM), the most current version, IV, of which is incorporated herein by reference in its entirety.

In another embodiment, recombinant chimeric toxin molecules comprising a recombinant tissue protective cytokine can be used for therapeutic delivery of toxins to treat a proliferative disorder, such as cancer, or viral disorder, such as subacute sclerosing panencephalitis.

The following table lists additional exemplary, non-limiting indications as to the various conditions and diseases amenable to treatment by the aforementioned recombinant tissue protective cytokines.

<i>Cell, tissue or organ</i>	<i>Dysfunction or pathology</i>	<i>Condition or disease</i>	<i>Type</i>
Heart	Ischemia	Coronary artery disease	Acute, chronic Stable, unstable

<i>Cell, tissue or organ</i>	<i>Dysfunction or pathology</i>	<i>Condition or disease</i>	<i>Type</i>
		Myocardial infarction	Dressler's syndrome
		Angina	
		Congenital heart disease	Valvular
		Prinzmetal angina	Cardiomyopathy
		Cardiac rupture	Aneurysmatic
		Angiitis	Septal perforation
	Arrhythmia	Tachy-, bradyarrhythmia Supraventricular, ventricular Conduction abnormalities	Stable, unstable Hypersensitive carotid sinus node
	Congestive heart failure	Left, right, bi-ventricular, systolic, diastolic	Cardiomyopathies, such as idiopathic familial, infective, metabolic, storage disease, deficiencies, connective tissue disorder, infiltration and granulomas, neurovascular
		Myocarditis	Autoimmune, infective, idiopathic
		Cor pulmonale	
	Blunt and penetrating trauma		
Vascular	Toxins	Cocaine toxicity	
	Hypertension	Primary, secondary	
	Decompression sickness		
	Fibromuscular hyperplasia		
	Aneurysm	Dissecting, ruptured, enlarging	
Lungs	Obstructive	Asthma	
		Chronic bronchitis, Emphysema and airway obstruction	
	Ischemic lung disease	Pulmonary embolism, Pulmonary thrombosis, Fat embolism	
	Environmental lung diseases		
	Ischemic lung disease	Pulmonary embolism Pulmonary thrombosis	
	Interstitial lung disease	Idiopathic pulmonary fibrosis	
	Congenital	Cystic fibrosis	
	Cor pulmonale		
	Trauma		
	Pneumonia and pneumonitides	Infectious, parasitic, toxic, traumatic, burn, aspiration	
Pancreas	Endocrine	Diabetes mellitus, type I and II	Beta cell failure, dysfunction Diabetic neuropathy
		Other endocrine cell failure of the pancreas	
	Exocrine	Exocrine pancreas failure	pancreatitis

<i>Cell, tissue or organ</i>	<i>Dysfunction or pathology</i>	<i>Condition or disease</i>	<i>Type</i>
Bone	Osteopenia	Primary secondary	Hypogonadism immobilisation Postmenopausal Age-related Hyperparathyroidism Hyperthyroidism Calcium, magnesium, phosphorus and/or vitamin D deficiency
	Osteomyelitis		
	Avascular necrosis		
	Trauma		
	Paget's disease		
Skin	Alopecia	Areata Totalis	Primary Secondary Male pattern baldness
	Vitiligo	Localized generalized	Primary secondary
	Diabetic ulceration		
	Peripheral vascular disease		
	Burn injuries		
Autoimmune disorders	Lupus erythematoses, Sjogren, Rheumatoid arthritis, Glomerulonephritis, Angiitis		
	Langerhan's histiocytosis		
Eye	Optic neuritis		
	Blunt and penetrating injuries, Infections, Sarcoid, Sickle C disease, Retinal detachment, Temporal arteritis		
	Retinal ischemia, Macular degeneration, Retinitis pigmentosa, Arteriosclerotic retinopathy, Hypertensive retinopathy, Retinal artery blockage, Retinal vein blockage, Hypotension, Diabetic retinopathy, and Macular edema		
Embryonic and fetal disorders	Asphyxia		
	Ischemia		
CNS	Chronic fatigue syndrome, acute and chronic hyposmolar and hyperosmolar syndromes, AIDS Dementia, Electrocution		
	Encephalitis	Rabies, Herpes	
	Meningitis		
	Subdural hematoma		
	Nicotine addiction		

<i>Cell, tissue or organ</i>	<i>Dysfunction or pathology</i>	<i>Condition or disease</i>	<i>Type</i>
	Drug abuse and withdrawal	Cocaine, heroin, crack, marijuana, LSD, PCP, poly-drug abuse, ecstasy, opioids, sedative hypnotics, amphetamines, caffeine	
	Obsessive-compulsive disorders		
	Spinal stenosis, Transverse myelitis, Guillian Barre, Trauma, Nerve root compression, Tumoral compression, Heat stroke		
ENT	Tinnitus Meuniere's syndrome Hearing loss		
	Traumatic injury, barotraumas		
Kidney	Renal failure	Acute, chronic	Vascular/ischemic, interstitial disease, diabetic kidney disease, nephrotic syndromes, infections, injury, contrast-induced, chemotherapy-induced, CPB-induced, or preventive
	Henoch S. Purpura		
Striated muscle	Autoimmune disorders	Myasthenia gravis Dermatomyositis Polymyositis	
	Myopathies	Inherited metabolic, endocrine and toxic	
	Heat stroke		
	Crush injury		
	Rhabdomyolysis		
	Mitochondrial disease		
	Infection	Necrotizing fasciitis	
Sexual dysfunction	Central and peripheral (e.g. erectile dysfunction)	Impotence secondary to medication, (diabetes)	
Liver	Hepatitis	Viral, bacterial, parasitic	
	Ischemic disease		
	Cirrhosis, fatty liver		
	Infiltrative/metabolic diseases		
Gastrointestinal	Ischemic bowel disease		
	Inflammatory bowel disease		
	Necrotizing enterocolitis		
Organ transplantation	Treatment of donor and recipient		
Reproductive tract	Infertility	Vascular Autoimmune Uterine abnormalities Implantation disorders	
Endocrine	Glandular hyper- and hypofunction		

As mentioned above, these diseases, disorders or conditions are merely illustrative of the range of benefits provided by the recombinant tissue protective cytokines of the invention. Accordingly, this invention generally provides therapeutic or prophylactic treatment of the consequences of mechanical trauma or of human diseases. Therapeutic or prophylactic treatment for diseases, disorders or conditions of the CNS and/or peripheral nervous system are preferred. Therapeutic or prophylactic treatment for diseases, disorders or conditions which have a psychiatric component is provided. Therapeutic or prophylactic treatment for diseases, disorders or conditions including, but not limited to, those having an ophthalmic, cardiovascular, cardiopulmonary, respiratory, kidney, urinary, reproductive, gastrointestinal, endocrine, or metabolic component is provided.

In one embodiment, such a pharmaceutical composition of a recombinant tissue protective cytokine may be administered systemically to protect or enhance the target cells, tissue, or organ. Such administration may be parenterally, via inhalation, or transmucosally, *e.g.*, orally, nasally, rectally, intravaginally, sublingually, submucosally or transdermally. Preferably, administration is parenteral, *e.g.*, via intravenous or intraperitoneal injection, and also including, but is not limited to, intra-arterial, intramuscular, intradermal and subcutaneous administration.

For other routes of administration, such as by use of a perfusate, injection into an organ, or other local administration, a pharmaceutical composition will be provided which results in similar levels of a recombinant tissue protective cytokine as described above. A level of about 0.01pM –30 nM is preferred.

The pharmaceutical compositions of the invention may comprise a therapeutically effective amount of a compound, and a pharmaceutically acceptable carrier. In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized foreign pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as saline solutions in water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. A saline solution is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical

excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents.

5 These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. The compounds of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with free amino groups such as those derived from

10 hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with free carboxyl groups such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective

15 amount of the compound, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

Pharmaceutical compositions adapted for oral administration may be provided as capsules or tablets; as powders or granules; as solutions, syrups or suspensions (in aqueous

20 or non-aqueous liquids); as edible foams or whips; or as emulsions. Tablets or hard gelatine capsules may comprise lactose, starch or derivatives thereof, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, stearic acid or salts thereof. Soft gelatine capsules may comprise vegetable oils, waxes, fats, semi-solid, or liquid polyols etc. Solutions and syrups may comprise water, polyols, and sugars.

25 An active agent intended for oral administration may be coated with or admixed with a material that delays disintegration and/or absorption of the active agent in the gastrointestinal tract (*e.g.*, glyceryl monostearate or glyceryl distearate may be used). Thus, the sustained release of an active agent may be achieved over many hours and, if necessary, the active agent can be protected from being degraded within the stomach. Pharmaceutical

30 compositions for oral administration may be formulated to facilitate release of an active agent at a particular gastrointestinal location due to specific pH or enzymatic conditions.

Pharmaceutical compositions adapted for transdermal administration may be provided as discrete patches intended to remain in intimate contact with the epidermis of the

recipient for a prolonged period of time. Pharmaceutical compositions adapted for topical administration may be provided as ointments, creams, suspensions, lotions, powders, solutions, pastes, gels, sprays, aerosols or oils. For topical administration to the skin, mouth, eye or other external tissues a topical ointment or cream is preferably used. When
5 formulated in an ointment, the active ingredient may be employed with either a paraffinic or a water-miscible ointment base. Alternatively, the active ingredient may be formulated in a cream with an oil-in-water base or a water-in-oil base. Pharmaceutical compositions adapted for topical administration to the eye include eye drops. In these compositions, the active ingredient can be dissolved or suspended in a suitable carrier, *e.g.*, in an aqueous
10 solvent. Pharmaceutical compositions adapted for topical administration in the mouth include lozenges, pastilles, and mouthwashes.

Pharmaceutical compositions adapted for nasal and pulmonary administration may comprise solid carriers such as powders (preferably having a particle size in the range of 20 to 500 microns). Powders can be administered in the manner in which snuff is taken, *i.e.*,
15 by rapid inhalation through the nose from a container of powder held close to the nose. Alternatively, compositions adopted for nasal administration may comprise liquid carriers, *e.g.*, nasal sprays or nasal drops. Alternatively, inhalation directly into the lungs may be accomplished by inhalation deeply or installation through a mouthpiece into the oropharynx. These compositions may comprise aqueous or oil solutions of the active
20 ingredient. Compositions for administration by inhalation may be supplied in specially adapted devices including, but not limited to, pressurized aerosols, nebulizers or insufflators, which can be constructed so as to provide predetermined dosages of the active ingredient. In a preferred embodiment, pharmaceutical compositions of the invention are administered into the nasal cavity directly or into the lungs via the nasal cavity or
25 oropharynx.

Pharmaceutical compositions adapted for rectal administration may be provided as suppositories or enemas. Pharmaceutical compositions adapted for vaginal administration may be provided as pessaries, tampons, creams, gels, pastes, foams or spray formulations.

Pharmaceutical compositions adapted for parenteral administration include aqueous
30 and non-aqueous sterile injectable solutions or suspensions, which may contain antioxidants, buffers, bacteriostats, and solutes that render the compositions substantially isotonic with the blood of an intended recipient. Other components that may be present in such compositions include water, alcohols, polyols, glycerine and vegetable oils, for

example. Compositions adapted for parenteral administration may be presented in unit-dose or multi-dose containers, for example sealed ampules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of a sterile liquid carrier, e.g., sterile saline solution for injections, immediately prior to use. Extemporaneous
5 injection solutions and suspensions may be prepared from sterile powders, granules, and tablets. In one embodiment, an autoinjector comprising an injectable solution of a recombinant tissue protective cytokine may be provided for emergency use by ambulances, emergency rooms, and battlefield situations, and even for self-administration in a domestic setting, particularly where the possibility of traumatic amputation may occur, such as by
10 imprudent use of a lawn mower. The likelihood that cells and tissues in a severed foot or toe will survive after reattachment may be increased by administering a recombinant tissue protective cytokine to multiple sites in the severed part as soon as practicable, even before the arrival of medical personnel on site, or arrival of the afflicted individual with severed toe at the emergency room.

15 In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lidocaine to ease pain at the site of the
20 injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water-free concentrate in a hermetically-sealed container such as an ampule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the
25 composition is administered by injection, an ampule of sterile saline can be provided so that the ingredients may be mixed prior to administration.

Suppositories generally contain active ingredient in the range of 0.5% to 10% by weight; oral formulations preferably contain 10% to 95% active ingredient.

A perfusate composition may be provided for use in transplanted organ baths, for *in*
30 *situ* perfusion, or for administration to the vasculature of an organ donor prior to organ harvesting. Such pharmaceutical compositions may comprise levels of a recombinant tissue protective cytokine or a form of a recombinant tissue protective cytokine not suitable for acute or chronic, local or systemic administration to an individual, but will serve the

functions intended herein in a cadaver, organ bath, organ perfusate, or *in situ* perfusate prior to removing or reducing the levels of the recombinant tissue protective cytokine contained therein before exposing or returning the treated organ or tissue to regular circulation.

5 The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use, or sale for human administration.

10 In another embodiment, for example, a recombinant tissue protective cytokine can be delivered in a controlled-release system. For example, the polypeptide may be administered using intravenous infusion, an implantable osmotic pump, a transdermal patch, liposomes, or other modes of administration. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, 1987, CRC Crit. Ref. Biomed. Eng. 14:201; Buchwald *et al.*, 1980, 15 Surgery 88:507; Saudek *et al.*, 1989, N. Engl. J. Med. 321:574). In another embodiment, the compound can be delivered in a vesicle, in particular a liposome (see Langer, Science 249:1527-1533 (1990); Treat *et al.*, in Liposomes in the Therapy of Infectious Disease and Cancer, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989); WO 91/04014; U.S. Patent No. 4,704,355; Lopez-Berestein, *ibid.*, pp. 317-327; see generally 20 *ibid.*). In another embodiment, polymeric materials can be used (see Medical Applications of Controlled Release, Langer and Wise (eds.), CRC Press: Boca Raton, Florida, 1974; Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley: New York (1984); Ranger and Peppas, J. Macromol. Sci. Rev. Macromol. Chem. 23:61, 1953; see also Levy *et al.*, 1985, Science 228:190; During *et al.*, 1989, Ann. 25 Neurol. 25:351; Howard *et al.*, 1989, J. Neurosurg. 71:105).

 In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, i.e., the target cells, tissue or organ, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, pp. 115-138 in Medical Applications of Controlled Release, vol. 2, *supra*, 1984). Other controlled release systems are discussed in the review 30 by Langer (1990, Science 249:1527-1533).

 In another embodiment, a recombinant tissue protective cytokine, as properly formulated, can be administered by nasal, oral, rectal, vaginal, or sublingual administration.

In a specific embodiment, it may be desirable to administer the recombinant tissue protective cytokine compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, *e.g.*, in conjunction with a wound dressing after surgery, by
5 injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as silastic membranes, or fibers.

Selection of the preferred effective dose will be determined by a skilled artisan based upon considering several factors which will be known to one of ordinary skill in the
10 art. Such factors include the particular form of recombinant tissue protective cytokine, and its pharmacokinetic parameters such as bioavailability, metabolism, half-life, etc., which will have been established during the usual development procedures typically employed in obtaining regulatory approval for a pharmaceutical compound. Further factors in considering the dose include the condition or disease to be treated or the benefit to be
15 achieved in a normal individual, the body mass of the patient, the route of administration, whether administration is acute or chronic, concomitant medications, and other factors well known to affect the efficacy of administered pharmaceutical agents. Thus the precise dosage should be decided according to the judgment of the practitioner and each patient's circumstances, *e.g.*, depending upon the condition and the immune status of the individual
20 patient, and according to standard clinical techniques.

In another aspect of the invention, a perfusate or perfusion solution is provided for perfusion and storage of organs for transplant, the perfusion solution including an amount of a recombinant tissue protective cytokine effective to protect responsive cells and associated cells, tissues, or organs. Transplant includes, but is not limited to, xenotransplantation,
25 where a organ (including cells, tissue or other bodily part) is harvested from one donor and transplanted into a different recipient; and autotransplant, where the organ is taken from one part of a body and replaced at another, including bench surgical procedures, in which an organ may be removed, and while *ex vivo*, resected, repaired, or otherwise manipulated, such as for tumor removal, and then returned to the original location. In one embodiment,
30 the perfusion solution is the University of Wisconsin (UW) solution (U.S. Patent No. 4,798,824) which contains from about 1 to about 25 U/ml erythropoietin, 5% hydroxyethyl starch (having a molecular weight of from about 200,000 to about 300,000 and substantially free of ethylene glycol, ethylene chlorohydrin, sodium chloride and acetone); 25mM

KH₂PO₄; 3mM glutathione; 5mM adenosine; 10mM glucose; 10mM HEPES buffer; 5mM magnesium gluconate; 1.5mM CaCl₂; 105mM sodium gluconate; 200,000 units penicillin; 40 units insulin; 16mg Dexamethasone; 12mg Phenol Red; and has a pH of 7.4-7.5 and an osmolality of about 320 mOSm/l. The solution is used to maintain cadaveric kidneys and pancreases prior to transplant. Using the solution, preservation can be extended beyond the 30-hour limit recommended for cadaveric kidney preservation. This particular perfusate is merely illustrative of a number of such solutions that can be adapted for the present use by inclusion of an effective amount of a recombinant tissue protective cytokine. In a further embodiment, the perfusate solution contains from about 0.01pg/ml to about 400 ng/ml recombinant tissue protective cytokine, or from about 40 to about 300 ng/ml recombinant tissue protective cytokine. As mentioned above, any form of recombinant tissue protective cytokine can be used in this aspect of the invention.

While the preferred recipient of a recombinant tissue protective cytokine for the purposes herein throughout is a human, the methods herein apply equally to other mammals, particularly domesticated animals, livestock, companion and zoo animals. However, the invention is not so limiting and the benefits can be applied to any mammal.

5.6. THERAPEUTIC AND PREVENTATIVE USES OF RECOMBINANT TISSUE PROTECTIVE CYTOKINES

As noted in Example 1 below, the presence of erythropoietin receptors in the brain capillary human endothelium indicates that the targets of the recombinant tissue protective cytokines of the invention are present in the human brain, and that the animal studies on these recombinant tissue protective cytokines of the invention are directly translatable to the treatment or prophylaxis of human beings.

In another aspect of the invention, methods and compositions for enhancing the viability of cells, tissues, or organs which are not isolated from the vasculature by an endothelial cell barrier are provided by exposing the cells, tissue or organs directly to a pharmaceutical composition comprising a recombinant tissue protective cytokine, or administering or contacting an recombinant tissue protective cytokine-containing pharmaceutical composition to the vasculature of the tissue or organ. Enhanced activity of responsive cells in the treated tissue or organ is responsible for the positive effects exerted.

As described above, the invention is based, in part, on the discovery that erythropoietin molecules can be transported from the luminal surface to the basement

membrane surface of endothelial cells of the capillaries of organs with endothelial cell tight junctions, including, for example, the brain, retina, and testis. Thus, responsive cells across the barrier are susceptible targets for the beneficial effects of a recombinant tissue protective cytokine, and others cell types or tissues or organs that contain and depend in whole or in part on responsive cells therein are targets for the methods of the invention. While not wishing to be bound by any particular theory, after transcytosis of a recombinant tissue protective cytokine, the recombinant tissue protective cytokine can interact with an erythropoietin receptor on an responsive cell, for example, neuronal, retinal, muscle, heart, lung, liver, kidney, small intestine, adrenal cortex, adrenal medulla, capillary endothelial, testes, ovary, pancreas, bone, skin, or endometrial cell, and receptor binding can initiate a signal transduction cascade resulting in the activation of a gene expression program within the responsive cell or tissue, resulting in the protection of the cell or tissue, or organ, from damage, such as by toxins, chemotherapeutic agents, radiation therapy, hypoxia, etc. Thus, methods for protecting responsive cell-containing tissue from injury or hypoxic stress, and enhancing the function of such tissue are described in detail herein below. As noted above, the methods of the invention are equally applicable to humans as well as to other animals.

In the practice of one embodiment of the invention, a mammalian patient is undergoing systemic chemotherapy for cancer treatment, including radiation therapy, which commonly has adverse effects such as nerve, lung, heart, ovarian, or testicular damage. Administration of a pharmaceutical composition comprising a recombinant tissue protective cytokine as described above is performed prior to and during chemotherapy and/or radiation therapy, to protect various tissues and organs from damage by the chemotherapeutic agent, such as to protect the testes. Treatment may be continued until circulating levels of the chemotherapeutic agent have fallen below a level of potential danger to the mammalian body.

In the practice of another embodiment of the invention, various organs were planned to be harvested from a victim of an automobile accident for transplant into a number of recipients, some of which required transport for an extended distance and period of time. Prior to organ harvesting, the victim was infused with a pharmaceutical composition comprising a recombinant tissue protective cytokine as described herein. Harvested organs for shipment were perfused with a perfusate containing a recombinant tissue protective cytokine as described herein, and stored in a bath comprising recombinant tissue protective cytokine. Certain organs were continuously perfused with a pulsatile perfusion device,

utilizing a perfusate containing a recombinant tissue protective cytokine in accordance with the present invention. Minimal deterioration of organ function occurred during the transport and upon implant and reperfusion of the organs *in situ*.

5 In another embodiment of the invention, a surgical procedure to repair a heart valve required temporary cardioplegia and arterial occlusion. Prior to surgery, the patient was infused with 4 μ g recombinant tissue protective cytokine per kg body weight. Such treatment prevented hypoxic ischemic cellular damage, particularly after reperfusion.

10 In another embodiment of the invention, in any surgical procedure, such as in cardiopulmonary bypass surgery, a recombinant tissue protective cytokine of the invention can be used. In one embodiment, administration of a pharmaceutical composition comprising a recombinant tissue protective cytokine as described above is performed prior to, during, and/or following the bypass procedure, to protect the function of brain, heart, and other organs.

15 In the foregoing examples in which a recombinant tissue protective cytokine of the invention is used for *ex-vivo* applications, or to treat responsive cells such as neuronal tissue, retinal tissue, heart, lung, liver, kidney, small intestine, adrenal cortex, adrenal medulla, capillary endothelial, testes, ovary, or endometrial cells or tissue, the invention provides a pharmaceutical composition in dosage unit form adapted for protection or enhancement of responsive cells, tissues, or organs distal to the vasculature which
20 comprises, per dosage unit, an effective non-toxic amount within the range from about 0.01 pg to 5 mg, 1 pg to 5 mg, 500pg to 5 mg, 1 ng to 5 mg, 500 ng to 5 mg, 1 μ g to 5 mg, 500 μ g to 5mg, or 1 mg to 5 mg of a recombinant tissue protective cytokine and a pharmaceutically acceptable carrier. In a preferred embodiment, the amount of recombinant tissue protective cytokine is within the range from about 1 ng to 5mg. In a preferred
25 embodiment, the recombinant tissue protective cytokine of the aforementioned composition is non-erythropoietic.

In a further aspect of the invention, EPO administration was found to restore cognitive function in animals having undergone brain trauma. Recombinant tissue protective cytokines of the invention would be expected to have the same cellular protective
30 effects as EPO. After a delay of either 5 days or 30 days, EPO was still able to restore function as compared to sham-treated animals, indicating the ability of a EPO to regenerate or restore brain activity. Thus, the invention is also directed to the use of a recombinant

tissue protective cytokine for the preparation of a pharmaceutical composition for the treatment of brain trauma and other cognitive dysfunctions, including treatment well after the injury (e.g. three days, five days, a week, a month, or longer). The invention is also directed to a method for the treatment of cognitive dysfunction following injury by
5 administering an effective amount of a recombinant tissue protective cytokine. Any recombinant tissue protective cytokine as described herein may be used for this aspect of the invention.

Furthermore, this restorative aspect of the invention is directed to the use of any of the recombinant tissue protective cytokines herein for the preparation of a pharmaceutical
10 composition for the restoration of cellular, tissue, or organ dysfunction, wherein treatment is initiated after, and well after, the initial insult responsible for the dysfunction. Moreover, treatment using recombinant tissue protective cytokines of the invention can span the course of the disease or condition during the acute phase as well as a chronic phase.

In the instance wherein a recombinant tissue protective cytokine of the invention has
15 erythropoietic activity, in a preferred embodiment, recombinant tissue protective cytokine may be administered systemically at a dosage between about 0.01 pg and about 100 $\mu\text{g/kg}$ body weight, preferably about 1-50 $\mu\text{g/kg}$ -body weight, most preferably about 5-30 $\mu\text{g/kg}$ -body weight, per administration. This effective dose should be sufficient to achieve serum levels of recombinant tissue protective cytokine greater than about 10,000, 15,000, or
20 20,000 mU/ml of serum after recombinant tissue protective cytokine administration. Such serum levels may be achieved at about 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 hours post-administration. Such dosages may be repeated as necessary. For example, administration may be repeated daily, as long as clinically necessary, or after an appropriate interval, e.g., every 1 to 12 weeks, but preferably, every 1 to 3 weeks. In one embodiment, the effective
25 amount of recombinant tissue protective cytokine and a pharmaceutically acceptable carrier may be packaged in a single dose vial or other container. In another embodiment, a recombinant tissue protective cytokine useful for the purposes herein is nonerythropoietic, i.e., it is capable of exerting the activities described herein without causing an increase in hemoglobin concentration or hematocrit. Such a non-erythropoietic form of a recombinant
30 tissue protective cytokine is preferred in instances wherein the methods of the present invention are intended to be provided chronically. In another embodiment, a recombinant tissue protective cytokine is given at a dose greater than that necessary to maximally stimulate erythropoiesis. As noted above, a recombinant tissue protective cytokine of the

invention does not necessarily have erythropoietic activity, and therefore the above dosages expressed in units are merely exemplary for recombinant tissue protective cytokines; herein above molar equivalents for dosages are provided which are applicable to any recombinant tissue protective cytokine.

5 The present invention is further directed to a method for facilitating the transport of a molecule across an endothelial cell barrier in a mammal by administering a composition which comprises the particular molecule in association with a recombinant tissue protective cytokine as described herein above. As described above, tight junctions between endothelial cells in certain organs in the body create a barrier to the entry of certain
10 molecules. For treatment of various conditions within the barriered organ, means for facilitating passage of pharmaceutical agents is desired. A recombinant tissue protective cytokine of the invention is useful as a carrier for delivering other molecules across the blood-brain and other similar barriers. A composition comprising a molecule desirous of crossing the barrier with a recombinant tissue protective cytokine is prepared, and
15 peripheral administration of the composition results in the transcytosis of the composition across the barrier. The association between the molecule to be transported across the barrier and the recombinant tissue protective cytokine may be a labile covalent bond, in which case the molecule is released from association with the recombinant tissue protective cytokine after crossing the barrier. If the desired pharmacological activity of the molecule is
20 maintained or unaffected by association with the recombinant tissue protective cytokine, such a complex can be administered.

 The skilled artisan will be aware of various means for associating molecules with a recombinant tissue protective cytokine of the invention and the other agents described above, by covalent, non-covalent, and other means; furthermore, evaluation of the efficacy
25 of the composition can be readily determined in an experimental system. Association of molecules with a recombinant tissue protective cytokine may be achieved by any number of means, including labile, covalent binding, cross-linking, etc. Biotin/avidin interactions may be employed. As mentioned above, a hybrid molecule may be prepared by recombinant or synthetic means, for example, which includes both the domain of the molecule with desired
30 pharmacological activity and the domain responsible for erythropoietin receptor activity modulation.

 A molecule may be conjugated to a recombinant tissue protective cytokine through a polyfunctional molecule, i.e., a polyfunctional crosslinker. As used herein, the term

"polyfunctional molecule" encompasses molecules having one functional group that can react more than one time in succession, such as formaldehyde, as well as molecules with more than one reactive group. As used herein, the term "reactive group" refers to a functional group on the crosslinker that reacts with a functional group on a molecule (e.g., peptide, protein, carbohydrate, nucleic acid, particularly a hormone, antibiotic, or anti-cancer agent to be delivered across an endothelial cell barrier) so as to form a covalent bond between the cross-linker and that molecule. The term "functional group" retains its standard meaning in organic chemistry. The polyfunctional molecules which can be used are preferably biocompatible linkers, i.e., they are noncarcinogenic, nontoxic, and substantially non-immunogenic *in vivo*. Polyfunctional cross-linkers such as those known in the art and described herein can be readily tested in animal models to determine their biocompatibility. The polyfunctional molecule is preferably bifunctional. As used herein, the term "bifunctional molecule" refers to a molecule with two reactive groups. The bifunctional molecule may be heterobifunctional or homobifunctional. A heterobifunctional cross-linker allows for vectorial conjugation. It is particularly preferred for the polyfunctional molecule to be sufficiently soluble in water for the cross-linking reactions to occur in aqueous solutions such as in aqueous solutions buffered at pH 6 to 8, and for the resulting conjugate to remain water soluble for more effective bio-distribution. Typically, the polyfunctional molecule covalently bonds with an amino or a sulfhydryl functional group. However, polyfunctional molecules reactive with other functional groups, such as carboxylic acids or hydroxyl groups, are contemplated in the present invention.

The homobifunctional molecules have at least two reactive functional groups, which are the same. The reactive functional groups on a homobifunctional molecule include, for example, aldehyde groups and active ester groups. Homobifunctional molecules having aldehyde groups include, for example, glutaraldehyde and subaraldehyde. The use of glutaraldehyde as a cross-linking agent was disclosed by Poznansky *et al.*, Science 223, 1304-1306 (1984). Homobifunctional molecules having at least two active ester units include esters of dicarboxylic acids and N-hydroxysuccinimide. Some examples of such N-succinimidyl esters include disuccinimidyl suberate and dithio-bis-(succinimidyl propionate), and their soluble bis-sulfonic acid and bis-sulfonate salts such as their sodium and potassium salts. These homobifunctional reagents are available from numerous commercial sources (Pierce, Rockford, Illinois).

The heterobifunctional molecules have at least two different reactive groups. The reactive groups react with different functional groups, *e.g.*, present on the erythropoietin mutein and the molecule. These two different functional groups that react with the reactive group on the heterobifunctional cross-linker are usually an amino group, *e.g.*, the epsilon amino group of lysine; a sulfhydryl group, *e.g.*, the thiol group of cysteine; a carboxylic acid, *e.g.*, the carboxylate on aspartic acid; or a hydroxyl group, *e.g.*, the hydroxyl group on serine. Of course, recombinant tissue protective cytokines of the invention may be lacking a particular amino acid residue that would facilitate cross-linking of native erythropoietin, but one of skill in the art will be aware of the available residue moieties in a mutein of the invention and cross-link accordingly.

Moreover, the various recombinant tissue protective cytokine molecules of the invention may not have suitable reactive groups available for use with certain cross-linking agents; however, one of skill in the art will be amply aware of the choice of cross-linking agents based on the available groups for cross-linking in an erythropoietin of the invention.

When a reactive group of a heterobifunctional molecule forms a covalent bond with an amino group, the covalent bond will usually be an amido or imido bond. The reactive group that forms a covalent bond with an amino group may, for example, be an activated carboxylate group, a halocarbonyl group, or an ester group. The preferred halocarbonyl group is a chlorocarbonyl group. The ester groups are preferably reactive ester groups such as, for example, an N-hydroxy-succinimide ester group.

The other functional group typically is either a thiol group, a group capable of being converted into a thiol group, or a group that forms a covalent bond with a thiol group. The covalent bond will usually be a thioether bond or a disulfide. The reactive group that forms a covalent bond with a thiol group may, for example, be a double bond that reacts with thiol groups or an activated disulfide. A reactive group containing a double bond capable of reacting with a thiol group is the maleimido group, although others, such as acrylonitrile, are also possible. A reactive disulfide group may, for example, be a 2-pyridyldithio group or a 5,5'-dithio-bis-(2-nitrobenzoic acid) group. Some examples of heterobifunctional reagents containing reactive disulfide bonds include N-succinimidyl 3-(2-pyridyldithio)propionate (Carlsson, *et al.*, 1978, *Biochem J.*, 173:723-737), sodium S-4-succinimidylloxycarbonyl-alpha-methylbenzylthiosulfate, and 4-succinimidylloxycarbonyl-alpha-methyl-(2-pyridyldithio)toluene. N-succinimidyl 3-(2-pyridyldithio)propionate is preferred. Some examples of heterobifunctional reagents comprising reactive groups

having a double bond that reacts with a thiol group include succinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate and succinimidyl m-maleimidobenzoate.

Other heterobifunctional molecules include succinimidyl 3-(maleimido)propionate, sulfosuccinimidyl 4-(p-maleimido-phenyl)butyrate, sulfosuccinimidyl 4-(N-maleimidomethyl-cyclohexane)-1-carboxylate, maleimidobenzoyl-N-hydroxy-succinimide ester. The sodium sulfonate salt of succinimidyl m-maleimidobenzoate is preferred. Many of the above-mentioned heterobifunctional reagents and their sulfonate salts are available from Pierce Chemical Co., Rockford, Illinois USA.

The need for the above-described conjugated to be reversible or labile may be readily determined by the skilled artisan. A conjugate may be tested *in vitro* for both the recombinant tissue protective cytokine activity, and for the desirable pharmacological activity. If the conjugate retains both properties, its suitability may then be tested *in vivo*. If the conjugated molecule requires separation from the recombinant tissue protective cytokine for activity, a labile bond or reversible association with the recombinant tissue protective cytokine will be preferable. The lability characteristics may also be tested using standard *in vitro* procedures before *in vivo* testing.

Additional information regarding how to make and use these as well as other polyfunctional reagents may be obtained from the following publications or others available in the art:

1. Carlsson, J. *et al.*, 1978, Biochem. J. 173:723-737.
2. Cumber, J.A. *et al.*, 1985, Methods in Enzymology 112:207-224.
3. Jue, R. *et al.*, 1978, Biochem 17:5399-5405.
4. Sun, T.T. *et al.*, 1974, Biochem. 13:2334-2340.
5. Blattler, W.A. *et al.*, 1985, Biochem. 24:1517-152.
6. Liu, F.T. *et al.*, 1979, Biochem. 18:690-697.
7. Youle, R.J. and Neville, D.M. Jr., 1980, Proc. Natl. Acad. Sci. U.S.A. 77:5483-5486.
8. Lerner, R.A. *et al.*, 1981, Proc. Natl. Acad. Sci. U.S.A. 78:3403-

3407.

9. Jung, S.M. and Moroi, M., 1983, *Biochem. Biophys. Acta* 761:162.
10. Caulfield, M.P. *et al.*, 1984, *Biochem.* 81:7772-7776.
11. Staros, J.V., 1982, *Biochem.* 21:3950-3955.
12. Yoshitake, S. *et al.*, 1979, *Eur. J. Biochem.* 101:395-399.
13. Yoshitake, S. *et al.*, 1982, *J. Biochem.* 92:1413-1424.
14. Pilch, P.F. and Czech, M.P., 1979, *J. Biol. Chem.* 254:3375-3381.
15. Novick, D. *et al.*, 1987, *J. Biol. Chem.* 262:8483-8487.
16. Lomant, A.J. and Fairbanks, G., 1976, *J. Mol. Biol.* 104:243-261.
17. Hamada, H. and Tsuruo, T., 1987, *Anal. Biochem.* 160:483-488.
18. Hashida, S. *et al.*, 1984, *J. Applied Biochem.* 6:56-63.

Additionally, methods of cross-linking are reviewed by Means and Feeney, 1990, *Bioconjugate Chem.* 1:2-12.

Barriers which are crossed by the above-described methods and compositions of the present invention include, but are not limited to, the blood-brain barrier, the blood-eye
5 barrier, the blood-testis barrier, the blood-ovary barrier, the blood-heart barrier, the blood-kidney barrier, and the blood-uterus barrier.

Candidate molecules for transport across an endothelial cell barrier include, for example, hormones, such as growth hormone, neurotrophic factors, antibiotics, antivirals, or antifungals such as those normally excluded from the brain and other barriered organs,
10 peptide radiopharmaceuticals, antisense drugs, antibodies and antivirals against biologically-active agents, pharmaceuticals, and anti-cancer agents. Non-limiting examples of such molecules include hormones such as growth hormone, nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), ciliary neurotrophic factor (CNTF), basic fibroblast growth factor (bFGF), transforming growth factor β 1 (TGF β 1), transforming
15 growth factor β 2 (TGF β 2), transforming growth factor β 3 (TGF β 3), interleukin 1,

interleukin 2, interleukin 3, and interleukin 6, AZT, antibodies against tumor necrosis factor, and immunosuppressive agents such as cyclosporin. Additionally, dyes or markers may be attached to erythropoietin or one of the tissue protective cytokines of the present invention in order to visualize cells, tissues, or organs within the brain and other barriered
5 organs for diagnostic purposes. As an example, a marker used to visualize plaque within the brain could be attached to erythropoietin or a tissue protective cytokine in order to determine the progression of Alzheimers disease within a patient.

The present invention is also directed to a composition comprising a molecule to be transported via transcytosis across a endothelial cell tight junction barrier and a recombinant
10 tissue protective cytokine as described above. The invention is further directed to the use of a conjugate between a molecule and a recombinant tissue protective cytokine as described above for the preparation of a pharmaceutical composition for the delivery of the molecule across a barrier as described above.

The present invention may be better understood by reference to the following non-
15 limiting examples, which are provided as exemplary of the invention. The following examples are presented in order to more fully illustrate the preferred embodiments of the invention. They should in no way be construed, however, as limiting the broad scope of the invention.

6. EXAMPLES

20 6.1. EXAMPLE 1: DISTRIBUTION OF ERYTHROPOIETIN RECEPTOR IN HUMAN BRAIN

Normal human brain removed during surgical procedures (e.g., to provide tumor-free margins in tumor resections) were immediately fixed in 5% acrolein in 0.1 M phosphate buffer (pH 7.4) for 3 h. Sections were cut with a vibrating microtome at 40
25 micrometer thickness. Immunohistochemical staining was performed using free-floating sections and the indirect antibody peroxidase-antiperoxidase method using a 1:500 dilution of erythropoietin receptor antiserum (obtained from Santa Cruz Biotechnology). Endogenous peroxidase activity was quenched by pretreatment of tissue sections with hydrogen peroxide (3% in methanol for 30 min). Tissue controls were also carried out by
30 primary antibody omission and by using the appropriate blocking peptide (from Santa Cruz Biotech.) to confirm that staining was specific for erythropoietin (EPO) receptor.

Figure 1 shows capillaries of the human brain express very high levels of EPO receptor, as determined by immunohistochemistry using specific anti-EPO receptor antibodies. This provides a mechanism whereby EPO is able to penetrate into the brain from the systemic circulation, in spite of the blood brain barrier.

5 **Figure 2** shows the EPO receptor is densely localized within and around capillaries forming the blood brain barrier in the human brain.

A similar protocol as for **Figures 1 & 2** was performed for **Figure 3**, except that 10 micrometer sections were cut from paraffin, the embedded sections fixed by immersion in 4% paraformaldehyde. **Figure 3** shows that there is a high density of EPO receptor at the
10 luminal and anti-luminal surfaces of human brain capillaries, forming the anatomical basis for transport of EPO from the circulation into the brain.

Figure 4 was obtained following a similar protocol as in **Figure 3** except that the tissue was sectioned on an ultramicrotome for electron microscopy and the secondary antibody was labeled with colloidal gold particles. This figure shows that EPO receptor is
15 found upon the endothelial surface (*), within cytoplasmic vesicles (arrows) and upon glial endfeet (+) in human brain, providing the anatomical basis for transport of EPO from within the circulation into the brain.

6.2. **EXAMPLE 2: ERYTHROPOIETIN CROSSES THE BLOOD-CEREBROSPINAL FLUID TIGHT BARRIER**

20 Adult male Sprague-Dawley rats were anesthetized and administered recombinant human erythropoietin intraperitoneally at 5000 U/kg body weight. Cerebrospinal fluid was sampled from the cisterna magna at 30 minute intervals up to 4 hrs and the erythropoietin concentration determined using a sensitive and specific enzyme-linked immunoassay. As illustrated in **Figure 5**, the baseline erythropoietin concentration in CSF is 8 mU/ml. After a
25 delay of several hours, the levels of erythropoietin measured in the CSF begin to rise and by 2.5 hours and later are significantly different from the baseline concentration at the $p < 0.01$ level. The peak level of about 100 mU/ml is within the range known to exert protective effects *in vitro* (0.1 to 100 mU/ml). The time to peak occurs at about 3.5 hrs, which is delayed significantly from the peak serum levels which occur at less than 1 hr. The results
30 of this experiment illustrate that significant levels of erythropoietin can be accomplished across a tight cellular junction by bolus parenteral administration of erythropoietin at appropriate concentrations.

6.3. EXAMPLE 3: RECOMBINANT TISSUE PROTECTIVE CYTOKINE

The following human erythropoietin constructs were made using the following procedures. The cDNA for the human erythropoietin was cloned by PCR from human brain cDNA by using primers based on the published human cDNA sequence (accession number
 5 NM_000799). The primers were designed to introduce a Xho I site in the 5' end and a Xba I site in the 3' end of the cDNA. The primer sequences are:

HEPO-5-Xho I 5'-AGCTCTCGAGGCGCGGAGATGGGGGTGCACGAATG-3'
 (SEQ. ID. 8)

HEPO-3-Xba I 5'-ATGCTCTAGACACACCTGGTCATCTGTCCCCTGTCC-3'
 10 (SEQ. ID. 9).

The PCR product was cloned between the Xho I and Xba I sites in pCiNeo mammalian expression vector (Promega). The clones were sequenced and the sequence was verified to match the sequence in NM_000799 with the exception of a single base. Base 418 in the coding sequence (starting the numbering from the ATG) was C instead of
 15 G, changing amino acid 140 in the full length EPO sequence starting from the first methionine from Arg to Gly. This is however, normal sequence variation from the original sequence and present in most forms of erythropoietin.

The coding sequence from the erythropoietin cDNA is below:

ATGGGGGTGCACGAATGTCCTGCCTGGCTGTGGCTTCTCCTGTCCCCTGCT
 20 GTCGCTCCCTCTGGGCCTCCCAGTCCTGGGCGCCCCACCGCCTCATCTGTGA
 CAGCCGAGTCCTGGAGAGGTACCTCTTGGAGGCCAAGGAGGCCGAGAATATCA
 CGACGGGCTGTGCTGAACACTGCAGCTTGAATGAGAATATCACTGTCCCAGACA
 CCAAAGTTAATTTCTATGCCTGGAAGAGGATGGAGGTCGGGCAGCAGGCCGTA
 GAAGTCTGGCAGGGCCTGGCCCTGCTGTGCGAAGCTGTCCTGCGGGGCCAGGC
 25 CCTGTTGGTCAACTCTTCCCAGCCGTGGGAGCCCCTGCACTGCATGTGGATAAA
 GCCGTCAGTGGCCTTCGCAGCCTCACCCTCTGCTTCGGGCTCTGGGAGCCCAG
 AAGGAAGCCATCTCCCCTCCAGATGCGGCCTCAGCTGCTCCACTCCGAACAATC
 ACTGCTGACACTTTCGCAAACCTCTTCCGAGTCTACTCCAATTTCTCCGGGGAA
 AGCTGAAGCTGTACACAGGGGAGGCCTGCAGGACAGGGGACAGATGA (SEQ ID
 30 NO: 7).

This cDNA codes for the full length amino acid sequence of erythropoietin, which is below

MGVHECPAWLWLLSLLSLPLGLPVLGAPPRLICDSRVLERYLLEAKEAENI
 TTGCAEHCSLNENITVPDTKVNFYAWKRMEVGQQAQAVEVWQGLALLSEAVLRGQA
 5 LLVNSSQPWEPLQLHVDKAVSGLRSLTLLRALGAQKEAISPPDAASAAPLRTITAD
 TFRKLFRVYSNFLRGKCLKLYTGEACRTGDR (SEQ ID NO: 10).

The first 27 amino acid residues of SEQ ID NO:10 comprise a leader sequence.

A 6xHis tag was introduced to the C-terminal end of the human EPO protein by designing a new oligonucleotide so that the 6 histidines would be joined to the Asp 192 in
 10 the full length sequence using the following oligonucleotide:

3'-6xhis- hEPO 5'-
 GGTCTAGATCAATGGTGATGGTGATGATGGTCCCCTGTCCTGCAGGCC-3' (SEQ
 ID NO:134)

The EPO cDNA was amplified by PCR using the HEPO-5-Xho I oligo and 6xHis-
 15 Tag oligo and cloned between the Xho I and Xba I sites in the pCiNeo mammalian expression vector. The insert was again sequenced and the sequence verified.

The mutations described above in section 5.2 to the human EPO cDNA sequence, with a C-terminal 6xHis tag, were introduced by oligo directed mutagenesis using the oligonucleotides described in this section. Mutant clones were sequenced to confirm the
 20 mutations.

Numerous methods for purification of the recombinant tissue protective cytokines of the invention may be used, including, but not limited to, the following protocol which was used in conjunction with the histidine tagged recombinantly expressed tissue protective cytokines of the invention. The recombinant cell (CHO-K1) supernatant (for example, resin
 25 from (Ni-CAM HC RESIN: High Capacity Nickel Chelate Affinity Matrix, Sigma, Cat no. N 3158)) was thoroughly resuspend with gentle inversion. Then, 100µl of the resin suspension (equivalent to 50µl of packed resin) was added to a microcentrifuge tube (1.7 ml size). The mixture was centrifuged at 8,000 rpm, at 4°C for 5 minutes to pellet the resin and then discard the supernatant. The microcentrifuge was Megafuge 1.0R (Heraeus
 30 Instruments). The mixture was washed twice with 1 ml of deionized water (0.2µm filtered) to remove the ethanol. The resin was resuspended in 500µl of equilibration buffer (50mM

sodium phosphate, pH 8.0, 0.3M NaCl, 10mM imidazole), and then transferred the mixture to a 50-ml conical tube. The microcentrifuge tube was rinsed with 500µl of equilibration buffer, and then added this amount to the mixture in the 50-ml conical tube. The mixture was centrifuged at 3,000 rpm, at 4°C for 5 minutes to pellet the resin. The supernatant was removed and discarded. The samples (CHO-KI supernatant) were centrifuged at 3,800 rpm, at 4°C for 5 minutes prior to binding. The cell supernatant was added to the resin. Sample addition buffer (50mM sodium phosphate, pH 8.0, 3M NaCl, 100mM imidazole) was added to 1X, and gently mixed in a rotating platform for 1 hour at 4°C. An example of such a platform used is Nutator (rotating platform) (Clay Adams Brand). The mixture was centrifuged at 3,000 rpm, at 4°C for 5 minutes. The supernatant was removed and saved for SDS-PAGE analysis and ELISA (unbound). The resin was resuspended in 500µl of wash buffer, and then the mixture was transferred to a microcentrifuge tube. The 50-ml conical tube was rinsed with 500µl of equilibration buffer, and then this amount was added to the mixture in the microcentrifuge tube. The resin suspension was then mixed in a rotating platform for 10 minutes at 4°C. The suspension was centrifuged at 8,000 rpm, at 4°C for 5 minutes (the first wash may be saved for ELISA). The resin was resuspended in 1 ml of wash buffer and the resin suspension was again then mixed in a rotating platform for 10 minutes at 4°C, to wash the resin one more time. The wash was discarded. Then, 75µl of elution buffer (50mM sodium phosphate, pH 8.0, 0.3M NaCl, 500mM imidazole) was added. The resin was mixed in a rotating platform for 10 minutes at 4°C. The mixture was centrifuged at 8,000 rpm, at 4°C for 5 minutes. The supernatant was removed and saved. The histidine tagged protein was in this fraction. To elute more protein, 75µl of elution buffer (50mM sodium phosphate, pH 8.0, 0.3M NaCl, 500mM imidazole) were again added. The resin was again mixed in a rotating platform for 10 minutes at 4°C. The mixture was again centrifuged at 8,000 rpm, at 4°C for 5 minutes. The eluted fractions were saved as a single pool or separate fractions.

Alternatively, the following procedure was used to isolate purified histidine-tagged cytokines. Conditioned media was collected and filtered through a 0.45 µm filter. A 50 ml aliquot was then applied to a 5 ml HiTrap chelating (Amersham biosciences) equilibrated with 20 mM sodium phosphate pH 7.4 and activated with 2.5 ml 100 mM NiSO₄. The column was washed with 20 mM sodium phosphate pH 7.4 and eluted with a gradient from 0 M to 0.5 M Imidazole in 20 mM sodium phosphate pH 7.4 over 25 column volumes. The flow was 5 ml/min and fraction size 5 ml.

Fractions were analyzed for the presence of recombinant tissue protective cytokine by SDS-PAGE and EPO ELISA. Positive fractions were pooled and dialyzed against 10 mM Tris pH 7.0. The pool was applied to a 1 ml HiTrap Q HP (Amersham biosciences) equilibrated with 10 mM Tris pH 7.0. After washing with equilibration buffer the sample
5 was eluted with a gradient of NaCl to 0.5 M over 10 column volumes at a flow of 1 ml/min. Fractions of 1 ml were collected and analyzed by SDS-PAGE, EPO ELISA and western blotting using antibodies against hexa-his tag (Anti-His₆, ROCHE). Fractions containing the recombinant tissue protective cytokine were pooled and concentrated using a centricon with a cut off size of 10 kDa (Amicon) to a final volume of 1-2 ml.

10 The final pool of recombinant tissue protective cytokine was analyzed by SDS-PAGE (NuPage 4-12%; Invitrogen) and visualized using NOVEX Colloidal Blue (Invitrogen) by the protocol recommended by the manufacturer. The purity of the recombinant tissue protective cytokine was judged from the resulting gel. Only one band corresponding to the glycosylated recombinant tissue protective cytokine was present in
15 each lane of the gel indicating a high purity of the isolated cytokine.

All the plasmids were transfected to either CHO-1 cells or COS-7 cells by using lipofectamine. Forty-eight to 72 hours post transfection media from the cells was collected. This media was tested for EPO by ELISA assay and used either directly or after purification in either the hematopoietic or neuronal assays.

20 Mutations K45D, S100E, and A30N/H32T to the human EPO cDNA sequence were introduced by oligo directed mutagenesis using following oligonucleotides:

HEPO-S100E-upper 5'-

CATGTGGATAAAGCCGTCGAGGGCCTTCGCAGCCTCACCACTCTG-3' (SEQ ID
NO: 11)

25 HEPO-S100E-lower 5'-

CAGAGTGGTGAGGCTGCGAAGGCCCTCGACGGCTTTATCCACATG-3' (SEQ ID
NO: 12)

HEPO-K45D-upper 5'-

GAGAATATCACTGTCCCAGACACCGACGTTAATTTCTATGCCTGG-3' (SEQ ID
30 NO: 13)

HEPO-K45D-lower 5'-

CCAGGCATAGAAATTAACGTCGGTGTCTGGGACAGTGATATTCTC-3' (SEQ ID NO: 14)

hEPO-A30N/H32T-upper 5'-

5 GAATATCACGACGGGCTGTAATGAAACCTGCAGCTTGAATGAG-3' (SEQ ID NO: 132)

hEPO-A30N/H32T-lower 5'-

CTCATTCAAGCTGCAGGTTTCATTACAGCCCGTCGTGATATTC-3' (SEQ ID NO: 133)

10 Oligonucleotide sequences used in oligo directed mutagenesis for the other erythropoietin muteins and recombinant tissue protective cytokines of the invention include:

For R150E mutein:

R150E-F GTCTACTCCAATTCCTCGAGGGAAAGCTGAAGCTG, (SEQ ID NO: 120)

15 R150E-R GCTTCAGCTTTCCTCGAGGAAATTGGAGTAGAC (SEQ ID NO: 121)

For R103E mutein:

R103E-F CCGTCAGTGGCCTTGAGAGCCTCACCACTCTG, (SEQ ID NO: 122)

20 R103E-R CAGAGTGGTGAGGCTCTCAAGGCCACTGACGG (SEQ ID NO: 123)

For R103E/L108S(103) combination mutein:

R103E-F CCGTCAGTGGCCTTGAGAGCCTCACCACTCTG (SEQ ID NO: 124)

R103E-R CAGAGTGGTGAGGCTCTCAAGGCCACTGACGG (SEQ ID NO: 125)

25 L108S(103)F CGCAGCCTCACCACTTCGCTTCGGGCTCTGG, (SEQ ID NO: 126)

L108S(103)R CCAGAGCCCGAAGCGAAGTGGTGAGGCTGCG (SEQ ID NO:
127)

For 44-49 deletion

5 d44-49F GAATATCACTGTCCCAGACGGTGGTGCCTGGAAGAGGATG, (SEQ
ID NO: 128)

d44-49R CATCCTCTTCCAGGCACCACCGTCTGGGACAGTGATATTC (SEQ
ID NO: 129)

10 For K20A mutein:

K20A-F TACCTCTTGGAGGCCGCGGAGGCCGAGAATATC, (SEQ ID NO:
130)

K20A-R GATATTCTCGGCCTCCGCGGCCTCCAAGAGGTA (SEQ ID NO: 131)

15 For K140A mutein:

K140A-F GCTGACACTTTCCGCGCACTCTTCCGAGTCTACTC, (SEQ ID NO:
132)

K140A-R GAGTAGACTCGGAAGAGTGCGCGGAAAGTGTACGC (SEQ ID
NO: 133)

20

For K152A mutein:

K152A-F ATTCCTCCGGGGAGCGCTGAAGCTGTACACAG, (SEQ ID NO:
134)

K152A-R CTGTGTACAGCTTCAGCGCTCCCCGGAGGAAAT (SEQ ID NO:
135)

For K154A mutein:

5 K154A-F CTCCGGGGAAAGCTGGCGCTGTACACAGGGGA, (SEQ ID NO:
136)

K154A-R TCCCCTGTGTACAGCGCCAGCTTTCCCCGGAG (SEQ ID NO: 137)

For K45A mutein:

10 K45A-F ACTGTCCCAGACACCGCAGTTAATTTCTATGCCTG, (SEQ ID NO:
138)

K45A-R CAGGCATAGAAATTAAGTGCAGGTGTCTGGGACAGT (SEQ ID NO:
139)

15 For K52A mutein:

K52A-F AGTTAATTTCTATGCCTGGGCGAGGATGGAGGTCG, (SEQ ID NO:
140)

K52A-R CGACCTCCATCCTCGCCCAGGCATAGAAATTAAGT (SEQ ID NO:
141)

20

For K97A mutein:

K97A-F TGCAGCTGCATGTGGATGCAGCCGTCAGTGGCC, (SEQ ID NO:
142)

K97A-R GGCCACTGACGGCTGCATCCACATGCAGCTGCA (SEQ ID NO:
25 143)

For K116A mutein:

K116A-F CTCTGGGAGCCCAGGCGGAAGCCATCTCCCCT, (SEQ ID NO: 144)

K116A-R AGGGGAGATGGCTTCCGCCTGGGCTCCCAGAG (SEQ ID NO: 145)

5 For K140A/K52A combination mutein:

K140A-F GCTGACACTTTCCGCGCACTCTTCCGAGTCTACTC, (SEQ ID NO: 146)

K140A-R GAGTAGACTCGGAAGAGTGCGCGGAAAGTGTCAGC (SEQ ID NO: 147)

10 K52A-F AGTTAATTTCTATGCCTGGGCGAGGATGGAGGTCG, (SEQ ID NO: 148)

K52A-R CGACCTCCATCCTCGCCCAGGCATAGAAATTA ACT (SEQ ID NO: 149)

15 For K140A/K52A/K45A combination mutein:

K140A-F GCTGACACTTTCCGCGCACTCTTCCGAGTCTACTC, (SEQ ID NO: 150)

K140A-R GAGTAGACTCGGAAGAGTGCGCGGAAAGTGTCAGC (SEQ ID NO: 151)

20 K52A-F AGTTAATTTCTATGCCTGGGCGAGGATGGAGGTCG, (SEQ ID NO: 152)

K52A-R CGACCTCCATCCTCGCCCAGGCATAGAAATTA ACT (SEQ ID NO: 153)

K45A-F ACTGTCCCAGACACCGCAGTTAATTTCTATGCCTG, (SEQ ID NO: 25 154)

K45A-R CAGGCATAGAAATTA ACTGCGGTGTCTGGGACAGT (SEQ ID NO: 155)

For K97A/K152A combination mutein:

K97A-F TGCAGCTGCATGTGGATGCAGCCGTCAGTGGCC, (SEQ ID NO:
156)
5 K97A-R GGCCACTGACGGCTGCATCCACATGCAGCTGCA (SEQ ID NO:
157)
K152A-F ATTCCTCCGGGGAGCGCTGAAGCTGTACACAG, (SEQ ID NO:
158)
K152A-R CTGTGTACAGCTTCAGCGCTCCCCGGAGGAAAT (SEQ ID NO:
10 159)

For K97A/K152A/K45A combination mutein:

K97A-F TGCAGCTGCATGTGGATGCAGCCGTCAGTGGCC, (SEQ ID NO:
160)
15 K97A-R GGCCACTGACGGCTGCATCCACATGCAGCTGCA (SEQ ID NO:
161)
K152A-F ATTCCTCCGGGGAGCGCTGAAGCTGTACACAG, (SEQ ID NO:
162)
K152A-R CTGTGTACAGCTTCAGCGCTCCCCGGAGGAAAT (SEQ ID NO:
20 163)
K45A-F ACTGTCCCAGACACCGCAGTTAATTTCTATGCCTG, (SEQ ID NO:
164)
K45A-R CAGGCATAGAAATTAAGTGCAGGTGTCTGGGACAGT (SEQ ID NO:
165)
25

For K97A/K152A/K45A/K52A combination mutein:

- 166) K97A-F TGCAGCTGCATGTGGATGCAGCCGTCAGTGGCC, (SEQ ID NO:
167) K97A-R GGCCACTGACGGCTGCATCCACATGCAGCTGCA (SEQ ID NO:
5 168) K152A-F ATTCCTCCGGGGAGCGCTGAAGCTGTACACAG, (SEQ ID NO:
169) K152A-R CTGTGTACAGCTTCAGCGCTCCCCGGAGGAAAT (SEQ ID NO:
10 170) K45A-F ACTGTCCCAGACACCGCAGTTAATTTCTATGCCTG, (SEQ ID NO:
171) K45A-R CAGGCATAGAAATTAAGTGCAGGTGTCTGGGACAGT (SEQ ID NO:
172) K52A-F AGTTAATTTCTATGCCTGGGCGAGGATGGAGGTCG, (SEQ ID NO:
15 173) K52A-R CGACCTCCATCCTCGCCCAGGCATAGAAATTAAGT (SEQ ID NO:

For K97A/K152A/K45A/K52A/K140A combination mutein:

- 20 174) K97A-F TGCAGCTGCATGTGGATGCAGCCGTCAGTGGCC, (SEQ ID NO:
175) K97A-R GGCCACTGACGGCTGCATCCACATGCAGCTGCA (SEQ ID NO:
176) K152A-F ATTCCTCCGGGGAGCGCTGAAGCTGTACACAG, (SEQ ID NO:
25 177) K152A-R CTGTGTACAGCTTCAGCGCTCCCCGGAGGAAAT (SEQ ID NO:

- 178) K45A-F ACTGTCCCAGACACCGCAGTTAATTTCTATGCCTG, (SEQ ID NO:
179) K45A-R CAGGCATAGAAATTAAGTGCAGGTGTCTGGGACAGT (SEQ ID NO:
5 180) K52A-F AGTTAATTTCTATGCCTGGGCGAGGATGGAGGTCG, (SEQ ID NO:
181) K52A-R CGACCTCCATCCTCGCCCAGGCATAGAAATTAAGT (SEQ ID NO:
10 182) K140A-F GCTGACACTTTCCGCGCACTCTTCCGAGTCTACTC, (SEQ ID NO:
K140A-R GAGTAGACTCGGAAGAGTGCAGCGGAAAGTGTCAGC (SEQ ID
NO: 183)
- For K97A/K152A/K45A/K52A/K140A/K154A combination mutein:
- 15 184) K97A-F TGCAGCTGCATGTGGATGCAGCCGTCAGTGGCC, (SEQ ID NO:
185) K97A-R GGCCACTGACGGCTGCATCCACATGCAGCTGCA (SEQ ID NO:
20 186) K152A-F ATTTCTCCGGGGAGCGCTGAAGCTGTACACAG, (SEQ ID NO:
187) K152A-R CTGTGTACAGCTTCAGCGCTCCCCGGAGGAAAT (SEQ ID NO:
188) K45A-F ACTGTCCCAGACACCGCAGTTAATTTCTATGCCTG, (SEQ ID NO:
25 189) K45A-R CAGGCATAGAAATTAAGTGCAGGTGTCTGGGACAGT (SEQ ID NO:

K52A-F AGTTAATTTCTATGCCTGGGCGAGGATGGAGGTCG, (SEQ ID NO:
190)

52A-R CGACCTCCATCCTCGCCCAGGCATAGAAATTA ACT (SEQ ID NO:
191)

5 K140A-F GCTGACACTTTCCGCGCACTCTTCCGAGTCTACTC, (SEQ ID NO:
192)

K140A-R GAGTAGACTCGGAAGAGTGCGCGGAAAGTGTCAGC (SEQ ID
NO: 193)

10 K154A(152)F CTCCGGGGAGCGCTGGCGCTGTACACAGGGGA, (SEQ ID
NO: 194)

154(152)R TCCCCTGTGTACAGCGCCAGCGCTCCCCGGAG (SEQ ID NO:
195)

For N24K/N38K/N83K combination mutein:

15 N24K-F CAAGGAGGCCGAGAAAATCACGACGGGCTGT, (SEQ ID NO: 196)

N24K-R ACAGCCCGTCGTGATTTTCTCGGCCTCCTTG (SEQ ID NO: 197)

N38K-F ACTGCAGCTTGAATGAGAAAATCACTGTCCCAGACAC, (SEQ ID
NO: 198)

20 N38K-R GTGTCTGGGACAGTGATTTTCTCATTCAAGCTGCAGT (SEQ ID
NO: 199)

N83K-F AGGCCCTGTTGGTCAAATCTTCCCAGCCGTG, (SEQ ID NO: 200)

N83K-R CACGGCTGGGAAGATTGACCAACAGGGCCT (SEQ ID NO: 201)

For K152W mutein:

25 K152W-F ATTCCTCCGGGGATGGCTGAAGCTGTACACAG, (SEQ ID NO:
202)

K152W-R CTGTGTACAGCTTCAGCCATCCCCGGAGGAAAT (SEQ ID NO: 203)

For R14A/Y15A combination mutein:

5 RY14AA-F AGCCGAGTCCTGGAGGCGGCCCTCTTGGAGGCCAA, (SEQ ID NO: 204)

RY14AA-R TTGGCCTCCAAGAGGGCCGCCTCCAGGACTCGGCT (SEQ ID NO: 205)

10 Y15A-F AGCCGAGTCCTGGAGAGGGCCCTCTTGGAGGCCAA (SEQ ID NO: 206)

Y15A-R TTGGCCTCCAAGAGGGCCCTCTCCAGGACTCGGCT (SEQ ID NO: 207)

The following are examples of constructs that were made: human EPO(hEPO)-6xHisTag-pCiNeo sequence (SEQ ID NO: 208); hEPO6xHisTag-A30N/H32T-pCiNeo (SEQ ID NO: 209); hEPO-6xHisTag-K45D-pCiNeo sequence (SEQ ID NO: 210); hEPO-6xHisTag-S100E-pCiNeo sequence (SEQ ID NO: 211); and hEPO-6xHisTag-K45D/S100E-pCiNeo sequence (SEQ ID NO: 212). The pCI-neo mammalian expression vector carries the human cytomegalovirus (CMV) immediate-early enhancer/promoter region to promote constitutive expression of cloned DNA inserts in mammalian cells.

20 These oligonucleotides were annealed to the original human erythropoietin cDNA clone in pCiNeo to introduce the mutations. Mutant clones were sequenced to confirm the mutations. All the plasmids were transfected to either CHO-1 cells or COS-7 cells by using lipofectamine. At 48 to 72 hours post-transfection media from the cells was collected. This media was tested for erythropoietin by ELISA assay and used either directly or after
25 purification in either the hematopoietic or neuronal assays.

Subsequently, both the K45D and S100E recombinant tissue protective cytokines were tested within a neuronal assay. Specifically, an *in vitro* neuroprotection assay using SK-N-SH neuroblastoma cells was used. SK-N-SH cells were plated at a density of 40,000 cells/well in 24 well plates for 24 hours. Then recombinant tissue protective cytokines were

added at a concentration of 3 nM for an additional 24 hours (Erythropoietin = commercial preparation; EPO = erythropoietin and recombinant tissue protective cytokines expressed in CHO cells; pure vector = cell supernatant from CHO cells transfected with vector without Epo construct). After this preincubation, cells were exposed to rotenone (5 μ M) for 4 hours, washed, and left to recover for 24 hours. The indicated EPO variants were present during all these steps. Cell viability was quantitated at the end of the experiment by incubation of cells with the tetrazolium dye WST-1 (according to manufacturer's instructions: Roche # 1644807) for 2 hours, and the viability was indicated as absorbance reading.

Figure 6A and 6B indicates the results of the SK-N-SH neuroblastoma cell neuroprotection assay (against rotenone) for erythropoietin as well as the recombinant tissue protective cytokines with the K45D and S100E recombinant tissue protective cytokines. The y-axis on the graph indicates the absorbance readings, and the data are means \pm range of duplicate determinations. The graph within **Figure 6A** clearly indicates that the viability of the cells within the K45D and S100E samples maintained their viability demonstrating their cellular protective effect. **Figure 6B** shows the plasmid map of hEPO-6xHisTag-PCiNeo.

6.4. EXAMPLE 4: TISSUE PROTECTIVE CYTOKINES

Recombinant tissue protective cytokines desirable for the uses described herein may be further modified by desialation, guanidination, carbamylation, amidination, trinitrophenylation, acetylation, succinylation, nitration, or modification of arginine residues or carboxyl groups, among other procedures. Alternatively, these modifications may be made to native erythropoietin or a derivative of erythropoietin, including, but not limited to, desialylated, guanidinated, carbamylated, amidinated, trinitrophenylated, acetylated, succinylated, or nitrated erythropoietin, prior to its mutation into a recombinant tissue protective cytokines. Some examples of further modifications to recombinant tissue protective cytokines are described below. One of ordinary skill in the art would understand that the procedures listed below may also be used to chemically modify native erythropoietin or its derivatives prior to the introduction of mutations to generate a recombinant tissue protective cytokine.

6.4.1. Desialylating Recombinant Tissue Protective Cytokines.

A recombinant tissue protective cytokines may be desialylated by the following exemplary procedure. Sialidase (isolated from *Streptococcus* sp 6646K.) is obtained from

SEIKAGAKU AMERICA (Code No. 120050). The recombinant tissue protective cytokine is subjected to desialylation by sialidase (0.025 U/mg EPO) at 37 C for 3 h. Desalt and concentrate the reaction mixture by Ultrafree Centrifugal Filter Unit. Apply sample to an ion exchange column in AKTAprime™ system. Elute protein with the selected buffers.

- 5 Perform IEF gel analysis of the eluted fractions containing a significant amount of protein. Pool the fractions containing only the top two bands (migrating at pI ~8.5 and ~7.9 on IEF gel). Determine the protein content and add 1/9 volumes of 10 x salt solution (1 M NaCl, 0.2 M sodium citrate, 3 mM citric acid). Determine the sialic acid content. No significant sialic content is detected.

- 10 Asialoerythropoietin was as effective as native erythropoietin for neural cells *in vitro* as shown in **Figures 7-8**. *In-vitro* testing was carried out using neural-like embryonal carcinoma cells (P19) that undergo apoptosis upon the withdrawal of serum. Twenty-four hours before the removal of serum, 1-1000 ng/ml of erythropoietin or a modified erythropoietin was added to the cultures. The following day the medium was removed, the
- 15 cells washed with fresh, non-serum containing medium, and medium containing the test substance (no serum) added back to the cultures for an additional 48 hours. To determine the number of viable cells, a tetrazolium reduction assay was performed (CellTiter 96; Promega, Inc.). As **Figures 7-8** illustrate, asialoerythropoietin appears to be of equal potency to erythropoietin itself in preventing cell death.

- 20 Retention of neuroprotective activity *in vivo* was confirmed using a rat focal ischemia model in which a reversible lesion in the territory of the middle cerebral artery is performed as described previously (Brines *et al.*, 2000, Proc. Nat. Acad. Sci. U.S.A. 97:10526-31). Adult male Sprague-Dawley rats were administered asialoerythropoietin or erythropoietin (5000 U/kgBW intraperitoneally) or vehicle at the onset of the arterial
- 25 occlusion. Twenty-four hours later, the animals were sacrificed and their brains removed for study. Serial sections were cut and stained with tetrazolium salts to identify living regions of the brain. As shown in **Figure 9**, asialoerythropoietin was as effective as native erythropoietin in providing neuroprotection, i.e. reducing infarct volume, from 1 hour of ischemia. **Figure 10** shows the results of another focal ischemia model in which a
- 30 comparative dose response was performed with erythropoietin and asialoerythropoietin. At the lowest dose of 4 µg /kg, asialoerythropoietin afforded protection whereas unmodified erythropoietin did not. The number of rats in each group, n, was greater than or equal to 4.

Similar results would be expected from asialo recombinant tissue protective cytokines of the present invention.

6.4.2. Carbamylating Recombinant Tissue Protective Cytokine.

The recombinant tissue protective cytokine may be used to prepare the respective
5 carbamylated molecules, in accordance with the following procedure, as described in Jin
Zeng (1991). Lysine modification of metallothionein by carbamylation and guanidination.
Methods in Enzymology 205: 433-437. Recrystallize potassium cyanate. Prepare 1 M
Borate buffer (pH 8.8). Mix a recombinant tissue protective cytokine solution with equal
volume of borate buffer. Add potassium cyanate directly to the reaction tube to a final
10 concentration of 0.5 M. Mix well and incubate at 37 C for 6-16 h. Dialyze thoroughly.
Remove the product from the dialysis tubing and collect into a fresh tube. Measure the
volume and add 1/9 volume of 10 X salt solution (1 M NaCl, 0.2 M sodium citrate, 3 mM
citric acid). Determine the protein content and calculate the product recovery rate. Analyze
the products by IEF gel followed by an *in vitro* test with TF-1 cells.

15 6.4.3. Succinylating Recombinant Tissue Protective Cytokines.

The recombinant tissue protective cytokine may be used to prepare the respective
succinylated molecule, in accordance with the following procedure, as described in Alcalde
et al. (2001). Succinylation of cyclodextrin glycosyltransferase from *Thermoanaerobacter*
sp. 501 enhances its transferase activity using starch as donor. J. Biotechnology 86: 71-80.
20 Recombinant tissue protective cytokine (100 ug) in 0.5 M NaHCO₃ (pH 8.0) was incubated
with a 15 molar excess of succinic anhydride at 15 C for 1 hour. The reaction was stopped
by dialysis against distilled water.

Dissolve succinic anhydride in dry acetone at 27 mg/ml. Do the reaction in an
eppendorf tube in 10 mM sodium phosphate buffer (pH 8.0). Add recombinant tissue
25 protective cytokine protein and 50-fold molar of succinic anhydride to the tube. Mix well
and rotate the tube at 4 C for 1 h. Stop the reaction by dialysis against 10 mM sodium
phosphate buffer, using the Dialysis cassette (Slide-A-Laze 7K, Pierce 66373). Remove the
product from the dialysis cassette and collect into a fresh tube. Measure the volume and add
1/9 volume of 10 X salt solution (1 M NaCl, 0.2 M sodium citrate, 3 mM citric acid).
30 Determine the protein content and calculate the product recovery rate. Analyze the products
by IEF gel followed by an *in vitro* test with TF-1 cells.

6.4.4. Acetylating Recombinant Tissue Protective Cytokine.

The recombinant tissue protective cytokine may be used to prepare the respective acetylated molecule, in accordance with the following procedure, as described in Satake *et al* (1990). Chemical modification of erythropoietin: an increase in *in-vitro* activity by
5 guanidination. *Biochimica et Biophysica Acta*. 1038: 125-129.

Perform the reaction in an eppendorf tube in 80 mM sodium phosphate buffer (pH 7.2). Add recombinant tissue protective cytokine and equal molar of acetic anhydride. Mix well and incubate on ice for 1 h. Stop the reaction by dialysis against water, using the Dialysis cassette (Slide-A-Laze 7K, Pierce 66373). Remove the product from the dialysis
10 cassette and collect into a fresh tube. Measure the volume and add 1/9 volume of 10 X salt solution (1 M NaCl, 0.2 M sodium citrate, 3 mM citric acid). Determine the protein content and calculate the product recovery rate. Analyze the products by IEF gel followed by an *in vitro* test with TF-1 cells.

6.4.5. Carboxymethylating Lysine of Recombinant Tissue Protective Cytokine.

15 The recombinant tissue protective cytokine may be used to prepare the respective N ϵ -(carboxymethyl)lysine (CML) modified molecules in which one or more lysyl residues of the recombinant tissue protective cytokine are modified, in accordance with the following procedure, as described in Akhtar *et al* (1999) Conformational study of N ϵ -(carboxymethyl)lysine adducts of recombinant α -crystallins. *Current Eye Research*, 18:
20 270-276.

Freshly prepare 200 mM of glyoxylic acid and 120 mM of NaBH₃CN in sodium phosphate buffer (50 mM, pH 7.5). In an eppendorf tube, add recombinant tissue protective cytokine (in phosphate buffer); calculate the lysine equivalent in the solution (about 8 lysine residues/mol). Add 3-times greater NaBH₃CN and 5 or 10-times greater glyoxylic acid to
25 the tube. Vortex each tube and incubate at 37 C for 5 h. Dialyze the samples against phosphate buffer overnight at 4 C. Measure the volume of each product after dialysis. Determine protein concentration and calculate the product recovery rate. Analyze the products by IEF gel followed by an *in vitro* test with TF-1 cells.

6.4.6. Iodinating Recombinant Tissue Protective Cytokine

30 A recombinant tissue protective cytokine may be used to prepare the respective iodinated molecule, in accordance with the following procedure, as described in instruction

provided by Pierce Chemical Company (Rockford, IL) for IODO-Gen Pre-Coated Iodination Tubes (product # 28601).

Prepare 0.1 M of NaI and perform iodination in IODO-Gen Pre-Coated Iodination Tube (Pierce, 28601), with total reaction volume of 0.1 ml/tube in sodium phosphate buffer (40 mM, pH 7.4). Mix the protein substrate (recombinant tissue protective cytokine) with sodium phosphate buffer and then transfer to an IODO-Gen Pre-Coated Iodination Tube. Add NaI to final concentration of 1 - 2 mM, making the molar ration of NaI/protein as 14-20. Mix well and incubate at room temperature for 15 min with gentle agitation. Stop the reaction by removing the reaction mixture and add to a tube containing 3.9 ml of sodium buffer (i.e., a 40-fold dilution). Concentrate the product by a pre-wet Ultrafree centrifugal filter unit. Measure the volume of concentrate and add 1/9 volume of 10 X salt solution (1 M NaCl, 0.2 M sodium citrate, 3 mM citric acid). Determine protein concentration and calculate the product recovery rate. Analyze the products by IEF gel followed by an *in vitro* test with TF-1 cells.

Alternatively the recombinant tissue protective cytokine may be iodinated using the following procedure. One Iodo Bead (Pierce, Rockford, IL) was incubated in 100 ul PBS (20mM sodium phosphate, 0.15M NaCl, pH7.5) containing 1 mCi free Na¹²⁵I for 5 minutes. One hundred micrograms of recombinant tissue protective cytokine in 100 ul PBS was then added to the mixture. After a ten minute incubation period at room temperature, the reaction was stopped by removing the 200 ul solution from the reaction vessel (leaving the iodo bead behind). The excess iodine was removed by gel filtration on a Centricon 10 column. As shown in **Figure 11**, iodo-erythropoietin produced in this manner is efficacious in protecting P19 cells from serum withdrawal. One of ordinary skill in the art would recognize that similar results would be expected from the iodination of a recombinant tissue protective cytokine of the present invention.

Yet another method for iodinating a recombinant tissue protective cytokine is outlined below. One Hundred micrograms of recombinant tissue protective cytokine in 100 ul PBS was added to 500 uCi Na¹²⁵I were mixed together in an eppendorf tube. Twenty-five microliters of chloramines T (2 mg/ml) was then added and the mixture was incubated for 1 minute at room temperature. Fifty microliters of Chloramine T stop buffer (2.4 mg/ml sodium metabisulfite, 10 mg/ml tyrosine, 10% glycerol, 0.1% xylene in PBS) was then added. The iodotyrosine and iodinated recombinant tissue protective cytokine were then separated by gel filtration on a Centricon 10 column.

6.4.7. Biotinylating Recombinant Tissue Protective Cytokine.

A recombinant tissue protective cytokine may be used to prepare the respective biotinylated molecules, in accordance with the following procedure, as described in instruction provided by Pierce Chemical Company (Rockford, IL) for EZ-Link NHS-LC-Biotin (product # 21336).

Immediately before the reaction, dissolve EZ-Link NHS-LC-Biotin (pierce, 21336) in DMSO at 2 mg/ml. Perform the reaction in a tube (17 x 100 mm) with total volume of 1 ml containing 50 mM sodium bicarbonate (pH 8.3). Add recombinant tissue protective cytokine and < 10% of EZ-Link NHS-LC-Biotin, with molar ratio of Biotin/protein at ~ 20. Mix well and incubate on ice for 2 h. Desalt and concentrate the reaction product by Ultrafree centrifugal filter unit. Collect the product into a fresh tube. Measure the volume and add 1/9 volume of 10 X salt solution (1 M NaCl, 0.2 M sodium citrate, 3 mM citric acid). Determine the protein content and calculate the product recovery rate. Analyze the products by IEF gel followed by an *in vitro* test with TF-1 cells.

A method for biotinylating the free amino groups of a recombinant tissue protective cytokine is disclosed below. 0.2 mg D-biotinoyl-e-aminocaproic acid-N-hydroxysuccinimide ester (Boehringer Mannheim #1418165) was dissolved in 100 ul DMSO. This solution was combined with 400 ul PBS containing approximately 0.2 mg recombinant tissue protective cytokine in a foil covered tube. After incubation for 4 hours at room temperature, the unreacted biotin was separated by gel filtration on a Centricon 10 column. As shown in **Figure 12**, this biotinylated erythropoietin protects p19 cells from serum withdrawal. One of ordinary skill in the art would recognize that similar results would be expected from the biotinylation of a recombinant tissue protective cytokine of the present invention.

Lastly, in "Biotinylated recombinant human erythropoietins: Bioactivity and Utility as a receptor ligand" by Wojchowski *et al.* Blood, 1989, 74(3):952-8, the authors use three different methods of biotinylating erythropoietin. Biotin is added to (1) the sialic acid moieties; (2) carboxylate groups; and (3) amino groups. The authors use a mouse spleen cell proliferation assay to demonstrate that (1) the addition of biotin to the sialic acid moieties does not inactivate the biological activity of erythropoietin; (2) the addition of biotin to carboxylate groups led to substantial biological inactivation of erythropoietin; (3) the addition of biotin to amino groups resulted in complete biological inactivation of

erythropoietin. These methods and modifications are fully embraced herein. **Figure 12** shows the activity of biotinylated erythropoietin and asialoerythropoietin in the serum-starved P19 assay. One of ordinary skill in the art would recognize that similar results would be expected from the iodination of a recombinant tissue protective cytokine of the present invention, see Section 6.15.

6.5. EXAMPLE 5: MODIFICATION OF RECOMBINANT TISSUE PROTECTIVE CYTOKINES BY OTHER METHODS

6.5.1. Trinitrophenylation

Recombinant tissue protective cytokine (100 ug) was modified with 2,4,6-trinitrobenzenesulfonate as described in Plapp *et al* ("Activity of bovine pancreatic deoxyribonuclease A with modified amino groups" 1971, J. Biol. Chem. 246, 939-845).

6.5.2. Arginine modifications

Recombinant tissue protective cytokine was modified with 2,3 butanedione as described in Riordan ("Functional arginyl residues in carboxypeptidase A. Modification with butanedione" Riordan JF, Biochemistry 1973, 12(20): 3915-3923).

In another modification wherein the amino acid residues of erythropoietin are modified, arginine residues were modified by using phenylglyoxal according to the protocol of Takahashi (1977, *J. Biochem.* 81:395-402) carried out for variable lengths of time ranging from 0.5 to 3 hrs at room temperature. The reaction was terminated by dialyzing the reaction mixture against water. Use of such modified forms of erythropoietin is fully embraced herein. The phenylglyoxal-modified erythropoietin was tested using the neural-like P19 cell assay described above. As **Figure 13** illustrates, this chemically-modified erythropoietin fully retains its neuroprotective effects. Similar results form a similarly modified recombinant tissue protective cytokine of the present invention.

A recombinant tissue protective cytokine was modified with cyclohexanone as in Patthy *et al* ("Identification of functional arginine residues in ribonuclease A and lysozyme" Patthy, L, Smith EL, J. Biol. Chem 1975 250(2): 565-9).

A recombinant tissue protective cytokine was modified with phenylglyoxal as described in Werber *et al.* ("Proceedings: Carboxypeptidase B: modification of functional arginyl residues" Werber, MM, Sokolovsky M Isr J Med Sci 1975 11(11): 1169-70).

6.5.3. Tyrosine modifications

Recombinant tissue protective cytokine (100 ug) was incubated with tetranitromethane as previously described in Nestler *et al* "Stimulation of rat ovarian cell steroidogenesis by high density lipoproteins modified with tetranitromethane" Nestler JE,
5 Chacko GK, Strauss JF 3rd. J Biol Chem 1985 Jun 25;260(12):7316-21).

6.5.4. Glutamic acid (and aspartic acid) modifications

In order to modify carboxyl groups, recombinant tissue protective cytokine (100 ug) was incubated with 0.02 M EDC in 1M glycylamide at pH 4.5 at room temperature for 60 minutes as described in Carraway *et al* "Carboxyl group modification in chymotrypsin and
10 chymotrypsinogen." Carraway KL, Spoerl P, Koshland DE Jr. J Mol Biol 1969 May 28;42(1):133-7.

6.5.5. Tryptophan residue modifications

A recombinant tissue protective cytokine (100 ug) was incubated with 20 uM n-bromosuccinimide in 20 mM potassium phosphate buffer (pH 6.5) at room temperature as
15 described in Ali *et al.*, J Biol Chem. 1995 Mar 3;270(9):4570-4. The number of oxidized tryptophan residues was determined by the method described in Korotchkina (Korotchkina, LG *et al* Protein Expr Purif. 1995 Feb;6(1):79-90).

6.5.6. Removal of amino groups

In order to remove amino groups of recombinant tissue protective cytokines 100 ug
20 was incubated with in PBS (pH 7.4) containing 20mM ninhydrin (Pierce Chemical, Rockford, Il), at 37 C for two hours as in Kokkini *et al* (Kokkini, G., *et al* "Modification of hemoglobin by ninhydrin" Blood, Vol. 556, No 4 1980: 701-705). Reduction of the resulting aldehyde was accomplished by reacting the product with Sodium borohydride or lithium aluminum hydride. Specifically, erythropoietin (100 ug) was incubated with 0.1M
25 sodium borohydride in PBS for 30 minutes at room temperature. The reduction was terminated by cooling the samples on ice for 10 minutes and dialyzing it against PBS, three times, overnight. (Kokkini, G., Blood, Vol. 556, No 4 1980: 701-705). Reduction using lithium aluminum hydride was accomplished by incubating recombinant tissue protective cytokine (100 ug) with 0.1M lithium aluminum hydride in PBS for 30 minutes at room
30 temperature. The reduction was terminated by cooling the samples on ice for 10 minutes and dialyzing it against PBS, three times, overnight.

6.5.7. Disulfide reduction and stabilization

A recombinant tissue protective cytokine (100 ug) was incubated with 500 mM DTT for 15 minutes at 60 C. 20 mM iodoacetamide in water was then added to the mixture and incubated for 25 minutes, at room temperature in the dark.

5 6.5.8. Limited proteolysis

A recombinant tissue protective cytokine can be subjected to a limited chemical proteolysis that targets specific residues. A recombinant tissue protective cytokine can be reacted with 2-(2-nitrophenylsulfenyl)-3-methyl-3'-bromoindolenine which cleaves specifically after tryptophan residues in a 50 times excess in 50% acetic acid for 48 hours in
10 the dark at room temperature in tubes capped under nitrogen pressure. The reaction was terminated by quenching with tryptophan and desalting.

As noted above, a recombinant tissue protective cytokine may be modified, yet multiple modifications as well as additional modifications of the tissue protective cytokine molecule may also be performed without deviating from the spirit of the present invention.

15

6.6. EXAMPLE 6: TISSUE PROTECTIVE CYTOKINES HAVE NEURO PROTECTIVE EFFECT

The neuroprotective affects of chemically modified erythropoietin was evaluated
20 using a water intoxication assay in accordance with Manley *et al.*, 2000, Aquaporin-4 deletion in mice reduces brain edema after acute water intoxication and ischemic stroke, Nat Med 2000 Feb;6(2):159-63. Female C3H/HEN mice were used. The mice were given 20% of their body weight as water IP with 400 ng/kg bw DDAVP (desmopressin). The mice were administered erythropoietin (A) or a tissue protective cytokine: asialoerythropoietin
25 (B), carbamylated asialoerythropoietin (C); succinylated asialoerythropoietin (D), acetylated asialoerythropoietin (E); iodinated asialoerythropoietin (F); carboxymethylated asialoerythropoietin (G); carbamylated erythropoietin (H); acetylated erythropoietin (I); iodinated erythropoietin (J) or N^ε-carboxy methyl erythropoietin (K). The mice were given a 100 microgram/kg dose of erythropoietin or chemically modified erythropoietin
30 intraperitoneally 24 hrs before administration of the water and again at the time of the water

administration. A modified scale from Manley *et al.* was used to evaluate the mice. The modified scale is as listed below:

Explores cage/table		
5	Yes	0
	No	1
Visually tracks objects		
	Yes	0
	No	1
Whisker movement		
10	Present	0
	Absent	1
Leg-tail movements		
15	Normal	0
	Stiff	1
	Paralyzed	2
Pain withdrawal (toe pinch)		
20	Yes	0
	No	1
Coordination of movement		
	Normal	0
	Abnormal	1
Stops at edge of table		
25	Yes	0
	No	1
Total score possible : 8		

The mice were scored at the following time points: 15, 30, 45, 60, 75, 90, 120, 150, and 180 minutes. **Figure 14** plots the performance of the mice that received erythropoietin or the chemically modified erythropoietin as a percentage of the neuronal deficit experienced by the control mice. **Figure 14** shows that the tissue protective cytokines protect the mice from the neurological trauma induced by the water intoxication. Similar results would be expected from recombinant tissue protective cytokines with similar chemical modifications. Statistical significance was also determined. Those administration regimens with significant differences, $p < 0.05$, in comparison to controls are indicated with *, while those with highly significant differences, $p < 0.01$, are indicated by **.

6.7. EXAMPLE 7: MAINTENANCE OF FUNCTION IN HEART PREPARED FOR TRANSPLANTATION

Wistar male rats weighing 300 to 330g are given erythropoietin (5000 U/kg body weight) or vehicle 24h prior to removal of the heart for *ex vivo* studies, done in accordance with the protocol of Delcayre *et al.*, 1992, *Amer. J. Physiol.* 263:H1537-45. Animals are sacrificed with pentobarbital (0.3mL), and intravenously heparinized (0.2mL). The hearts are initially allowed to equilibrate for 15 min. The left ventricular balloon is then inflated to a volume that gives an end-diastolic pressure of 8 mm Hg. A left ventricular pressure-volume curve is constructed by incremental inflation of the balloon volume by 0.02 ml aliquots. Zero volume is defined as the point at which the left ventricular end-diastolic pressure is zero. On completion of the pressure-volume curve, the left ventricular balloon is deflated to set end-diastolic pressure back to 8mmHg and the control period is pursued for 15 min, after check of coronary flow. Then the heart is arrested with 50 mL Celsior + molecule to rest at 4°C under a pressure of 60cm H₂O. The heart is then removed and stored for 5 hours at 4°C in plastic container filled with the same solution and surrounded with crushed ice.

On completion of storage, the heart is transferred to a Langendorf apparatus. The balloon catheter is reinserted into the left ventricle and re-inflated to the same volume as during preischemic period. The heart is re-perfused for at least 2 hours at 37°C. The re-perfusion pressure is set at 50 cm H₂O for 15min of re-flow and then back to 100 cm H₂O for the 2 next hours. Pacing (320 beats per minute) is re-instituted. Isovolumetric measurements of contractile indexes and diastolic pressure are taken in triplicate at 25, 45, 60, and 120 min of reperfusion. At this time point pressure volume curves are performed and coronary effluent during the 45mn reperfusion collected to measure creatine kinase

leakage. The two treatment groups are compared using an unpaired t-test, and a linear regression using the end-diastolic pressure data is used to design compliance curves. As shown in **Figure 15**, significant improvement of left ventricular pressure developed occurs after treatment with erythropoietin, as well as improved volume-pressure curve, decrease of left diastolic ventricular pressure and decrease of creatine kinase leakage. Similar results would be expected from treatment with recombinant tissue protective cytokines of the present invention.

6.8. EXAMPLE 8: ERYTHROPOIETIN PROTECTS MYOCARDIUM FROM ISCHEMIC INJURY

Adult male rats given recombinant human erythropoietin (5000 U/kg body weight) 24 hrs previously are anesthetized and prepared for coronary artery occlusion. An additional dose of erythropoietin is given at the start of the procedure and the left main coronary artery occluded for 30 minutes and then released. The same dose of erythropoietin is given daily for one week after treatment. The animals are then studied for cardiac function. As **Figure 16** illustrates, animals receiving a sham injection (saline) demonstrated a large increase in the left end diastolic pressure, indicative of a dilated, stiff heart secondary to myocardial infarction. In contradistinction, animals receiving erythropoietin suffered no decrement in cardiac function, compared to sham operated controls (difference significant at the $p < 0.01$ level). Similar results would be expected from treatment with recombinant tissue protective cytokines of the present invention.

6.9. EXAMPLE 9: PROTECTION OF RETINAL ISCHEMIA BY PERIPHERALLY-ADMINISTERED ERYTHROPOIETIN.

Retinal cells are very sensitive to ischemia such that many will die after 30 minutes of ischemic stress. Further, subacute or chronic ischemia underlies the deterioration of vision which accompanies a number of common human diseases, such as diabetes mellitus, glaucoma, and macular degeneration. At the present time there are no effective therapies to protect cells from ischemia. A tight endothelial barrier exists between the blood and the retina that excludes most large molecules. To test whether peripherally-administered erythropoietin will protect cells sensitive to ischemia, an acute, reversible glaucoma rat model was utilized as described by Rosenbaum *et al.* (1997; Vis. Res. 37:3443-51). Specifically, saline was injected into the anterior chamber of the eye of adult male rats to a pressure above systemic arterial pressure and maintained for 60 minutes. Animals were

administered saline or 5000 U erythropoietin/kg body weight intraperitoneally 24 hours before the induction of ischemia, and continued as a daily dose for 3 additional days. Electoretinography was performed on dark-adapted rats 1 week after treatment. **Figure 17-18** illustrate that the administration of erythropoietin is associated with good preservation of the electroretinogram (ERG) (Figure 17, Panel D), in contrast to animals treated with saline alone (Figure 17, Panel C), for which very little function remained. Figure 18 compares the electroretinogram a- and b-wave amplitudes after 60 minutes ischemia for the erythropoietin-treated and saline-treated groups, and shows significant protection afforded by erythropoietin. Similar results are obtainable from treatment with recombinant tissue protective cytokines of the present invention.

6.10. EXAMPLE 10: RESTORATIVE EFFECTS OF ERYTHROPOIETIN ON DIMINSHED COGNITIVE FUNCTION ARISING FROM BRAIN INJURY

In a study to demonstrate the ability of erythropoietin to restore diminished cognitive function in mice after receiving brain trauma, female Balb/c mice were subject to blunt brain trauma as described in Brines *et al.* PNAS 2000, 97; 10295-10672 and five days later, daily erythropoietin administration of 5000 U/kg-bw intraperitoneally began. Twelve days after injury, animals were tested for cognitive function in the Morris water maze, with four trials per day. While both treated and untreated animals performed poorly in the test (with swim times of about 80 seconds out of a possible 90 seconds), **Figure 19** shows the results of the Morris Water maze test, with each group of mice, n=16, after blunt brain trauma with EPO administration beginning on day 5 after injury. The first test began 1 week after EPO dosing began (12 days after injury). Both groups of animals did poorly with swim times ~ 80 out of 90 seconds possible. The erythropoietin-treated animals performed better (in this presentation, a negative value is better). Means of 4 trials per day were used. Figure 19 shows that. Even if the initiation of erythropoietin treatment is delayed until 30 days after trauma (**Figure 20**), restoration of cognitive function is also seen. In Figure 20, the each group of mice, n=7, were treated with 5000 U/kg EPO daily except on weekends, beginning one month after injury. Means of trials were also 4 trials each day. Similar results are obtainable from treatment with recombinant tissue protective cytokines of the present invention.

6.11. EXAMPLE 11: KAINATE MODEL

In the kainate neurotoxicity model, asialoerythropoietin was administered according to the protocol of Brines *et al.* Proc. Nat. Acad. Sci. U.S.A. 2000, 97; 10295-10672 at a dose of 5000U/kg-bw given intraperitoneally 24 hours before the administration of 25 mg/kg kainate is shown to be as effective as erythropoietin, as shown by time to death (Figure 21). Similar results are obtainable from treatment with tissue protective cytokines of the present invention.

6.12. EXAMPLE 12: SPINAL CORD INJURY MODELS

6.12.1. Rat Spinal Cord Compression Testing Erythropoietin and Tissue Protective Cytokines

Wistar rats (female) weighing 180–300g were used in this study. The animals were fasted for 12 h before surgery, and were humanely restrained and anesthetized with an intraperitoneal injection of thiopental sodium (40 mg/kg-bw). After infiltration of the skin (bupivacaine 0.25%), a complete single level (T-3) laminectomy was performed through a 2 cm incision with the aid of a dissecting microscope. Traumatic spinal cord injury was induced by the extradural application of a temporary aneurysm clip exerting a 0.6 newton (65 grams) closing force on the spinal cord for 1 minute. After removal of the clip, the skin incision was closed and the animals allowed to recover fully from anesthesia and returned to their cages. The rats were monitored continuously with bladder palpation at least twice daily until spontaneous voiding resumed.

Forty animals were randomly divided into five groups. Animals in the control group (I) (n= 8) received normal saline (via intravenous injection) immediately after the incision is closed. Group (II; n= 8) received rhEPO @ 16 micrograms/kg-bw iv; group (III) received an asialo tissue protective cytokine of the present invention (asialoerythropoietin) @ 16 micrograms/kg-bw iv, group (IV) received an asialo tissue protective cytokine @ 30 micrograms/kg-bw iv, and group (V) received an asialo tissue protective cytokine of the present invention (asialoerythropoietin) @ 30 micrograms/kg-bw; all as a single bolus intravenous injection immediately after removal of the aneurysm clip.

Motor neurological function of the rats will be evaluated by use of the locomotor rating scale of Basso *et al.* In this scale, animals are assigned a score ranging from 0 (no observable hindlimb movements) to 21 (normal gait). The rats will be tested for functional deficits at 1, 12, 24, 48, and 72 hours and then at 1 week after injury by the same examiner who is blind to the treatment each animal receives.

Figure 22 is a graph demonstrating the locomotor ratings of the rats recovering from the spinal cord trauma over a period of thirty days. As can be seen from the graph, the rats that were given erythropoietin (group II) or tissue protective cytokines (groups III-V) recovered from the injury more readily and demonstrated better overall recovery from the injury than the control rats. Similar results would be expected from the therapeutic treatment with the recombinant tissue protective cytokines of the present invention.

In a second related study animals were injured in the same way. Forty animals were randomly divided into three groups. Animals in the control group (n= 8) received normal saline (via intravenous injection) immediately after the incision is closed. The second group (n= 8) received methylprednisolone @ 30 mg/kg per day x3 then biweekly, a common therapeutic for spinal cord injury; the third group received an recombinant tissue protective cytokine, S100E, of the present invention at a dose of 10ug/kg immediately following injury, all as a single bolus intravenous injection immediately after removal of the aneurysm clip.

Motor neurological function of the rats will be evaluated by use of the locomotor rating scale of Basso *et al.* In this scale, animals are assigned a score ranging from 0 (no observable hind limb movements) to 21 (normal gait). The rats will be tested for functional deficits at 1, 12, 24 , 48, and 72 hours and then at 1 week after injury by the same examiner who is blind to the treatment each animal receives.

Figure 37 is a graph demonstrating the locomotor ratings of the rats recovering from the spinal cord trauma over a period of forty-two days. As can be seen from the graph, the rats that were given S100E recovered from the injury more readily and demonstrated better overall recovery from the injury than the control rats and rats administered methylprednisolone.

6.12.2. Rabbit Spinal Cord Ischemia Testing Erythropoietin and a Tissue Protective Cytokine.

Thirty-six New Zealand White rabbits (8-12 months old, male) weighing 1.5-2.5 kg were used in this study. The animals were fasted for 12 hours and humanely restrained. Anesthesia induction was via 3% halothane in 100% oxygen and maintained with 0.5 – 1.5 % halothane in a mixture of 50% oxygen and 50% air. An intravenous catheter (22 gauge) was placed in the left ear vein. Ringers lactate was infused at a rate of 4 ml/kg body weight (bw) per hour during the surgical procedure. Preoperatively, cefazoline 10 mg/kg-bw was

administered intravenously for prophylaxis of infection. The animals were placed in the right lateral decubitus position, the skin prepared with povidone iodine, infiltrated with bupivacaine (0.25%) and a flank skin incision was made parallel to the spine at the 12th costal level. After incision of the skin and subcutaneous thoracolumbar fascia, the
5 longissimus lumborum and iliocostalis lumborum muscles were retracted. The abdominal aorta was exposed via a left retroperitoneal approach and mobilized just inferior to the left renal artery. A piece of PE-60 tubing was looped around the aorta immediately distal to the left renal artery and both ends passed through a larger rubber tube. By pulling on the PE tubing, the aorta was non-traumatically occluded for 20 minutes. Heparin (400 IU) was
10 administered as an intravenous bolus before aortic occlusion. After 20 minutes of occlusion, the tube and catheter were removed, the incision was closed and the animals were monitored until full recovery and then were serially assessed for neurological function.

Thirty-six animals were randomly divided into six groups. In a control group (I), animals (n = 6) received normal saline intravenously immediately after release of aortic
15 occlusion. Group (II) received rhEPO @ 6.5 microgram/kg-bw; group (III) received a tissue protective cytokine (carbamylated erythropoietin) @ 6.5 microgram/kg-bw; group (IV) received another tissue protective cytokine (asialoerythropoietin) @ 6.5 microgram/kg-bw; group (V) received the same tissue protective cytokine as group (IV) but @ 20 microgram/kg-bw; and group (VI) received yet another tissue protective cytokine
20 (asialocarbamylatederythropoietin) @ 20 microgram/kg-bw all intravenously immediately after reperfusion (n = 6 for each group).

Motor function was assessed according to the criteria of Drummond and Moore by an investigator blind to the treatment at 1, 24 and 48 h after reperfusion. A score of 0 to 4 was assigned to each animal as follows: 0 = paraplegic with no evident lower extremity
25 motor function; 1 = poor lower extremity motor function, weak antigravity movement only; 2 = moderate lower extremity function with good antigravity strength, but inability to draw legs under body; 3 = excellent motor function with the ability to draw legs under body and hop, but not normally; 4 = normal motor function. The urinary bladder was evacuated manually in paraplegic animals twice a day.

30 **Figure 23** is a graph plotting motor function of the recovering rabbits. The graph demonstrates that even over a period of only two days erythropoietin and the tissue protective cytokines of the present invention permit the rabbits to recover more fully from

the spinal cord injury. Similar results would be expected from the therapeutic treatment with the recombinant tissue protective cytokines of the present invention.

6.13. EXAMPLE 13: ANTI-INFLAMMATORY AFFECTS OF ERYTHROPOIETIN

5 In-Vivo Studies:

MCAO in Rats

Male Crl:CD(SD)BR rats weighing 250-280 g were obtained from Charles River, Calco, Italy. Surgery was performed on these rats in accordance with the teachings of Brines, M.L., Ghezzi, P., Keenan, S., Agnello, D., de Lanerolle, N.C., Cerami, C., Itri, L.M., and Cerami, A. 2000 Erythropoietin crosses the blood-brain barrier to protect against experimental brain injury, Proc Natl Acad Sci USA 97:10526-10531. Briefly, the rats were anesthetized with chloral hydrate (400 mg/kg-bw, i.p.), the carotid arteries were visualized, and the right carotid was occluded by two sutures and cut. A burr hole adjacent and rostral to the right orbit allowed visualization of the MCA, which was cauterized distal to the rhinal artery. To produce a penumbra (borderzone) surrounding this fixed MCA lesion, the contralateral carotid artery was occluded for 1 hour by using traction provided by a fine forceps and then re-opened. PBS or rhEPO (5,000 U/kg-bw, i.p.; previously shown to be protective in this model (1)) were administered immediately after the MCAO. When indicated, TNF and IL-6 were quantified in brain cortex homogenates as previously described (8). MCP-1 was measured in the homogenates using a commercially available ELISA kit (biosource, Camarillo, CA).

Twenty-four hours after MCAO, the rats were anesthetized as described above and transcardially perfused with 100 ml saline followed by 250 ml of sodium phosphate buffered 4% paraformaldehyde solution. Brains were rapidly removed, fixed in sodium phosphate buffered 4% paraformaldehyde solution for two hours, transferred to 20% sucrose solution in PBS overnight, then in 30% sucrose solution until they sank and were then frozen in 2-methylbutane at -45° C. Sections (30 μ m) were cut on a cryostat (HM 500 OM, Microm) at -20° C in the transverse plane through the brain and selected every fifth section for histochemistry against the different antigens, or hematoxylin-eosin staining. Free floating sections were processed for immunoreactivity both with anti-glial fibrillary acid protein (GFAP) mouse monoclonal antibody (1:250, Boehringer Mannheim, Monza, Italy) and with anti-cd11b (MRC OX-42) mouse monoclonal antibody (1:50, Serotec, UK),

according to the protocols described by Houser *et al.* and the manufacturer's protocol respectively. All sections were mounted for light microscopy in saline on coated slides, dehydrated through graded alcohols, fixed in xylene and coverslipped using DPX mountant (BDH, Poole, UK). Adjacent sections were stained with hematoxylin-eosin as described

5 (10).

Figure 24 shows a coronal section of the brain cortical layer stained by hematoxylin and eosin. Control rat (A), ischemic rat treated with PBS (B), ischemic rat treated with rhEPO (5,000 U/kg-bw, i.p., immediately following MCAO) (C). Section B shows a marked decrease in tissue staining consistent with inflammation, accompanied by a loss of neuronal component compared to the control (A). Systemic rhEPO administration largely reduces the ischemic damage localizing the cell death or injury in a restricted area (C). (Magnification 2.5x. Size bar = 800 μm .)

Figure 25 shows coronal sections of frontal cortex adjacent to the region of infarction stained by GFAP antibody. Control rat (A), ischemic rat treated with PBS (B), and ischemic rat treated with rhEPO (C). Activated astrocytes are visualized by their GFAP-positive processes (Panel B). Note the marked reduction in number as well as staining intensity in a representative rhEPO-treated animal (Panel C). (Magnification 10x. Size bar = 200 μm .)

Figure 26 shows coronal sections of brain cortical layer stained by OX-42 antibody. Ischemic rat treated with PBS (A), and ischemic rat treated with rhEPO (B). In the ischemic cerebral hemisphere, the cellular staining is especially prominent around the infarcted tissue in both treatment groups, but it is much denser and extends further in the saline treated group. (Magnification 20x; Size bar = 100 μm).

Figure 27 shows coronal sections of brain cortical layer adjacent to the region of infarction stained by OX-42 antibody. A much higher density of mononuclear inflammatory cells are observed in the tissue from an ischemic rat treated with PBS (A) compared to an ischemic rat treated with rhEPO (B). The infiltrating leukocytes, with typical round shape, potentially will extend the volume of infarction. (Magnification 10x; Size bar = 200 μm)

Similar results would be expected from the therapeutic treatment with the recombinant tissue protective cytokines of the present invention.

Acute Experimental Allergic Encephalomyelitis (EAE) in Lewis rats

Female Lewis rats, 6-8 weeks of age, were purchased from Charles River (Calco, Italy). EAE was induced in rats by injecting 50 µg of guinea pig MBP (Sigma, St. Louis, MO) in water emulsified in equal volumes of complete Freund's adjuvant (CFA, Sigma) 5 added with 7 mg/ml of heat-killed *M. tuberculosis* H37Ra (Difco, Detroit, MI) in a final volume of 100 µ under light ether anesthesia into both hind footpads. 1. Rats were examined in a blinded fashion for signs of EAE and scored as follows: 0, no disease; 1, flaccid tail; 2, ataxia; 3, complete hind limb paralysis with urinary incontinence. Starting from day 3 after immunization, rats were given r-Hu-EPO (EPOetin alfa, Procrit, Ortho 10 Biotech, Raritan, NJ) intraperitoneally (i.p.) once a day at the indicated doses, in PBS. Since the clinical-grade EPO contained human serum albumin, control animals were always given PBS containing an identical amount of human serum albumin. Daily administration of 5,000 U/kg-bw of EPO increased the hematocrit by 30%. When indicated, rats were injected s.c. 15 once a day from day 3 until day 18 with 1.3 mg/kg-bw dexamethasone (DEX) phosphate disodium salt (Sigma) corresponding to 1 mg/kg-bw of DEX, dissolved in PBS. When indicated, TNF and IL-6 were quantified in brain and spinal cord homogenates as previously described [Agnello, 2000 #10].

Figure 28 shows the protective effect on the clinical signs of EAE of different doses of EPO, given from day 3 after immunization with MBP until day 18. EPO, in a dose- 20 dependent fashion, delayed the onset of disease and decreased disease severity, as summarized in Table 1, but did not delay the time to greatest severity. As shown in this table, EPO at the doses of 2,500 and 5,000 U/kg-bw significantly decreased the mean cumulative score.

In experiments where treatment of EPO was discontinued after the disease regressed 25 and the rats were monitored up to two months, no relapse was observed, in contrast with DEX which induces an exacerbation of disease after suspending its administration (**Figure 29**). Similar results would be expected from the therapeutic treatment with the recombinant tissue protective cytokines of the present invention.

In Vitro Studies:

30 Primary cultures of glial cells were prepared from newborn Sprague-Dawley rats 1-2 days old. Cerebral hemispheres were freed from the meninges and mechanically disrupted. Cells were dispersed in a solution of trypsin 2.5% and DNAase 1%, filtered through a 100

μ m nylon mesh and plated (140,000 cells per 35 mm dish) in Eagle's minimum essential medium supplemented with 10% fetal calf serum, 0.6% glucose, streptomycin (0.1 mg/ml) and penicillin (100 U/ml). Glial cultures were fed twice a week and grown at 37° C in a humidified incubator with 5% CO₂. All experiments were performed on 2-3 week-old glial cell cultures with 97% astrocytes and 3% microglia, as assessed by immunocytochemistry of GFAP and Griffonia simplicifolia isolectin B4. Neuronal cultures were established from the hippocampus of 18-day rat fetuses. Brains were removed and freed from meninges and the hippocampus was isolated. Cells were dispersed by incubation for 15-20 min at 37° C in a 2.5% trypsin solution followed by titration. The cell suspension was diluted in the medium used for glial cells and plated onto polyornithine-coated coverslips at a density of 160,000 cells per coverslip. The day after plating, coverslips were transferred to dishes containing a glial monolayer in neuron maintenance medium (Dulbecco's modified Eagle's medium and Ham's nutrient mix F12 supplemented with 5 μ g/ml insulin, 100 μ g/ml transferrin, 100 μ g/ml putrescine, 30 nM Na selenite, 20 nM progesterone and penicillin 100 U/ml) supplemented with cytosine arabinoside 5 μ M. Coverslips were inverted so that the hippocampal neurons faced the glia monolayer. Paraffin dots adhering to the coverslips supported them above the glia, creating a narrow gap that prevented the two cell types from contacting each other but allowed the diffusion of soluble substances. These culture conditions allowed the growth of differentiated neuronal cultures with >98% homogeneity, as assessed by immunocytochemistry of microtubule-associated protein 2 and GFAP. Cells were then treated for 24 hours with 1 μ M Trimethyl tin (TMT), in the presence or absence of rhEPO (10 U/ml), the supernatants used for TNF assay and cellular viability evaluated as described below. When indicated, glial cells were cultured in the presence of LPS for 24 hours, with or without rhEPO, and TNF measured in the cultured supernatants. Cell viability was measured by the 3-(4,5-dimethyl-thiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. Denizot, F., and Lang, R. 1986. Rapid colorimetric assay for cell growth and survival. Modifications to the tetrazolium dye procedure giving improved sensitivity and reliability. J Immunol Methods 89:271-277. Briefly, MTT tetrazolium salt was dissolved in serum-free medium to a final concentration of 0.75 mg/ml and added to the cells at the end of the treatment for 3 h at 37° C. The medium was then removed and the formazan was extracted with IN HCl:isopropanol (1:24). Absorbance at 560 nm was read on a microplate reader.

Figure 30 shows that rhEPO prevents neuronal death-induced TNF production in mixed neuron-glia cultures. Panel A: Percentage of neural cell death induced by TMT 1

μ M without or with treatment with rhEPO (10 U/ml). Panel B: Release of TNF- α from glial cells exposed to TMT 1 μ M in the presence (hatched bars) or absence (filled bars) of neurons, with or without rhEPO (10 U/ml). Similar results would be expected from the therapeutic treatment with the recombinant tissue protective cytokines of the present invention.

6.14. EXAMPLE 14: NMDA INDUCED CELL DEATH ASSAY

Excitotoxicity can be defined as the excessive activation of glutamate receptors, such as the N-methyl-D-aspartate (NMDA) receptor. The NMDA receptor exhibits increased activity in response to ischemia and other traumas (Fauci et al., 1998, Harrison's Principles of Internal Medicine), (Nishizawa, 2001, Life Sci. 69, 369-381), (White et al., 2000, J. Neurol. Sci. 179, 1-33). Thus, the assay serves as a model for assessing a compound's effect on cell injury and death.

Protocol of NMDA excitotoxicity in primary hippocampal neurons

Primary hippocampal neuronal cultures were prepared from new born mice (less than 24 hours old) essentially as previously described by Krohn et al. (1998). Briefly, the hippocampi were dissected out in DMEM containing 0.02% BSA. The tissue was transferred to DMEM containing 0.1 % papain and incubated for 20 minutes at 37°C. The digestion was stopped by aspiration of the papain containing medium and addition of MEMII and the hippocampal cells were dissociated by titration with a 1000 μ l pipet tip. The tissue pieces were allowed to settle and the supernatant, containing single cells, was transferred into MEMII containing 1% trypsin inhibitor (type II-O) and 1% BSA. The titration-step was repeated three times before the single cells were centrifuged at 600U/minute for 10 minutes and resuspended in growth medium (MEMII, 20mM D-glucose, 100U/ml penicillin, 100 μ g streptomycin, 2 mM L-glutamine, 10% Nu-serum (bovine), 2% B27 supplement, 26.2 mM NaHCO₃). Cells from 10 hippocampi were used to seed one 24 well plate. One day after seeding, the cells were treated with cytosine-arabino-furanoside (1 μ M). On day two, the medium was changed and cytosine-arabino-furanoside (1 μ M) was added.

Excitotoxicity assay

Twelve day old cultures were pre-incubated with test compound (vehicle, R103E, R150E, or EPO) at 5nM for 24 hours. On day 13, the medium was removed from the cells

and kept while the cultures were challenged with 300 μ M NMDA for 5 minutes at room temperature. After the excitotoxic insult, the pre-conditioned medium was returned to the cultures and the injury was quantified by trypan blue exclusion after another 24 hours of incubation. Approximately 300 neurons were counted per condition in at least four separate wells and the experiments were repeated at least twice (Krohn, A.J., Preis, E. and Prehn, J.H.M. (1998) *J. Neurosci.* 18(20):8186-8197).

Figure 31 shows that human erythropoietin and recombinant tissue protective cytokines R130E and R150E effectively reduce cell death induced by NMDA when added to the primary hippocampal neuron cell cultures prior to NMDA treatment. Cells treated with R103E (5 nM) exhibited significantly less cell death in comparison to vehicle control cells ($p=0.01$). Cells treated with R103E (5 nM) exhibited significantly less cell death in comparison to vehicle control cells ($p=0.01$). Cells treated with R150E (5 nM) exhibited approximately a 20% decrease in cell death in comparison to solvent control cells ($p=0.001$). Statistics: ANOVA plus Tukey's post-hoc test.

6.15. EXAMPLE 15: NEURONAL PROTECTION OF SERUM WITHDRAWAL IN P19 CELLS

To examine the neuronal protective effect of the recombinant tissue protective cytokines of the invention, withdrawal of serum from PC19 cell cultures was used as a model. The clone P19S1801A1 was kindly provided by Dr. W.H. Fischer. The cells were maintained in Dulbecco's Modified Eagles Medium (DMEM) supplemented with 2 mM L glutamine, 100 U/ml penicillin G, 100 μ g/ml streptomycin sulfate and 10% fetal calf serum (FCS; all from Gibco, Paisley, Scotland, UK), containing 1.2 g/l NaHCO_3 , 10 mM Hepes buffer (Carlo Erba, Milano, Italy), hereafter referred to as complete medium, in a humidified incubator under an atmosphere of 7% CO_2 in air. Serum free medium (N2) has the same constituents as above with the deletion of serum, and the addition of the following: 5 μ g/ml of insulin, 100 μ g/ml of transferrin, 20 nM progesterone, 100 μ M putrescine and 30 nM Na_2SeO_3 (all from Sigma). For the death experiments, cells were dissociated with 10% pancreatin (Gibco), washed once with complete medium, twice with N2 medium and plated, unless otherwise indicated, plated in 25 cm^2 tissue culture flasks (Falcon Becton Dickinson, Lincoln Park, New Jersey) at a final density of 104 cells/ cm^2 in 5 ml of serum-free medium. L acetylcarnitine (100 μ M) is taken as a positive control, that confers protection, reducing by 50% the percentage of apoptotic nuclei 24 h after serum deprivation. Twenty-four h after serum deprivation, cells were detached by tapping on the flask (without trypsin)

seeded on microscope slides by cytospin centrifugation (Shandon Southern, USA) at 600 rpm for 10 min, and fixed in Carnoy solution (methanol:acetic acid, 3:1) for 10 min, stained with Hoechst 33258 (0.1 µg/ml PBS) for 1 h at 37° C, washed with tap water for 15 min, air dried and mounted. Slides were observed with a fluorescence microscope (Zeiss, Germany) at an excitation wavelength of 365 nm. The percentage of apoptotic nuclei was determined by counting in blind a total of 100 cells in at least 5 determinations.

P19 cells were pre-incubated with 3 nM Epo or recombinant tissue protective cytokine S100E for 24h. This treatment resulted in significant ($p < 0.001$) protection from apoptosis triggered by serum withdrawal. Data are means from triplicate determinations within one experiment. The experiment was performed twice with similar results.

Figure 32 shows neuronal protection from serum withdrawal in P19 cells. The percent of apoptotic cells decreased for cells pretreated with Epo, EpoWT, and recombinant tissue protective cytokine S100E. Cells treated with Epo exhibited approximately a 20% decrease in apoptotic cell death in comparison to untreated control cells. Cells treated with EpoWT and S100E both exhibited approximately a 10% decrease in apoptotic cell death in comparison to untreated control cells.

6.16. EXAMPLE 16: NGF WITHDRAWAL IN DIFFERENTIATED PC12 CELLS

To examine the neuronal protective effect of the recombinant tissue protective cytokines of the invention, withdrawal of NGF in differentiated PC12 cells was used as a model. The assay is a well-established model of apoptosis. This PC12 rat cell line was derived from an adrenal medullary pheochromocytoma and can be differentiated into neuronal-like cells in the presence of NGF (Masuda et al., 1993, J Biol Chem 268, 11208-11216). The PC12 cell line is a neuroendocrine cell line, which in the presence of NGF can be differentiated to express a neuronal-like phenotype (Vaudry et al., 2002, Science 296, 1648-1649). Once the cells are fully differentiated they become NGF-dependent and withdrawal of NGF induces apoptosis.

PC12 cells were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% heat inactivated horse serum, 5% heat inactivated fetal bovine serum, 1% sodium-pyruvate and 1% penicillin-streptomycin (P/S) (Invitrogen, Carlsbad, USA).

For experiments, cells were differentiated for 7 days in collagen G-coated 48 well plates at a density of 24,000 cells/well in DMEM supplemented with 1% heat inactivated horse serum, 1% sodium-pyruvate, 1% P/S and 100ng/ml NGF (7S nerve growth factor, mouse submaxillary glands, purchased from Calbiochem, Cat. No.480354) with medium changed every 2-3 days. At day 6, the Epo mutant at amino acid 100 (=S100E) was added to the cells in the indicated concentrations for 24 hours, after which medium was replaced with RPMI1640, 1% P/S, to remove NGF from all cells. S100E was re-added, as was NGF (100 ng/ml) as positive control (+NGF). After 24 h, viability was measured by a tetrazolium (MTT)-reduction assay.

Figure 33A and 33B Show the effect of pre-incubation with S100E in differentiated PC12 cells submitted to NGF withdrawal in two independent experiments. Differentiated PC12 cells were pre-treated with S100E at the indicated concentrations for 24 h, Figure 33A (3 pM) Figure 33B (0.00003 pM-3pM). Viability was measured in the MTT assay. NGF (100 ng/ml) was used as a positive control and NGF-free medium (-NGF) as a negative control. Data presented in Figure 33 are presented as % viability of positive control (+NGF) (n=8 in both experiments). There is a statistically significant increase in viability of S100E treated cells compared to negative control cells (-NGF) by use of one-way ANOVA and Bonferroni post-hoc test. ***p<0.001, *p<0.05. The effects observed with S100E were similar to those of Epo in this test system with respect to potency and efficacy.

Figure 34 Shows the effect of pre-incubation with Epo in differentiated PC12 cells submitted to NGF withdrawal. Differentiated PC12 cells were pre-treated with Epo, S100E, or carbamylated Epo (30 pM-30 nM) for 24 h. The chemically modified Epo molecule, AA24496, has a 10000 times lower activity than EPO in the UT-7 cell assay. Viability was measured in the MTT assay. NGF (100 ng/ml) was used as a positive control and NGF-free medium (-NGF) as a negative control.

6.17. EXAMPLE 17: EPO BIO-ASSAY UT-7 CELL PROLIFERATION

UT-7 is a leukaemia Epo-dependent cell line used for the determination of the erythroid effect of recombinant tissue protective cytokine such as K45D. The UT-7 cells (Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ), Cat. No. ACC 363) were normally grown in the presence of 10% FBS and 5 ng/ml Epo. The proliferation/survival (= viability increase) response of the cells exposed to Epo is mediated by the classical peripheral-type Epo receptor. The proliferation response is a quantitative

measure of and correlates with the capacity of Epo-variants to stimulate the classical Epo receptor.

Methods for UT-7 cell Viability assay

The human leukemia cell line UT7 was made Epo dependent, and the proliferative
5 response to added Epo/recombinant tissue protective cytokines was used as a measurement for their biological activity. On day one of the assay the cells were transferred to fresh complete RPMI 1640 media with 10% serum containing Epo (5 ng/ml) (10% donor calf serum, 4mM L-glutamine, supplemented with 5ng/ml of rhuEPO). The cells were grown in the 75cm² flasks with 20 ml of culture/flask. On day two of the assay the cells were
10 transferred from the flask(s) into a 50-ml conical tube and centrifuge at 1,000 rpm for 5 minutes at room temperature. The old media was discard and the cells were washed two times with 10 ml of starvation media (3% donor calf serum, 4mM L-glutamine). The cells were re-suspended in starvation media, using pipet action up and down to obtain a single cell suspension. To determine the cell density, the re-suspended cells were diluted with
15 starvation media to a density of 4×10^5 cells/ml with a total culture volume of 10 ml and placed in a 25 cm² flask. The mixture was incubated for 4 h in a humidified incubator with 5% CO₂ at 37°C. During the last hour of incubation, a 96 well plate was prepared. At the end of the 4-hour incubation, the cell cultures were removed from the incubator, and the cells were transferred from a flask to a 50-ml conical tube. The contents were mixed by
20 hand to keep the cells suspended. 50 ml of starvation media was added as the media blank without cells. Five wells were the control cells without reagent. The next adjacent row of wells contained the lowest concentration of recombinant tissue protective cytokines. Each adjacent row of wells thereafter was filled with sequentially greater concentrations. The cell cultures that were incubated in media with 3% serum and without Epo were plated out
25 at 200,000 cells/ml and 100µl per well in 96-well plates. The contents were mixed briefly and carefully, using the orbital vibrating platform seated on top of the stir plate. The plate was incubated with different concentrations of Epo variants (from 0.2 pM to 20 nM) for 48 h in RPMI 1640 medium containing 3% serum in a humidified incubator with 5% CO₂ at 37°C. On day four of the assay, the 96-well plate was taken out from the Incubator and
30 placed at room temperature in the laminar flow hood. Immediately, the bioactivity is quantified (spectrophotometric absorption at 450 nm, subtracted from background absorption at 620 nm) by measuring the formazan product formed during cellular

metabolism of the tetrazolium dye WST1, which correlates with cellular viability/number of cells.

Results

The UT7 cells showed stable and reliable growth in Epo containing media for 3
5 months.

K45D induced a viability increase of the Epo-dependent UT-7 cells in a dose dependent way, with an EC_{50} of 294.0. In comparison, the EC_{50} was 58.13 for Epo (Figure 35) and 608 for His-tagged Epo (EpoWT). S100E did not increase viability (more than 50%) of the Epo-dependent UT-7 cells at concentrations < 50 nM (i.e. within the
10 measurable range). Hence, K45D showed potency within the same order of magnitude as Epo, while S100E showed at least 1000-fold lower potency as compared to Epo.

R103E did not increase survival of the Epo-dependent UT-7 cells at concentrations up to 20nM, i.e. its potency compared to Epo was at least four orders of magnitude lower. R150E induced survival of the Epo-dependent UT-7 cells in a dose dependent way, with an
15 EC_{50} of 20 nM. In comparison, the EC_{50} was 66.5 for Epo (Epo#4) (Figure 36). Hence, R150E showed three orders of magnitude lower potency as compared to Epo.

Figure 35 shows concentration-response curves of Epo, K45D and S100E in UT-7 cells. Different concentrations of Epo, EpoWT, K45D and S100E were added to UT-7 cells. Viability was measured 48 h later in the WST-1 assay. Data are mean \pm SD of three
20 different experiments each performed in duplicate. The curve is a non-linear regression curve fit.

Figure 36 shows dose response curves of Epo, R103E and R150E in UT-7 cells. Different concentrations of Epo, EpoWT, R103E and R150E were added to UT-7 cells. Viability was measured 48 h later in the WST-1 assay. Data are mean \pm SD of three
25 different experiments each performed in duplicate. The curve is a non-linear regression curve fit.

6.18. EXAMPLE 18: PROTECTION OF RETINAL ISCHEMIA BY PERIPHERALLY-ADMINISTERED RECOMBINANT TISSUE PROTECTIVE CYTOKINES.

As described in Section 6.9, retinal cells are very sensitive to ischemia such that many will die after 30 minutes of ischemic stress. In this experiment, the rat reversible glaucoma model was again utilized as described by Rosenbaum *et al.* (1997; Vis. Res. 37:3443-51). The effects of recombinant tissue protective cytokines on ischemic stress were examined.

One eye in each of the rats was injured in accordance with the protocol outlined in the example presented in Section 6.9 for saline injection into the anterior chamber of the adult male rat eye. At the time of reperfusion, i.e. when the pressure in the anterior chamber of the eye is released, the rats were administered 10 μ g/kg of EPO, one of four recombinant tissue protective cytokines: R103E, R150E, S100E, and S100e/K45D, or saline intravenously. On days 1, 3, 5 and 6 following the injury, electroretinograms were performed on both the injured and normal eye of each rat. The latency in the damaged eye of each rat was compared to the latency in the normal eye of the same rat. The data was recorded as a ratio of the latency of the injured eye over the latency the normal eye resulting in a ratio of one when the damaged eye has normal function. There are two components to the injury results: Amplitude (the difference from the peak to the trough as shown in Figure 17, Panel A, indicated by 'b' and Latency, the time that it takes to achieve the peak in response to the stimulus.

Figure 38 shows the ratio of the latency of the injured eye over the latency the normal eye for the various treatment regimens. The rat treated with EPO exhibited a latency of 1.2, which is better than the rat treated with saline. Each of the four recombinant tissue protective cytokines resulted in latency results equal to or better than EPO with R103E, R150E, and S100E showing a statistical improvement over saline.

The invention is not to be limited in scope by the specific embodiments described which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

All references cited herein are incorporated by reference herein in their entireties for all purposes.

WHAT IS CLAIMED IS:

1. A mutein recombinant tissue protective cytokine lacking at least one activity selected from the group consisting of increasing hematocrit, vasoactive action, hyperactivating platelets, pro-coagulant activities and increasing production of
5 thrombocytes, the cytokine comprising at least one responsive cellular protective activity selected from the group consisting of protecting, maintaining, enhancing or restoring the function or viability of a responsive mammalian cell, tissue or organ.
2. The recombinant tissue protective cytokine of claim 1, comprising one or more altered amino acid residue between position 11 to 15 of SEQ ID NO:10 [SEQ ID NO:1],
10 position 44 to 51 of SEQ ID NO 10 [SEQ ID NO:2], position 100-108 of SEQ ID NO [SEQ ID NO:3], or position 146-151 of SEQ ID NO 10 [SEQ ID NO:4].
3. The recombinant tissue protective cytokine of claim 1, comprising an altered amino acid residue at one or more of the following positions of SEQ ID NO: 10: 7, 20, 21, 29, 33, 38, 42, 59, 63, 67, 70, 83, 96, 126, 142, 143, 152, 153, 155, 156, or 161.
- 15 4. The recombinant tissue protective cytokine of claim 1, comprising the amino acid sequence of SEQ ID NO: 10 with one or more of the amino acid residue substitutions of SEQ ID NOs: 15-105 and 119.
5. The recombinant tissue protective cytokine of claim 1, comprising the amino acid sequence of SEQ ID NO: 10 with a deletion of amino acid residues 44-49 of SEQ ID NO:
20 10.
6. The recombinant tissue protective cytokine of claim 1, comprising, the amino acid sequence of SEQ ID NO: 10 with at least one of the following amino acid residue substitutions of SEQ ID NOs: 106-118.
7. The recombinant tissue protective cytokine of any one of claims 1-6, further
25 comprising a chemical modification of one or more amino acids.
8. The recombinant tissue protective cytokine of claim 7, wherein the chemical modification comprises altering the charge of the recombinant tissue protective cytokine.

9. The recombinant tissue protective cytokine of claim 8, wherein a positive or negative charge is chemically added to an amino acid residue where a charged amino acid residue is modified to an uncharged residue.
10. The recombinant tissue protective cytokine of any one of claims 1-6, wherein said
5 cytokine is a human erythropoietin mutein.
11. The recombinant tissue protective cytokine of any one of claims 1-6, wherein said cytokine is a human phenylglyoxal erythropoietin mutein.
12. The recombinant tissue protective cytokine of any one of claims 1-6, wherein the responsive mammalian cell comprises a neuronal, muscle, heart, lung, liver, kidney, small
10 intestine, adrenal cortex, adrenal medulla, capillary, endothelial, testis, ovary, endometrial, or stem cell.
13. The recombinant tissue protective cytokine responsive mammalian cell of any one of claims 1-6, comprising a photoreceptor, ganglion, bipolar, horizontal, amacrine, Muieller, myocardium, pace maker, sinoatrial node, sinus node, atrioventricular node, bundle of His,
15 hepatocyte, stellate, Kupffer, mesangial, goblet, intestinal gland, enteral endocrine, glomerulosa, fasciculate, reticularis, chromaffin, pericyte, Leydig, Sertoli, sperm, Graffian follicles, primordial follicles, endometrial stroma, and endometrial cell.
14. The recombinant tissue protective cytokine of any one of claims 1-6, wherein said cytokine is capable of traversing an endothelial cell barrier.
- 20 15. The recombinant tissue protective cytokine of claim 14, wherein the endothelial cell barrier comprises the blood-brain barrier, the blood-eye barrier, the blood testes barrier, the blood-ovary barrier, and the blood-uterus barrier.
16. The recombinant tissue protective cytokine of any one of claims 1-6, wherein said cytokine is selected from the group consisting of:
- 25 i. a cytokine having a reduced number or no sialic acid moieties;
ii. a cytokine having a reduced number or no N-linked or O-linked carbohydrates;
iii. a cytokine having at least a reduced carbohydrate content by virtue of treatment of native cytokine with at least one glycosidase;

- iv. a cytokine having at least one or more oxidized carbohydrates;
 - v. a cytokine having at least one or more oxidized carbohydrates and is chemically reduced;
 - vi. a cytokine having at least one or more modified arginine residues;
 - 5 vii. a cytokine having at least one or more modified lysine residues or a modification of the N-terminal amino group of a cytokine molecule;
 - viii. a cytokine having at least a modified tyrosine residue;
 - ix. a cytokine having at least a modified aspartic acid or glutamic acid residue;
 - x. a cytokine having at a modified tryptophan residue;
 - 10 xi. a cytokine having at least one amino acid group removed;
 - xii. a cytokine having at least one opening of at least one of the cystine linkages in the cytokine molecule;
 - xiii. a truncated cytokine;
 - xiv. a cytokine having at least one polyethylene glycol molecule attached;
 - 15 xv. a cytokine having at least one fatty acid attached;
 - xvi. a cytokine having a non-mammalian glycosylation pattern by virtue of the expression of a recombinant cytokine in non-mammalian cells; and
 - xvi. a cytokine having at least one histidine tagged amino acid to facilitate purification.
- 20 17. The recombinant tissue protective cytokine of claim 16 wherein said cytokine is an asialoerythropoietin.
18. The recombinant tissue protective cytokine of claim 17, wherein said asialoerythropoietin is human asialoerythropoietin.
19. The recombinant tissue protective cytokine of claim 16, wherein said cytokine is
25 hyposialylated or hypersialylated.
20. The recombinant tissue protective cytokine of claim 16, wherein said cytokine comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, or 13 sialic acid moieties.
21. The recombinant tissue protective cytokine of claim 16, wherein said cytokine comprises more than the fourteen sialic acid moieties present in native erythropoietin.

22. The recombinant tissue protective cytokine of claim 16, wherein said cytokine is an erythropoietin with no N-linked carbohydrates.
23. The recombinant tissue protective cytokine of claim 22, wherein said cytokine is an erythropoietin with no O-linked carbohydrates.
- 5 24. The recombinant tissue protective cytokine of claim 16, wherein said cytokine is treated with at least one glycosidase.
25. The recombinant tissue protective cytokine of claim 16, wherein said cytokine is periodate-oxidized erythropoietin.
26. The recombinant tissue protective cytokine of claim 25, wherein said periodate-oxidized erythropoietin is chemically reduced with sodium cyanoborohydride.
- 10 27. The recombinant tissue protective cytokine of claim 16, wherein said cytokine comprises an R-glyoxal moiety on the one or more arginine residues, wherein R is aryl or alkyl moiety.
28. The recombinant tissue protective cytokine of claim 27, wherein said cytokine is phenylglyoxal-erythropoietin.
- 15 29. The recombinant tissue protective cytokine of claim 16, wherein said cytokine is an erythropoietin in which an arginine residue is modified by reaction with a vicinal diketone selected from the group consisting of 2,3-butanedione and cyclohexanedione.
30. The recombinant tissue protective cytokine of claim 16, wherein said cytokine is an erythropoietin in which an arginine residue is reacted with 3-deoxyglucosone.
- 20 31. The recombinant tissue protective cytokine of claim 16, wherein said cytokine is a molecule having at least one biotinylated lysine or N-terminal amino group.
32. The recombinant tissue protective cytokine of claim 16, wherein said cytokine is a glucitolyl lysine erythropoietin or fructosyl lysine erythropoietin.
- 25 33. The recombinant tissue protective cytokine of claim 16, wherein said cytokine comprises at least one carbamylated lysine residue.

34. The recombinant tissue protective cytokine of claim 33, wherein said carbamylated cytokine is comprised of alpha-N-carbamoylerythropoietin; N-epsilon-carbamoylerythropoietin; alpha-N-carbamoyl, N-epsilon-carbamoylerythropoietin; alpha-N-carbamoylasialoerythropoietin; N-epsilon-carbamoylasialoerythropoietin; alpha-N-carbamoyl, N-epsilon-carbamoylasialoerythropoietin; alpha-N-carbamoylhyposialoerythropoietin; N-epsilon-carbamoylhyposialoerythropoietin; and alpha-N-carbamoyl, N-epsilon-carbamoylhyposialoerythropoietin.
35. The recombinant tissue protective cytokine of claim 16, wherein said cytokine comprises at least one acylated lysine residue.
36. The recombinant tissue protective cytokine of claim 35, wherein said cytokine comprises at least one acylated lysine residue.
37. The recombinant tissue protective cytokine of claim 36, wherein said cytokine comprises at least one acylated lysine residue.
38. The recombinant tissue protective cytokine of claim 37, wherein a said acetylated cytokine is comprised of alpha-N-acetylerythropoietin; N-epsilon-acetylerythropoietin; alpha-N-acetyl, N-epsilon-acetylerythropoietin; alpha-N-acetylasialoerythropoietin; N-epsilon-acetylasialoerythropoietin; alpha-N-acetyl, N-epsilon-acetylasialoerythropoietin; alpha-N-acetylhyposialoerythropoietin; N-epsilon-acetylhyposialoerythropoietin; and alpha-N-acetyl, N-epsilon-acetylhyposialoerythropoietin.
39. The recombinant tissue protective cytokine of claim 35, wherein a lysine residue of said cytokine is succinylated.
40. The recombinant tissue protective cytokine of claim 39, wherein said succinylated cytokine is comprised of alpha-N-succinylerythropoietin; N-epsilon-succinylerythropoietin; alpha-N-succinyl, N-epsilon-succinylerythropoietin; alpha-N-succinylasialoerythropoietin; N-epsilon-succinylasialoerythropoietin; alpha-N-succinyl, N-epsilon-succinylasialoerythropoietin; alpha-N-succinylhyposialoerythropoietin; N-epsilon-succinylhyposialoerythropoietin; and alpha-N-succinyl, N-epsilon-succinylhyposialoerythropoietin.

41. The recombinant tissue protective cytokine of claim 16, wherein said cytokine comprises at least one lysine residue modified by 2, 4, 6 trinitrobenzenesulfonate sodium or another salt thereof.
42. The recombinant tissue protective cytokine of claim 16, wherein said cytokine
5 comprises at least one nitrated or iodinated tyrosine residue.
43. The recombinant tissue protective cytokine of claim 16, wherein said cytokine comprises an aspartic acid or glutamic acid residue that is reacted with a carbodiimide followed by reaction with an amine.
44. The recombinant tissue protective cytokine of claim 16, wherein a said amine is
10 glycynamide.
45. An isolated nucleic acid molecule that comprises a nucleotide sequence which encodes a polypeptide comprising the recombinant tissue protective cytokine of any one of claims 1-6.
46. A vector comprising a nucleic acid molecule of claim 45.
- 15 47. An expression vector comprising a nucleic acid molecule of claim 45 and at least one regulatory region operably linked to the nucleic acid molecule.
48. The vector of claim 46 or 47 that is a pCiNeo vector.
49. A genetically-engineered cell which comprises a nucleic acid molecule of claim 45.
50. A cell comprising the expression vector of claim 45.
- 20 51. A pharmaceutical composition comprising a recombinant tissue protective cytokine of any one of claims 1-6, lacking at least one activity selected from the group consisting of increasing hematocrit, vasoactive action, hyperactivating platelets, pro-coagulant activities and increasing production of thrombocytes, the cytokine having at least one responsive cellular protective activity selected from the group consisting of protecting, maintaining,
25 enhancing or restoring the function or viability of a responsive mammalian cell, tissue or organ.

52. The pharmaceutical composition of claim 51, formulated for oral, intranasal, or parenteral administration.
53. The pharmaceutical composition of claim 51, formulated as a perfusate solution.
54. A method for protecting, maintaining or enhancing the viability of a cell, tissue or organ isolated from a mammalian body comprising exposing said cell, tissue or organ to a pharmaceutical composition comprising a mutein recombinant tissue protective cytokine.
55. The method of claim 54, wherein the protection does not effect bone marrow.
56. A method for protecting, maintaining or enhancing the viability of a cell, tissue or organ isolated from a mammalian body comprising exposing said cell, tissue or organ to a pharmaceutical composition comprising a recombinant tissue protective cytokine of any one of claims 1-6, that lacks at least one activity selected from the group consisting of increasing hematocrit, vasoactive action, hyperactivating platelets, pro-coagulant activity and increasing production of thrombocytes.
57. Use of a recombinant tissue protective cytokine of any one of claims 1-6, that lacks at least one activity selected from the group consisting of increasing hematocrit, vasoactive action, hyperactivating platelets, pro-coagulant activity and increasing production of thrombocytes, for the preparation of a pharmaceutical composition for the protection against and prevention of a tissue injury as well as the restoration of and rejuvenation of tissue and tissue function in a mammal.
58. The use of claim 57, wherein the injury is caused by a seizure disorder, multiple sclerosis, stroke, hypotension, cardiac arrest, ischemia, myocardial infarction, inflammation, age-related loss of cognitive function, radiation damage, cerebral palsy, neurodegenerative disease, Alzheimer's disease, Parkinson's disease, Leigh disease, AIDS dementia, memory loss, amyotrophic lateral sclerosis, alcoholism, mood disorder, anxiety disorder, attention deficit disorder, autism, Creutzfeld-Jakob disease, brain or spinal cord trauma or ischemia, heart-lung bypass, chronic heart failure, macular degeneration, diabetic neuropathy, diabetic retinopathy, glaucoma, retinal ischemia, or retinal trauma.
59. A method for facilitating the transcytosis of a molecule across an endothelial cell barrier in a mammal comprising administration to said mammal a composition comprising

said molecule in association with a recombinant tissue protective cytokine of any one of claims 1-6, lacking at least one activity selected from the group consisting of increasing hematocrit, increasing blood pressure, hyperactivating platelets, and increasing production of thrombocytes.

5 60. The method of claim 59, wherein said association is a labile covalent bond, a stable covalent bond, or a non-covalent association with a binding site for said molecule.

61. The method of claim 59, wherein said endothelial cell barrier is selected from the group consisting of the blood-brain barrier, the blood-eye barrier, the blood-testis barrier, the blood-ovary barrier, the blood-heart barrier, the blood-kidney barrier, and the blood-placenta barrier.

10

62. The method of claim 59, wherein said molecule is a receptor agonist or antagonist hormone, a neurotrophic factor, an antimicrobial agent, an antiviral agent, a radiopharmaceutical, an antisense oligonucleotide, an antibody, an immunosuppressant, a dye, a marker, or an anti-cancer drug.

15 63. A composition for transporting a molecule via transcytosis across an endothelial cell barrier comprising said molecule in association with a recombinant tissue protective cytokine, of any one of claims 1-6, lacking at least one activity selected from the group consisting of increasing hematocrit, vasoactive action, hyperactivating platelets, pro-coagulant activity and increasing production of thrombocytes.

20 64. The composition of claim 63, wherein said association is a labile covalent bond, a stable covalent bond, or a non-covalent association with a binding site for said molecule.

65. The composition of claim 63, wherein said molecule is a receptor agonist or antagonist hormone, a neurotrophic factor, an antimicrobial agent, a radiopharmaceutical, an antisense oligonucleotide, an antibody, an immunosuppressant, a dye, a marker, or an anti-cancer drug.

25

66. Use of an recombinant tissue protective cytokine of any one of claims 1-6, lacking at least one activity selected from the group consisting of increasing hematocrit, vasoactive action, hyperactivating platelets, pro-coagulant activities and increasing production of thrombocytes.

67. The use of claim 66, wherein said association is a labile covalent bond, a stable covalent bond, or a non-covalent association with a binding site for said molecule.

68. The use of claim 66, wherein said molecule is a receptor agonist or antagonist hormone, a neurotrophic factor, an antimicrobial agent, a radiopharmaceutical, an antisense
5 oligonucleotide, an antibody, an immunosuppressant, a dye, or a marker, or an anti-cancer drug.

1/37




FIG. 1

2/37




FIG.2

4/37



FIG.4

5/37




FIG.5

6/37


SK-N-SH NEUROBLASTOMA CELLS
NEUROPROTECTION ASSAY (AGAINST ROTENONE)

FIG.6A

7/37




FIG.6B

8/37




FIG.7

9/37




FIG.8





FIG.9



n FOR EACH GROUP IS

11/37

PROTECTION OF P19 CELLS FROM SERUM
DEPRIVATION BY IODO-EPO




FIG.11

12/37



BIOTINYLATED-EPO AND ASIALO-EPO RETAIN
ACTIVITY IN P19 ASSAY

FIG.12

SURVIVAL OF P19 CELLS AFTER SERUM WITHDRAWAL






FIG.14

14/37


ERYTHROPOIETIN IMPROVES CARDIAC FUNCTION IN
A HEART ISOLATED FOR TRANSPLANTATION

FIG.15

15/37

RAT HEART 7 DAYS AFTER 30 MINUTES OF ISCHEMIA




FIG.16




FIG.17D




FIG.17C




FIG.17B





FIG.17A

17/37

RETINOGRAM AMPLITUDE AFTER 60 min. ISCHEMIA



* $p < 0.05$
n = 5 PER GROUP

FIG. 18

18/37




FIG.19

20/37




FIG.21

21/37




FIG.22

22/37




FIG.23

23/37




FIG.24A




FIG.24B




FIG.24C

24/37





FIG. 25A



FIG. 25B



25/37




FIG. 26A




FIG. 26B

26/37




FIG.27A




FIG.27B

27/37




FIG.28

28/37




FIG.29




FIG.30A






FIG. 30B

30/37



31/37




FIG.33A




FIG.33B

32/37




FIG.34A




FIG.34B

33/37





FIG.34C

UT7 BIOASSAY
Epo MUTANTS



	Epo#4	K45D	S100E	EpoWT
EC50	58.13	294.0	5.3840e+006	608.0

FIG.35

35/37




FIG.36

36/37




FIG.37

37/37




FIG.38

SEQUENCE LISTING

<110> Kenneth S. Warren Industries, Inc.
H. Lundbeck A/S

<120> RECOMBINANT TISSUE PROTECTIVE CYTOKINES AND ENCODING NUCLEIC
ACIDS.THEREOF FOR PROTECTION, RESTORATION, AND ENHANCEMENT OF
RESPONSIVE CELLS, TISSUES AND ORGANS

<130> 10165-022-228

<140>

<141>

<150> 60/392,455

<151> 2002-07-01

<150> 60/393,423

<151> 2002-07-03

<160> 212

<170> PatentIn version 3.2

<210> 1

<211> 5

<212> PRT

<213> Homo sapiens

<400> 1

Val Leu Gln Arg Tyr
1 5

<210> 2

<211> 8

<212> PRT

<213> Homo sapiens

<400> 2

Thr Lys Val Asn Phe Tyr Ala Trp
1 5

<210> 3

<211> 9

<212> PRT

<213> Homo sapiens

<400> 3

Ser Gly Leu Arg Ser Leu Thr Thr Leu
1 5

<210> 4

<211> 6

<212> PRT

<213> Homo sapiens

<400> 4

Ser Asn Phe Leu Arg Gly
1 5

<210> 5

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 5

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Glu Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 6
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 6

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Asp Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 7
 <211> 580
 <212> DNA
 <213> Homo sapiens

<400> 7
 atgggggtgc acgaatgtcc tgcctggctg tggtctctcc tgtccctgct gtcgctccct 60
 ctgggcctcc cagtcctggg cgtccaccca cgctcatct gtgacagccg agtcctggag 120
 aggtacctct tggaggccaa ggaggccgag aatatcacga cgggctgtgc tgaacactgc 180
 agcttgaatg agaatatcac tgtcccagac accaaagtta atttctatgc ctggaagagg 240
 atggaggtcg ggcagcaggc cgtagaagtc tggcagggcc tggccctgct gtcggaagct 300
 gtccctgcggg gccaggccct gttggtcaac tcttccagc cgtgggagcc cctgcactgc 360
 atgtggataa agccgtcagt ggccttcgca gcctcaccac tctgcttcgg gctctgggag 420
 cccagaagga agccatctcc cctccagatg cggcctcagc tgctccactc cgaacaatca 480
 ctgctgacac ttctgcaaac tcttccgagt ctactccaat ttctccggg gaaagctgaa 540
 gctgtacaca ggggaggcct gcaggacagg ggacagatga 580

<210> 8
 <211> 35
 <212> DNA
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: primer

<400> 8
 agctctcgag gcgcggagat ggggggtgcac gaatg 35

<210> 9
 <211> 36
 <212> DNA
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: primer

<400> 9
 atgctctaga cacacctggt catctgtccc ctgtcc 36

<210> 10

<211> 193
 <212> PRT
 <213> Homo sapiens

<400> 10

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 11
 <211> 45
 <212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 11

catgtggata aagccgtcga gggccttcgc agcctcacca ctctg

45

<210> 12

<211> 45

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 12

cagagtgggtg aggctgcgaa gggcctcgac ggctttatcc acatg

45

<210> 13

<211> 45

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 13

gagaatatca ctgtcccaga caccgacgtt aattttctatg cctgg

45

<210> 14

<211> 45

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 14

ccaggcatag aaattaacgt cgggtgtctgg gacagtgata ttctc

45

<210> 15

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 15

Met	Gly	Val	His	Glu	Cys	Pro	Ala	Trp	Leu	Trp	Leu	Leu	Leu	Ser	Leu
1				5				10						15	

Leu	Ser	Leu	Pro	Leu	Gly	Leu	Pro	Val	Leu	Gly	Ala	Pro	Pro	Arg	Leu
			20					25					30		

Ala Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 16

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 16

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu

20 25 30
 Ile Ala Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60
 Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95
 Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110
 Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125
 Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140
 Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 17
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 17

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Ile Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 18
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 18

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Ser Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 19
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein
 <400> 19

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Ala Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 20
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 20

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu

1	5	10	15
Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu	20	25	30
Ile Cys Asp Ser Arg Val Leu Ala Arg Tyr Leu Leu Glu Ala Lys Glu	35	40	45
Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu	50	55	60
Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg	65	70	75
Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu	85	90	95
Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser	100	105	110
Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly	115	120	125
Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu	130	135	140
Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile	145	150	155
Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu	165	170	175
Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp	180	185	190

Arg

```
<210> 21
<211> 193
<212> PRT
<213> Artificial
```

```
<220>
<223> Description of Artificial Sequence: mutein
<400> 21
```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15
 Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30
 Ile Cys Asp Ser Arg Val Leu Glu Ala Tyr Leu Leu Glu Ala Lys Glu
 35 40 45
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60
 Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95
 Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110
 Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125
 Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140
 Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 22
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein
 <400> 22


```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu
1           5           10           15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
          20           25           30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
          35           40           45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
          50           55           60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65           70           75           80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
          85           90           95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
          100          105          110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
          115          120          125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
          130          135          140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145          150          155          160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
          165          170          175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
          180          185          190

```

Arg

```

<210> 23
<211> 193
<212> PRT
<213> Artificial

```

<220>

<223> Description of Artificial Sequence: mutein

<400> 23

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Glu Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 24

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 24

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Gln Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 25

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 25

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Ala Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 26

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 26

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Phe Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 27

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 27

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Ile Leu Leu Glu Ala Glu Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 28

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 28

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Glu Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 29

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 29

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Ala Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 30

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 30

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Ala
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 31

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 31

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Lys Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 32
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 32

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Ser Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 33
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 33

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Tyr Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 34
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 34

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Asn Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 35
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 35

```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1              5              10              15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
                20              25              30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
          35              40              45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu Thr Cys Ser Leu Asn Glu
      50              55              60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65              70              75              80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
          85              90              95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
          100              105              110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
          115              120              125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
          130              135              140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145              150              155              160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
          165              170              175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
          180              185              190

```

Arg

<210> 36

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 36

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Ser Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 37

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 37

Met	Gly	Val	His	Glu	Cys	Pro	Ala	Trp	Leu	Trp	Leu	Leu	Leu	Ser	Leu
1				5					10					15	

Leu	Ser	Leu	Pro	Leu	Gly	Leu	Pro	Val	Leu	Gly	Ala	Pro	Pro	Arg	Leu
			20					25					30		

Ile	Cys	Asp	Ser	Arg	Val	Leu	Glu	Arg	Tyr	Leu	Leu	Glu	Ala	Lys	Glu
		35					40					45			

Ala	Glu	Asn	Ile	Thr	Thr	Gly	Cys	Ala	Glu	His	Tyr	Ser	Leu	Asn	Glu
	50					55					60				

Asn	Ile	Thr	Val	Pro	Asp	Thr	Lys	Val	Asn	Phe	Tyr	Ala	Trp	Lys	Arg
65					70					75					80

Met	Glu	Val	Gly	Gln	Gln	Ala	Val	Glu	Val	Trp	Gln	Gly	Leu	Ala	Leu
				85					90					95	

Leu	Ser	Glu	Ala	Val	Leu	Arg	Gly	Gln	Ala	Leu	Leu	Val	Asn	Ser	Ser
			100					105					110		

Gln	Pro	Trp	Glu	Pro	Leu	Gln	Leu	His	Val	Asp	Lys	Ala	Val	Ser	Gly
		115					120					125			

Leu	Arg	Ser	Leu	Thr	Thr	Leu	Leu	Arg	Ala	Leu	Gly	Ala	Gln	Lys	Glu
	130					135					140				

Ala	Ile	Ser	Pro	Pro	Asp	Ala	Ala	Ser	Ala	Ala	Pro	Leu	Arg	Thr	Ile
145					150					155					160

Thr	Ala	Asp	Thr	Phe	Arg	Lys	Leu	Phe	Arg	Val	Tyr	Ser	Asn	Phe	Leu
				165					170					175	

Arg	Gly	Lys	Leu	Lys	Leu	Tyr	Thr	Gly	Glu	Ala	Cys	Arg	Thr	Gly	Asp
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

180

185

190

Arg

<210> 38

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 38

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Lys Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 39

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 39

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Lys Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 40

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 40

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Asn Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 41
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 41

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Ala Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu

165 170 175
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 42
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 42

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Ala Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 43
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 43

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Ile Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 44
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 44

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Asp Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 45
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 45

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Ala Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile

145					150					155					160	
Thr	Ala	Asp	Thr	Phe	Arg	Lys	Leu	Phe	Arg	Val	Tyr	Ser	Asn	Phe	Leu	
				165					170					175		
Arg	Gly	Lys	Leu	Lys	Leu	Tyr	Thr	Gly	Glu	Ala	Cys	Arg	Thr	Gly	Asp	
			180					185					190			

Arg

```
<210> 46
<211> 193
<212> PRT
<213> Artificial
```

<220>
<223> Description of Artificial Sequence: mutein

<400> 46

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Ala Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 47

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 47

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Ala Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 48
 <211> 193
 <212> PRT
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 48

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Ile Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 49
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 49

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Ala Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu

130						135				140					
Ala	Ile	Ser	Pro	Pro	Asp	Ala	Ala	Ser	Ala	Ala	Pro	Leu	Arg	Thr	Ile
145					150					155					160
Thr	Ala	Asp	Thr	Phe	Arg	Lys	Leu	Phe	Arg	Val	Tyr	Ser	Asn	Phe	Leu
				165					170					175	
Arg	Gly	Lys	Leu	Lys	Leu	Tyr	Thr	Gly	Glu	Ala	Cys	Arg	Thr	Gly	Asp
			180					185					190		

Arg

```
<210> 50
<211> 193
<212> PRT
<213> Artificial

<220>
<223> Description of Artificial Sequence: mutein

<400> 50
```

Met	Gly	Val	His	Glu	Cys	Pro	Ala	Trp	Leu	Trp	Leu	Leu	Leu	Ser	Leu
1				5					10					15	
Leu	Ser	Leu	Pro	Leu	Gly	Leu	Pro	Val	Leu	Gly	Ala	Pro	Pro	Arg	Leu
			20					25					30		
Ile	Cys	Asp	Ser	Arg	Val	Leu	Glu	Arg	Tyr	Leu	Leu	Glu	Ala	Lys	Glu
		35					40					45			
Ala	Glu	Asn	Ile	Thr	Thr	Gly	Cys	Ala	Glu	His	Cys	Ser	Leu	Asn	Glu
	50					55					60				
Asn	Ile	Thr	Val	Pro	Asp	Thr	Lys	Val	Asn	Phe	Ala	Ala	Trp	Lys	Arg
65					70					75					80
Met	Glu	Val	Gly	Gln	Gln	Ala	Val	Glu	Val	Trp	Gln	Gly	Leu	Ala	Leu
				85					90					95	
Leu	Ser	Glu	Ala	Val	Leu	Arg	Gly	Gln	Ala	Leu	Leu	Val	Asn	Ser	Ser
			100					105					110		
Gln	Pro	Trp	Glu	Pro	Leu	Gln	Leu	His	Val	Asp	Lys	Ala	Val	Ser	Gly
		115					120					125			

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 51
 <211> 193
 <212> PRT
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 51

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Ser Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 52
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 52

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Phe Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 53
 <211> 193
 <212> PRT
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 53

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Asn Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly

115 120 125
 Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140
 Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 54
 <211> 193
 <212> PRT
 <213> Artificial
 <220>
 <223> Description of Artificial Sequence: mutein
 <400> 54

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15
 Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30
 Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60
 Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Ala Arg
 65 70 75 80
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95
 Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 55

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 55

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Asn Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 56
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 56

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Thr Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 57

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 57

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Ser Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser

100 105 110
 Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125
 Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140
 Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 58
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein
 <400> 58

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15
 Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30
 Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60
 Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Ala Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 59
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 59

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Arg Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 60
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 60

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Ala Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 61

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 61

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu


```

      85              90              95
Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
      100              105              110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Arg Gly
      115              120              125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
      130              135              140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
      145              150              155              160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
      165              170              175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
      180              185              190

```

Arg

<210> 62

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 62

```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1              5              10              15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
      20              25              30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
      35              40              45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
      50              55              60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65              70              75              80

```

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Glu Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 63

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 63

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ala Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 64

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 64

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Thr Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 65
<211> 193
<212> PRT
<213> Artificial

<220>
<223> Description of Artificial Sequence: mutein

<400> 65

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg

```

65              70              75              80
Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
      85              90              95
Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
      100             105             110
Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Ala
      115             120             125
Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
      130             135             140
Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
      145             150             155             160
Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
      165             170             175
Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
      180             185             190

```

Arg

```

<210> 66
<211> 193
<212> PRT
<213> Artificial

```

```

<220>
<223> Description of Artificial Sequence: mutein

```

```

<400> 66

```

```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1              5              10              15
Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
      20              25              30
Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
      35              40              45
Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
      50              55              60

```

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Ile
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 67
<211> 193
<212> PRT
<213> Artificial

<220>
<223> Description of Artificial Sequence: mutein

<400> 67

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115 120 125

Ala Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 68
<211> 193
<212> PRT
<213> Artificial

<220>
<223> Description of Artificial Sequence: mutein

<400> 68

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115 120 125

Leu Ala Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 69
<211> 193
<212> PRT
<213> Artificial

<220>
<223> Description of Artificial Sequence: mutein

<400> 69

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu


```

50              55              60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65              70              75              80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85              90              95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100             105             110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115             120             125

Leu Glu Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130             135             140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145             150             155             160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165             170             175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180             185             190

```

Arg

<210> 70

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 70

```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1              5              10              15

```

```

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20              25              30

```

```

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35              40              45

```

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115 120 125

Leu Arg Ala Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 71

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 71

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ile Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 72

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 72

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115 120 125

Leu Arg Ser Ala Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 73

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 73

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu

35 40 45
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60
 Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95
 Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110
 Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125
 Leu Arg Ser Leu Ala Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140
 Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 74

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 74

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115 120 125

Leu Arg Ser Leu Ile Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 75

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 75

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Ala Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 76

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 76

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60
 Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95
 Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110
 Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125
 Leu Arg Ser Leu Thr Leu Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140
 Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 77
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 77

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu


```

                20                25                30
Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
   35                40                45
Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
   50                55                60
Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
   65                70                75                80
Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
   85                90                95
Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
  100                105                110
Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
  115                120                125
Leu Arg Ser Leu Thr Thr Lys Leu Arg Ala Leu Gly Ala Gln Lys Glu
  130                135                140
Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
  145                150                155                160
Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
  165                170                175
Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
  180                185                190

```

Arg

<210> 78

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 78

```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu
1          5          10          15

```

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Ala Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 79

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 79

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Ser Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 80
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 80

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Ala Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 81
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 81

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu

1				5					10						15
Leu	Ser	Leu	Pro	Leu	Gly	Leu	Pro	Val	Leu	Gly	Ala	Pro	Pro	Arg	Leu
			20					25					30		
Ile	Cys	Asp	Ser	Arg	Val	Leu	Glu	Arg	Tyr	Leu	Leu	Glu	Ala	Lys	Glu
		35					40					45			
Ala	Glu	Asn	Ile	Thr	Thr	Gly	Cys	Ala	Glu	His	Cys	Ser	Leu	Asn	Glu
	50					55					60				
Asn	Ile	Thr	Val	Pro	Asp	Thr	Lys	Val	Asn	Phe	Tyr	Ala	Trp	Lys	Arg
65					70					75					80
Met	Glu	Val	Gly	Gln	Gln	Ala	Val	Glu	Val	Trp	Gln	Gly	Leu	Ala	Leu
				85					90					95	
Leu	Ser	Glu	Ala	Val	Leu	Arg	Gly	Gln	Ala	Leu	Leu	Val	Asn	Ser	Ser
			100					105					110		
Gln	Pro	Trp	Glu	Pro	Leu	Gln	Leu	His	Val	Asp	Lys	Ala	Val	Ser	Gly
		115					120					125			
Leu	Arg	Ser	Leu	Thr	Thr	Leu	Leu	Arg	Ala	Leu	Gly	Ala	Gln	Lys	Glu
	130					135					140				
Ala	Ile	Ser	Pro	Pro	Asp	Ala	Ala	Ala	Ala	Ala	Pro	Leu	Arg	Thr	Ile
145					150						155				160
Thr	Ala	Asp	Thr	Phe	Arg	Lys	Leu	Phe	Arg	Val	Tyr	Ser	Asn	Phe	Leu
				165					170					175	
Arg	Gly	Lys	Leu	Lys	Leu	Tyr	Thr	Gly	Glu	Ala	Cys	Arg	Thr	Gly	Asp
			180					185					190		

Arg

```
<210> 82
<211> 193
<212> PRT
<213> Artificial
```

```
<220>
<223> Description of Artificial Sequence: mutein
<400> 82
```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Ala Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 83
 <211> 193
 <212> PRT
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 83

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ala
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 84

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 84

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Ala Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 85

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 85

```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1           5           10           15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
          20           25           30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
          35           40           45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
          50           55           60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65           70           75           80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
          85           90           95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
          100          105          110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
          115          120          125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
          130          135          140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145           150           155           160

Thr Ala Asp Thr Phe Arg Ala Leu Phe Arg Val Tyr Ser Asn Phe Leu
          165          170          175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
          180          185          190

```

Arg

<210> 86

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 86

```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1           5           10           15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
          20           25           30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
          35           40           45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
          50           55           60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65           70           75           80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
          85           90           95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
          100          105          110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
          115          120          125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
          130          135          140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
          145          150          155          160

Thr Ala Asp Thr Phe Arg Lys Leu Ile Arg Val Tyr Ser Asn Phe Leu
          165          170          175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
          180          185          190

Arg

```

<210> 87

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 87

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Ala Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 88

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 88

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ala Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 89

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 89

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Lys Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 90

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 90

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Ala Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 91

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 91

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Tyr Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 92
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 92

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Ala Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 93
 <211> 193
 <212> PRT
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 93

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Ala
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 94
 <211> 193
 <212> PRT
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 94

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Ala Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 95
 <211> 193
 <212> PRT
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 95

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Glu Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 96

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 96

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Ala Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 97
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 97

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Ala Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 98

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 98

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Trp Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp

180

185

190

Arg

<210> 99

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 99

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Ala Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 100

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 100

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Ala Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 101

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 101

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Ala Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 102

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 102

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu

				165					170					175				
Arg	Gly	Lys	Leu	Lys	Leu	Tyr	Thr	Ala	Glu	Ala	Cys	Arg	Thr	Gly	Asp			
			180					185					190					

Arg

```
<210> 103
<211> 193
<212> PRT
<213> Artificial
```

<220>
<223> Description of Artificial Sequence: mutein

<400> 103

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Ser Arg Thr Gly Asp
 180 185 190

Arg

<210> 104
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 104

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Ala Arg Thr Gly Asp
 180 185 190

Arg

<210> 105

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 105

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Ala Thr Gly Asp
 180 185 190

Arg

<210> 106
 <211> 192
 <212> PRT
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 106

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Glu His Cys Ser Leu Asn Glu Asn
 50 55 60

Ile Thr Val Pro Asp Thr Asp Val Asn Phe Tyr Ala Trp Lys Arg Met
 65 70 75 80

Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu Leu
 85 90 95

Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser Gln
 100 105 110

Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Glu Gly Leu
 115 120 125

Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu Ala
 130 135 140

Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile Thr

```

145                      150                      155                      160

Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu Arg
                165                      170                      175

Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp Arg
                180                      185                      190

<210> 107
<211> 193
<212> PRT
<213> Artificial

<220>
<223> Description of Artificial Sequence: mutein

<400> 107

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1                      5                      10                      15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
                20                      25                      30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
                35                      40                      45

Ala Glu Asn Ile Thr Thr Gly Cys Asn Glu Thr Cys Ser Leu Asn Glu
                50                      55                      60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65                      70                      75                      80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
                85                      90                      95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
                100                      105                      110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
                115                      120                      125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
                130                      135                      140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145                      150                      155                      160

```

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 108
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 108

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Asp Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Glu Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 109
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 109

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Glu Ser Leu Thr Thr Ser Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 110
 <211> 193
 <212> PRT
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 110

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Ala Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile

145 150 155 160

Thr Ala Asp Thr Phe Arg Ala Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 111

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 111

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Ala Val Asn Phe Tyr Ala Trp Ala Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Ala Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 112

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 112

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 113
<211> 193
<212> PRT
<213> Artificial

<220>
<223> Description of Artificial Sequence: mutein

<400> 113

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Ala Ala Val Ser Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
165 170 175

Arg Gly Ala Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
180 185 190

Arg

<210> 114

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 114

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
50 55 60

Asn Ile Thr Val Pro Asp Thr Ala Val Asn Phe Tyr Ala Trp Lys Arg
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Ala Ala Val Ser Gly
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu

130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Ala Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 115
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 115

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Ala Val Asn Phe Tyr Ala Trp Ala Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Ala Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Ala Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Ala Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 116
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein

<400> 116

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Ala Val Asn Phe Tyr Ala Trp Ala Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Ala Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Ala Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Ala Leu Ala Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 117
 <211> 193
 <212> PRT
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: mutein
 <400> 117

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Lys Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Lys Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Lys Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 118
 <211> 193
 <212> PRT
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 118

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Ala Leu Leu Glu Ala Lys Glu
 35 40 45

Ala Glu Lys Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly

115 120 125
 Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140
 Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 119
 <211> 193
 <212> PRT
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 119

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15
 Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30
 Ile Cys Asp Ser Arg Val Leu Glu Ala Ala Leu Leu Glu Ala Lys Glu
 35 40 45
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
 50 55 60
 Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
 65 70 75 80
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
 85 90 95
 Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
 180 185 190

Arg

<210> 120
 <211> 36
 <212> DNA
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: primer

<400> 120
 gtctactcca atttcctcga gggaaagctg aagctg

36

<210> 121
 <211> 34
 <212> DNA
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: primer

<400> 121
 gcttcagctt tccctcgagg aaattggagt agac

34

<210> 122
 <211> 32
 <212> DNA
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: primer

<400> 122
 ccgtcagtg ccttgagagc ctcaccactc tg

32

<210> 123

<211> 32
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 123
cagagtgggtg aggctctcaa ggccactgac gg 32

<210> 124
<211> 32
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 124
ccgtcagtg ccttgagagc ctcaccactc tg 32

<210> 125
<211> 32
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 125
cagagtgggtg aggctctcaa ggccactgac gg 32

<210> 126
<211> 31
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 126
cgcagcctca ccacttcgct tcgggctctg g 31

<210> 127
<211> 31
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 127
ccagagcccg aagcgaagtg gtgaggctgc g 31

<210> 128
<211> 40

<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 128
gaatatcact gtcccagacg gtggtgctg gaagaggatg 40

<210> 129
<211> 40
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 129
catcctcttc caggcaccac cgtctgggac agtgatattc 40

<210> 130
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 130
tacctcttgg aggcgcgga ggccgagaat atc 33

<210> 131
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 131
gatattctcg gcctccgcg cctccaagag gta 33

<210> 132
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 132
gctgacactt tccgcgcact cttccgagtc tactc 35

<210> 133
<211> 35
<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 133

gagtagactc ggaagagtgc gcggaaagtg tcagc 35

<210> 134

<211> 33

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 134

atttcctccg gggagcgcgtg aagctgtaca cag 33

<210> 135

<211> 33

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 135

ctgtgtacag cttcagcgcct ccccgaggga aat 33

<210> 136

<211> 32

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 136

ctccggggaa agctggcgcgt gtacacaggg ga 32

<210> 137

<211> 32

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 137

tcccctgtgt acagcgccag ctttccccgg ag 32

<210> 138

<211> 35

<212> DNA

<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 138
actgtcccag acaccgcagt taattttctat gcctg 35

<210> 139
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 139
caggcataga aattaactgc ggtgtctggg acagt 35

<210> 140
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 140
agttaatttc tatgcctggg cgaggatgga ggtcg 35

<210> 141
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 141
cgacctccat cctcgcccag gcatagaaat taact 35

<210> 142
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 142
tgcagctgca tgtggatgca gccgtcagtg gcc 33

<210> 143
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 143
ggccactgac ggctgcatcc acatgcagct gca 33

<210> 144
<211> 32
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 144
ctctgggagc ccaggcggaa gccatctccc ct 32

<210> 145
<211> 32
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 145
aggggagatg gcttccgcct gggctcccag ag 32

<210> 146
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 146
gctgacactt tccgcgcact cttccgagtc tactc 35

<210> 147
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 147
gagtagactc ggaagagtgc gcggaaagtg tcagc 35

<210> 148
<211> 35
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 148

agttaatttc tatgcctggg cgaggatgga ggtcg

35

<210> 149

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 149

cgacctccat cctcgcccag gcatagaaat taact

35

<210> 150

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 150

gctgacactt tccgcgcact cttccgagtc tactc

35

<210> 151

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 151

gagtagactc ggaagagtgc gcggaaagtg tcagc

35

<210> 152

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 152

agttaatttc tatgcctggg cgaggatgga ggtcg

35

<210> 153

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 153
cgacctccat cctcgcccag gcatagaaat taact 35

<210> 154
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 154
actgtcccag acaccgcagt taattttatat gcctg 35

<210> 155
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 155
caggcataga aattaactgc ggtgtctggg acagt 35

<210> 156
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 156
tgcagctgca tgtggatgca gccgtcagtg gcc 33

<210> 157
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 157
ggccactgac ggctgcatcc acatgcagct gca 33

<210> 158
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 158
atttcctccg gggagcgctg aagctgtaca cag 33

<210> 159
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 159
ctgtgtacag cttcagcgct ccccgaggga aat 33

<210> 160
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 160
tgcagctgca tgtggatgca gccgtcagtg gcc 33

<210> 161
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 161
ggccactgac ggctgcatcc acatgcagct gca 33

<210> 162
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 162
atttcctccg gggagcgctg aagctgtaca cag 33

<210> 163
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 163

ctgtgtacag cttcagcgct ccccgaggga aat 33

<210> 164

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 164

actgtcccag acaccgcagt taatttctat gcctg 35

<210> 165

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 165

caggcataga aattaactgc ggtgtctggg acagt 35

<210> 166

<211> 33

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 166

tgcagctgca tgtggatgca gccgtcagtg gcc 33

<210> 167

<211> 33

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 167

ggccactgac ggctgcatcc acatgcagct gca 33

<210> 168

<211> 33

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 168

atttcctccg gggagcgctg aagctgtaca cag 33

<210> 169
<211> 33
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 169
ctgtgtacag cttcagcgct ccccgaggga aat

33

<210> 170
<211> 35
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 170
actgtcccag acaccgcagt taattttctat gcctg

35

<210> 171
<211> 35
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 171
caggcataga aattaactgc ggtgtctggg acagt

35

<210> 172
<211> 35
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 172
agttaatttc tatgcctggg cgaggatgga ggtcg

35

<210> 173
<211> 35
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 173
cgacctccat cctcgcccag gcatagaaat taact

35

<210> 174
<211> 33
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 174
tgcagctgca tgtggatgca gccgtcagtg gcc

33

<210> 175
<211> 33
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 175
ggccactgac ggctgcatcc acatgcagct gca

33

<210> 176
<211> 33
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 176
atttcctccg gggagcgcgtg aagctgtaca cag

33

<210> 177
<211> 33
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 177
ctgtgtacag cttcagcgt ccccgaggga aat

33

<210> 178
<211> 35
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 178
actgtccag acaccgcagt taattttctat gcctg

35

<210> 179
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 179
cagggcataga aattaactgc ggtgtctggg acagt 35

<210> 180
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 180
agttaatttc tatgcctggg cgaggatgga ggtcg 35

<210> 181
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 181
cgacctccat cctcgcccag gcatagaaat taact 35

<210> 182
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 182
gctgacactt tccgcgcact cttccgagtc tactc 35

<210> 183
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 183
gagtagactc ggaagagtgc gcggaaagtg tcagc 35

<210> 184

<211> 33
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 184
tgcagctgca tgtggatgca gccgtcagtg gcc

33

<210> 185
<211> 33
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 185
ggccactgac ggctgcatcc acatgcagct gca

33

<210> 186
<211> 33
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 186
atttcctccg gggagcgctg aagctgtaca cag

33

<210> 187
<211> 33
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 187
ctgtgtacag cttcagcgct ccccgaggga aat

33

<210> 188
<211> 35
<212> DNA
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 188
actgtcccag acaccgcagt taatttctat gcctg

35

<210> 189
<211> 35

<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 189
caggcataga aattaactgc ggtgtctggg acagt 35

<210> 190
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 190
agttaatttc tatgcctggg cgaggatgga ggtagc 35

<210> 191
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 191
cgacctccat cctcgcccag gcatagaaat taact 35

<210> 192
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 192
gctgacactt tccgcgcact cttccgagtc tactc 35

<210> 193
<211> 35
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 193
gagtagatc ggaagagtgc gcggaaagtg tcagc 35

<210> 194
<211> 32
<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 194

ctccggggag cgctggcgct gtacacaggg ga

32

<210> 195

<211> 32

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 195

tcccctgtgt acagcgccag cgctccccgg ag

32

<210> 196

<211> 31

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 196

caaggaggcc gagaaaatca cgacgggctg t

31

<210> 197

<211> 31

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 197

acagcccgtc gtgattttct cggcctcctt g

31

<210> 198

<211> 37

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 198

actgcagctt gaatgagaaa atcactgtcc cagacac

37

<210> 199

<211> 37

<212> DNA

<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 199
gtgtctggga cagtgatttt ctcattcaag ctgcagt 37

<210> 200
<211> 31
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 200
aggccctgtt ggtcaaattct tcccagccgt g 31

<210> 201
<211> 31
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 201
cacggctggg aagatttgac caacagggcc t 31

<210> 202
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 202
atttcctccg gggatggctg aagctgtaca cag 33

<210> 203
<211> 33
<212> DNA
<213> Artificial

<220>
<223> Description of Artificial Sequence: primer

<400> 203
ctgtgtacag cttcagccat ccccggagga aat 33

<210> 204
<211> 35
<212> DNA
<213> Artificial

<220>
 <223> Description of Artificial Sequence: primer

<400> 204
 agccgagtcc tggaggcggc cctcttggag gccaa 35

<210> 205
 <211> 35
 <212> DNA
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: primer

<400> 205
 ttggcctcca agagggccgc ctccaggact cggct 35

<210> 206
 <211> 35
 <212> DNA
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: primer

<400> 206
 agccgagtcc tggagagggc cctcttggag gccaa 35

<210> 207
 <211> 35
 <212> DNA
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: primer

<400> 207
 ttggcctcca agagggccct ctccaggact cggct 35

<210> 208
 <211> 6059
 <212> DNA
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: plasmid

<400> 208
 ctagagtcca cccgggcggc cgcttccctt tagtgagggt taatgcttcg agcagacatg 60
 ataagataca ttgatgatgt tggacaaacc acaactagaa tgcagtgaaa aaaatgcttt 120
 atttgtgaaa tttgtgatgc tattgcttta tttgtaacca ttataagctg caataaacia 180
 gttaacaaca acaattgcat tcattttatg tttcagggtc agggggagat gtgggaggtt 240
 ttttaaagca agtaaaacct ctacaaatgt ggtaaaatcc gataaggatc gatccgggct 300

ggcgtaatag cgaagaggcc cgcaccgatc gcccttccca acagttgcgc agcctgaatg	360
gcgaatggac gcgccctgta gcggcgcatt aagcgcggcg ggtgtggtgg ttacgcgcag	420
cgtgaccgct acacttgcca gcgccctagc gcccgctcct ttcgctttct tcccttcctt	480
tctcgccacg ttcgccggct tccccgtca agctctaaat cgggggctcc ctttaggggt	540
ccgatttagt gctttacggc acctcgacct caaaaaactt gattaggggtg atggttcacg	600
tagtggggcca tcgccctgat agacggtttt tcgccctttg acgttggagt ccacgttctt	660
taatagtgga ctcttggtcc aaactggaac aacactcaac cctatctcgg totattcttt	720
tgatttataa gggattttgc cgatttcggc ctattgggta aaaaatgagc tgatttaaca	780
aaaattttaac gcgaatttta acaaaatatt aacgcttaca atttcctgat gcgggtatttt	840
ctccttacgc atctgtgcgg tatttcacac cgcatacgcg gatctgcgca gcaccatggc	900
ctgaaataac ctctgaaaga ggaacttggg taggtacctt ctgaggcgga aagaaccagc	960
tgtggaatgt gtgtcagtta ggggtgtggaa agtccccagg ctccccagca ggcagaagta	1020
tgcaaagcat gcatctcaat tagtcagcaa ccagggtgtg aaagtcccca ggctccccag	1080
caggcagaag tatgcaaagc atgcatctca attagtcagc aaccatagtc ccgcccctaa	1140
ctccgccccat cccgccccta actccgcccc gttccgcccc ttctccgccc catgggtgac	1200
taattttttt tattttatgca gagggcgagg ccgcctcggc ctctgagcta ttccagaagt	1260
agtgaggagg ctttttttga ggccctaggct tttgcaaaaa gcttgattct tctgacacaa	1320
cagtctcgaa cttaaggcta gagccaccat gattgaacaa gatggattgc acgcagggtc	1380
tccggccgct tgggtggaga ggctattcgg ctatgactgg gcacaacaga caatcggtg	1440
ctctgatgoc gccgtgttcc ggctgtcagc gcagggggcg ccggttcttt ttgtcaagac	1500
cgacctgtcc ggtgccctga atgaactgca ggacgaggca gcgcggctat cgtggctggc	1560
cacgacgggc gttccttgcg cagctgtgct cgacgttgct actgaagcgg gaagggactg	1620
gctgctattg ggcgaagtgc cggggcagga tctcctgtca tctcaccttg ctctgcca	1680
gaaagtatcc atcatggctg atgcaatgcg gcggctgcat acgcttgatc cggtacctg	1740
cccatctgac caccaagcga aacatcgcat cgagcgagca cgtactcgga tggaaagccgg	1800
tcttgtcgat caggatgatc tggacgaaga gcatcagggg ctcgcgccag ccgaactgtt	1860
cgccaggctc aaggcgcgca tgcccagcgg cgaggatctc gtogtgacct atggcgatgc	1920
ctgcttgccg aatatcatgg tggaaaatgg ccgcttttct ggattcatcg actgtggccg	1980
gctgggtgtg gcggaccgct atcaggacat agcgttggct acccgtgata ttgctgaaga	2040
gcttggcgggc gaatgggctg accgcttcct cgtgctttac ggtatcgccg ctcccgattc	2100

gcagcgcac	gccttctatc	gccttcttga	cgagttcttc	tgagcgggac	tctgggggttc	2160
gaaatgaccg	accaagcgac	gccaacctg	ccatcacgat	ggccgcaata	aaatatcttt	2220
atcttcatta	catctgtgtg	ttggtttttt	gtgtgaatcg	atagcgataa	ggatccgcgt	2280
atggtgcact	ctcagtacaa	tctgctctga	tgccgcatag	ttaagccagc	cccgcacccc	2340
gccaacaccc	gctgacgcgc	cctgacgggc	ttgtctgctc	ccggcatccg	cttacagaca	2400
agctgtgacc	gtctccggga	gctgcatgtg	tcagaggttt	tcaccgtcat	caccgaaacg	2460
cgcgagacga	aagggcctcg	tgatacgcc	atcttttatag	gttaatgtca	tgataataat	2520
ggtttcttag	acgtcagggtg	gcacttttctg	gggaaatgtg	cgcggaaccc	ctatttgttt	2580
atctttctaa	atacattcaa	atatgtatcc	gctcatgaga	caataaccct	gataaatgct	2640
tcaataatat	tgaaaaagga	agagtatgag	tattcaacat	ttccgtgtcg	cccttattcc	2700
cttttttgcg	gcattttgcc	ttcctgtttt	tgctcaccca	gaaacgctgg	tgaaagtaaa	2760
agatgctgaa	gatcagttgg	gtgcacgagt	gggttacatc	gaactggatc	tcaacagcgg	2820
taagatcctt	gagagttttc	gccccgaaga	acgttttcca	atgatgagca	cttttaaagt	2880
tctgctatgt	ggcgcggtat	tatcccgtat	tgacgcgggg	caagagcaac	tcggtcgccg	2940
catacactat	tctcagaatg	acttggttga	gtactcacca	gtcacagaaa	agcatcttac	3000
ggatggcatg	acagtaagag	aattatgcag	tgctgccata	accatgagtg	ataacactgc	3060
ggccaactta	cttctgacaa	cgatcggagg	accgaaggag	ctaaccgctt	ttttgcacaa	3120
catgggggat	catgtaactc	gccttgatcg	ttgggaaccg	gagctgaatg	aagccatacc	3180
aaacgcagag	cgtgacacca	cgatgcctgt	agcaatggca	acaacgttgc	gcaaactatt	3240
aactggcgaa	ctacttactc	tagcttcccc	gcaacaatta	atagactgga	tggaggcgga	3300
taaagttgca	ggaccacttc	tgcgctcggc	ccttccggct	ggctggttta	ttgtgataa	3360
atctggagcc	ggtagcgtg	ggtctcggg	tatcattgca	gcactggggc	cagatggtaa	3420
gccctcccgt	atcgtagtta	tctacacgac	ggggagtcag	gcaactatgg	atgaacgaaa	3480
tagacagatc	gctgagatag	gtgcctcact	gattaagcat	tggttaactgt	cagaccaagt	3540
ttactcatat	atactttaga	ttgatttaaa	acttcatttt	taatttaaaa	ggatctaggt	3600
gaagatcctt	tttgataatc	tcattgacaa	aatcccctaa	cgtgagtttt	cgttccactg	3660
agcgtcagac	cccgtagaaa	agatcaaagg	atcttcttga	gatccttttt	ttctgcgcgt	3720
aatctgctgc	ttgcaaacaa	aaaaaccacc	gctaccagcg	gtggtttgtt	tgccggatca	3780
agagctacca	actctttttc	cgaaggtaac	tggttcagc	agagcgcaga	taccaaatat	3840
tgcttctcta	gtgtagccgt	agttaggcca	ccacttcaag	aactctgtag	caccgcctac	3900
atacctcgct	ctgctaatac	tgttaccagt	ggctgctgcc	agtggcgata	agtcgtgtct	3960

taccgggttg gactcaagac gatagttacc ggataaggcg cagcggtcgg gctgaacggg	4020
gggttcgtgc acacagccca gcttggagcg aacgacctac accgaactga gatacctaca	4080
gcgtgagcta tgagaaagcg ccacgcttcc cgaagggaga aaggcggaca ggtatccggt	4140
aagcggcagg gtcggaacag gagagcgcac gagggagctt ccagggggaa acgcctggta	4200
tctttatagt cctgtcgggt ttgccacct ctgacttgag cgtcgatttt tgtgatgctc	4260
gtcagggggg cggagcctat ggaaaaacgc cagcaacgcg gcctttttac ggttcctggc	4320
cttttgctgg ccttttgctc acatggctcg acagatcttc aatattggcc attagccata	4380
ttattcattg gttatatagc ataaatcaat attggctatt ggccattgca tacgttgtat	4440
ctatatcata atatgtacat ttatatggc tcatgtccaa tatgaccgcc atgttggcat	4500
tgattattga ctagttatta atagtaatca attacggggg cattagttca tagcccatat	4560
atggagttcc gcgttacata acttacggta aatggcccg cctggctgacc gcccaacgac	4620
ccccgccc atgagtcagt aatgacgtat gttcccatag taacgccaat agggactttc	4680
cattgacgtc aatgggtgga gtattttacg taaactgccc acttggcagt acatcaagtg	4740
tatcatatgc caagtccgcc ccctattgac gtcaatgacg gtaaattggc cgcttggcat	4800
tatgcccagt acatgacctt acgggacttt cctacttggc agtacatcta cgtattagtc	4860
atcgctatta ccatgggtgat gcggttttgg cagtacacca atgggcgtgg atagcggttt	4920
gactcacggg gattttccaag tctccacccc attgacgtca atgggagttt gttttggcac	4980
caaaatcaac gggactttcc aaaatgtcgt aacaactgcg atcgcccgcc ccgttgacgc	5040
aaatgggogg taggcgtgta cgggtggagg tctatataag cagagctcgt ttagtgaacc	5100
gtcagatcac tagaagcttt attgcggtag tttatcacag ttaaattgct aacgcagtca	5160
gtgcttctga cacaacagtc tcgaacttaa gctgcagtga ctctcttaag gtagccttgc	5220
agaagttggt cgtgaggcac tgggcaggta agtatcaagg ttacaagaca ggtttaaggga	5280
gaccaataga aactgggctt gtcgagacag agaagactct tgcgtttctg ataggcacct	5340
attggtctta ctgacatcca ctttgccctt ctctccacag gtgtccactc ccagttcaat	5400
tacagctctt aaggctagag tacttaatac gactcactat aggctagcct cgagcgcgga	5460
gatgggggtg cacgaatgtc ctgcctggct gtggcttctc ctgtccctgc tgtcgctccc	5520
tctgggcctc ccagtcctgg gcgccccacc acgcctcatc tgtgacagcc gagtcctgga	5580
gaggtacctc ttggaggcca aggaggccga gaatatcacg acgggctgtg ctgaacactg	5640
cagcttgaat gagaatatca ctgtcccaga caccaaagtt aatttctatg cctggaagag	5700
gatggaggtc gggcagcagg ccgtagaagt ctggcagggc ctggccctgc tgtcggaagc	5760

tgtcctgctg	ggccaggccc	tgttggtcaa	ctcttcccag	ccgtgggagc	ccctgcagct	5820
gcatgtggat	aaagccgtca	gtggccttcg	cagcctcacc	actctgcttc	gggctctgctg	5880
agcccagaag	gaagccatct	cccctccaga	tgcggcctca	gctgctccac	tccgaacaat	5940
cactgctgac	actttccgca	aactcttccg	agtctactcc	aatttcctcc	ggggaaagct	6000
gaagctgtac	acaggggagg	cctgcaggac	aggggaccat	catcaccatc	accattgat	6059

<210> 209

<211> 6059

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: plasmid

<400> 209

ctagagtcga	cccgggcggc	cgcttccctt	tagtgagggt	taatgcttcg	agcagacatg	60
ataagataca	ttgatgagtt	tggacaaacc	acaactagaa	tgcaagtga	aaaatgcttt	120
atttgtgaaa	tttgtgatgc	tattgcttta	tttgtaacca	ttataagctg	caataaacia	180
gttaacaaca	acaattgcat	tcatttttatg	tttcagggttc	agggggagat	gtgggagggtt	240
ttttaaaagca	agtaaaacct	ctacaaatgt	ggtaaaatcc	gataaggatc	gatccgggct	300
ggcgtaatag	cgaagaggcc	cgcaccgatc	gcccttccca	acagttgcgc	agcctgaatg	360
gcgaatggac	gcgccctgta	gcggcgcat	aagcgcgcg	ggtgtggtgg	ttacgcgcag	420
cgtgaccgct	acacttgcca	gcgccctagc	gcccgctcct	ttcgctttct	tcccttcctt	480
tctcgccacg	ttcgccggct	ttccccgtca	agctctaaat	cgggggctcc	ctttagggtt	540
ccgatttagt	gctttacggc	acctcgaccc	caaaaaactt	gatttaggggtg	atgggttcacg	600
tagtgggcca	tcgccctgat	agacgggtttt	tcgccctttg	acgttgaggat	ccacgttctt	660
taatagtgga	ctcttggtcc	aaactggaac	aacactcaac	cctatctcgg	tctattcttt	720
tgattttataa	gggattttgc	cgatttcggc	ctattgggtta	aaaaatgagc	tgatttaaca	780
aaaattttaac	gcgaatttta	acaaaatatt	aacgcttaca	atttcctgat	gcgggtatttt	840
ctccttacgc	atctgtgcgg	tatttcacac	cgcatacgcg	gatctgcgca	gcaccatggc	900
ctgaaataac	ctctgaaaga	ggaacttggt	taggtacctt	ctgaggcgga	aagaaccagc	960
tgtggaatgt	gtgtcagtta	gggtgtggaa	agtccccagg	ctccccagca	ggcagaagta	1020
tgcaaagcat	gcatctcaat	tagtcagcaa	ccagggtgtg	aaagtccccca	ggctccccag	1080
caggcagaag	tatgcaaagc	atgcatctca	attagtcagc	aaccatagtc	ccgccccctaa	1140
ctccgccccat	cccgccccct	actccgcccc	gttccgcccc	ttctccgccc	catggctgac	1200
taattttttt	tatttatgca	gaggccgagg	ccgcctcggc	ctctgagcta	ttccagaagt	1260

agtgaggagg ctttttttga ggctagggt tttgcaaaaa gcttgattct tctgacacaa 1320
 cagtctcgaa cttaaggcta gagccaccat gattgaacaa gatggattgc acgcagggtc 1380
 tccggccgct tgggtggaga ggctattcgg ctatgactgg gcacaacaga caatcggctg 1440
 ctctgatgcc gccgtgttcc ggctgtcagc gcagggggcgc ccggttcttt ttgtcaagac 1500
 cgacctgtcc ggtgccctga atgaactgca ggacgaggca gcgcggctat cgtggctggc 1560
 cacgacgggc gttccttgcg cagctgtgct cgacgttgtc actgaagcgg gaagggactg 1620
 gctgctattg ggcgaaagtgc cggggcagga tctcctgtca tctcaccttg ctctgccga 1680
 gaaagtatcc atcatggctg atgcaatgcg gcggctgcat acgcttgatc cggctacctg 1740
 cccattcgac caccaagcga aacatogcat cgagcgagca cgtactcgga tggaagccgg 1800
 tcttgtcgat caggatgatc tggacgaaga gcatcagggg ctgcgccag ccgaactgtt 1860
 cgccaggctc aaggcgcgca tgcccgcgcg cgaggatctc gtcgtgacct atggcgatgc 1920
 ctgcttgccg aatatcatgg tggaaaatgg ccgcttttct ggattcatcg actgtggccg 1980
 gctgggtgtg ggggaccgct atcaggacat agcgttggct acccgtgata ttgctgaaga 2040
 gcttggcggc gaatgggctg accgcttctc cgtgctttac ggtatcgccg ctcccgatc 2100
 gcagcgcac gccttctatc gccttcttga cgagttcttc tgagcgggac tctggggttc 2160
 gaaatgaacc accaagcgac gcccaacctg ccatcacgat ggccgcaata aaatatcttt 2220
 attttcatca catctgtgtg ttgggttttt gtgtgaatcg atagcgataa ggatccgct 2280
 atgggtgcaat ctcaagtacaa tctgctctga tgccgcatag ttaagccagc ccgcacaccc 2340
 gccaacaccc gctgacgcgc cctgacgggc ttgtctgtc ccggcatccg cttacagaca 2400
 agctgtgacc gtctccggga gctgcatgtg tcagaggttt tcaccgtcat caccgaaacg 2460
 cgcgagacga aagggcctcg tgatacgctt atttttatag gttaatgtca tgataataat 2520
 ggtttcttag acgtcagggt gcacttttct gggaaatgtg cgcggaaccc ctattttgtt 2580
 atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct gataaatgct 2640
 tcaataatat tgaaaaagga agagtatgag tattcaacat ttccgtgtcg cccttattcc 2700
 cttttttgcg gcattttgcc ttctgtttt tgctcacca gaaacgctgg tgaaagtaaa 2760
 agatgctgaa gatcagttgg gtgcacgagt gggttacatc gaactggatc tcaacagcgg 2820
 taagatcctt gagagttttc gcccgaaga acgttttcca atgatgagca cttttaaagt 2880
 tctgctatgt ggcgcgggtat tatcccgat tgacgccggg caagagcaac tcggtcgccg 2940
 catacactat tctcagaatg acttggttga gtactacca gtcacagaaa agcatcttac 3000
 ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg ataacactgc 3060

ggccaaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt ttttgcacaa	3120
catgggggat catgtaactc gccttgatcg ttgggaaccg gagctgaatg aagccatacc	3180
aaacgacgag cgtgacacca cgatgcctgt agcaatggca acaacgttgc gcaaactatt	3240
aactggcgaa ctacttactc tagcttcccg gcaacaatta atagactgga tggaggcgga	3300
taaagttgca ggaccacttc tgcgctcggc ccttcgggct ggctggttta ttgctgataa	3360
atctggagcc ggtgagcgtg ggtctcggc tatcattgca gcactggggc cagatggtaa	3420
gccctcccg atcgtagtta tctacacgac ggggagtcag gcaactatgg atgaacgaaa	3480
tagacagatc gctgagatag gtgcctcact gattaagcat tggtaactgt cagaccaagt	3540
ttactcatat atactttaga ttgatttaaa acttcatttt taatttaaaa ggatctaggt	3600
gaagatcctt tttgataatc tcatgaccaa aatcccttaa cgtgagtttt cgttccactg	3660
agcgtcagac cccgtagaaa agatcaaagg atctcttga gatccttttt ttctgcgcgt	3720
aatctgctgc ttgcaaacaa aaaaaccacc gctaccagcg gtggtttgtt tgccggatca	3780
agagctacca actctttttc cgaaggtaac tggttcagc agagcgcaga taccaaatac	3840
tgttcttcta gtgtagccgt agttaggcca ccacttcaag aactctgtag caccgcctac	3900
atacctcgtc ctgctaatac tgttaccagt ggctgctgcc agtggcgata agtcgtgtct	3960
taccgggttg gactcaagac gatagttacc ggataaggcg cagcggtcgg gctgaacggg	4020
gggttcgtgc acacagccca gcttggagcg aacgacctac accgaactga gatacctaca	4080
gcgtgagcta tgagaaagcg ccacgcttcc cgaagggaga aaggcggaca ggtatccggt	4140
aagcggcagg gtcggaacag gagagcgcac gagggagctt ccagggggaa acgcctggta	4200
tctttatagt cctgtcgggt ttgcgccact ctgacttgag cgtcgatttt tgtgatgctc	4260
gtcagggggg cggagcctat ggaaaaacgc cagcaacgcg gcctttttac ggttcctggc	4320
cttttgctgg ctttttgctc acatggctcg acagatcttc aatattggcc attagccata	4380
ttattcattg gttatatagc ataaatcaat attggctatt ggccattgca tacgttgtat	4440
ctatatcata atatgtacat ttatatggc tcatgtccaa tatgacgcgc atgttggcat	4500
tgattattga ctagttatta atagtaatca attacggggg cattagttca tagccatat	4560
atggagttcc gcgttacata acttacggta aatggccgcg ctggctgacc gcccaacgac	4620
ccccgccc atgacgtcaat aatgacgtat gttcccatag taacgccaat agggactttc	4680
cattgacgtc aatgggtgga gtatttacgg taaactgccc acttggcagt acatcaagtg	4740
tatcatatgc caagtccgcc ccctattgac gtcaatgacg gtaaatggcc cgcctggcat	4800
tatgccagtc acatgacctt acgggacttt cctacttggc agtacatcta cgtattagtc	4860
atcgctatta ccatgggtgat gcggttttgg cagtacacca atgggcgtgg atagcggttt	4920

gactcacggg gatttccaag tctccacccc attgacgtca atgggagttt gttttggcac 4980
 caaaatcaac gggactttcc aaaatgtcgt aacaactgcg atcgcccgcc ccgttgacgc 5040
 aaatgggcgg taggcgtgta cgggtgggagg tctatataag cagagctcgt ttagtgaacc 5100
 gtcagatcac tagaagcttt attgcggtag tttatcacag ttaaattgct aacgcagtca 5160
 gtgcttctga cacaacagtc tcgaacttaa gctgcagtga ctctcttaag gtagccttgc 5220
 agaagttggt cgtgaggcac tgggcaggta agtatcaagg ttacaagaca ggtttaagga 5280
 gaccaataga aactgggctt gtcgagacag agaagactct tgcgtttctg ataggcacct 5340
 attggtctta ctgacatcca ctttgccttt ctctccacag gtgtccactc ccagttcaat 5400
 tacagctctt aaggctagag tacttaatac gactcactat aggctagcct cgagcgcgga 5460
 gatgggggtg cacgaatgtc ctgcctggct gtggcttctc ctgtccctgc tgtcgtctcc 5520
 tctgggcctc ccagtcctgg gcgccccacc acgcctcatc tgtgacagcc gagtcctgga 5580
 gaggtacctc ttggaggcca aggaggccga gaatatcacg acgggctgta atgaaacctg 5640
 cagcttgaat gagaatatca ctgtcccaga caccaaagtt aatttctatg cctggaagag 5700
 gatggaggtc gggcagcagg ccgtagaagt ctggcagggc ctggccctgc tgtcgggaagc 5760
 tgtcctgcgg ggccaggccc tgttgggtcaa ctcttcccag ccgtgggagc ccctgcagct 5820
 gcatgtggat aaagccgtca gtggccttcg cagcctcacc actctgcttc gggctctgcg 5880
 agcccagaag gaagccatct cccctccaga tgcggcctca gctgctccac tccgaacaat 5940
 cactgctgac actttccgca aactcttccg agtctactcc aatttcctcc ggggaaagct 6000
 gaagctgtac acaggggagg cctgcaggac aggggaccat catcaccatc accattgat 6059

<210> 210
 <211> 6059
 <212> DNA
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: plasmid

<400> 210
 ctagagtcga cccgggcggc cgttccctt tagtgagggt taatgcttcg agcagacatg 60
 ataagataca ttgatgagtt tggacaaacc acaactagaa tgcagtgaaa aaaatgcttt 120
 atttgtgaaa tttgtgatgc tattgcttta tttgtaacca ttataagctg caataaacia 180
 gttaacaaca acaattgcat tcattttatg tttcagggtc agggggagat gtgggagggt 240
 ttttaaagca agtaaaacct ctacaaatgt ggtaaaatcc gataaggatc gatccgggct 300
 ggcgtaatag cgaagaggcc cgcaccgatc gcccttccca acagttgcgc agcctgaatg 360

gcgaatggac gcgcacctgta ggggcgcatt aagcgcgggcg ggtgtggtgg ttacgcgcag 420
 cgtgaccgct acacttgcca gcgcacctagc gcccgctcct ttcgctttct tcccttcctt 480
 tctcgccacg ttgcgcggct ttccccgtca agctctaaat cgggggctcc ctttaggggtt 540
 ccgatttagt gctttadggc acctcgaccc caaaaaactt gattaggggtg atggttcacg 600
 tagtgggcca tcgcacctgat agacgggtttt tcgcacctttg acgttggagt ccacgttctt 660
 taatagtgga ctcttggtcc aaactggaac aacactcaac cctatctcgg tctattcttt 720
 tgatttataa gggatttttg cgatttcggc ctattgggtta aaaaatgagc tgatttaaca 780
 aaaatttaac gcgaatttta acaaaatatt aacgcttaca atttcctgat gcggtatttt 840
 ctccctacgc atctgtgagg tatttcacac cgcatacgcg gatctgcgca gcaccatggc 900
 ctgaaataac ctctgaaaga ggaacttggg taggtacctt ctgaggcggg aagaaccagc 960
 tgtggaatgt gtgtcagtta ggggtgtgaa agtccccagg ctccccagca ggcagaagta 1020
 tgcaaagcat gcactcaat tagtcagcaa ccagggtgtg aaagtcccca ggctccccag 1080
 caggcagaag tatgcaaagc atgcactca attagtcagc aaccatagtc ccgcccctaa 1140
 ctccgcccct cccgccccta actccgcccc gttccgcccc ttctccgccc catggctgac 1200
 taatTTTTTT tatttatgca gaggccgagg ccgcctcggc ctctgagcta ttccagaagt 1260
 agtgaggagg ctttttttga ggcctaggct tttgcaaaaa gcttgattct tctgacacaa 1320
 cagtctcgaa cttaaggcta gagccaccat gattgaacaa gatggattgc acgcaggttc 1380
 tcgggcgct tgggtggaga ggctattcgg ctatgactgg gcacaacaga caatcggtg 1440
 ctctgatgcc gcctgttcc ggctgtcagc gcaggggcgc ccggttcttt ttgtcaagac 1500
 cgacctgtcc ggtgcacctga atgaactgca ggacgaggca gcgcggctat cgtggctggc 1560
 cacgacgggc gttccttgcg cagctgtgct cgacgttgtc actgaagcgg gaagggactg 1620
 gctgctattg ggogaagtgc cggggcagga tctcctgtca tctcaccttg ctctgccga 1680
 gaaagtatcc atcatggctg atgcaatgcg gcggctgcat acgcttgatc cggctacctg 1740
 cccattcgac caccaagcga aacatcgcat cgagcgagca cgtactcgga tggaagccgg 1800
 tottgtcgat caggatgatc tggacgaaga gcatcagggg ctcgcgccag ccgaactgtt 1860
 cgccaggctc aaggcgcgca tgcccagcgg cgaggatctc gtcgtgacct atggcgatgc 1920
 ctgcttgccg aatatcatgg tggaaaatgg ccgcttttct ggattcatcg actgtggccg 1980
 gctgggtgtg gcggaccgct atcaggacat agcgttggct acccgtgata ttgctgaaga 2040
 gcttggcggc gaatgggctg accgcttctt cgtgctttac ggtatcgccg ctcccgattc 2100
 gcagcgcata gccttctatc gccttcttga cgagttcttc tgagcgggac tctgggggtc 2160
 gaaatgaccg accaagcgac gcccaacctg ccatcacgat ggccgcaata aaatatcttt 2220

attttcatta catctgtgtg ttgggtttttt gtgtgaatcg atagcgataa ggatccgcgt	2280
atggtgcact ctcaagtacaa tctgctctga tgccgcataag ttaagccagc cccgacaccc	2340
gccaacaccc gctgacgcgc cctgacgggc ttgtctgctc ccggcatccg cttacagaca	2400
agctgtgacc gtctccggga gctgcatgtg tcagaggttt tcaccgtcat caccgaaacg	2460
cgcgagacga aagggcctcg tgatacgctt atttttatag gttaatgtca tgataataat	2520
ggtttcttag acgtcagggtg gcacttttctg gggaaatgtg cgcggaaccc ctatttgttt	2580
atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct gataaatgct	2640
tcaataatat tgaaaaagga agagtatgag tattcaacat ttccgtgtcg cccttattcc	2700
cttttttgcg gcattttgcc ttctgtttt tgctcaccga gaaacgctgg tgaaagtaaa	2760
agatgctgaa gatcagttgg gtgcacgagt gggttacatc gaactggatc tcaacagcgg	2820
taagatcctt gagagttttc gccccgaaga acgttttcca atgatgagca cttttaagt	2880
tctgctatgt ggcgcggtat tatcccgat tgacgcggg caagagcaac tcggtcgccg	2940
catacactat tctcagaatg acttggttga gtactacca gtcacagaaa agcatcttac	3000
ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg ataacactgc	3060
ggccaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt ttttgacaaa	3120
catgggggat catgtaactc gccttgatcg ttgggaaccg gagctgaatg aagccatacc	3180
aaacgacgag cgtgacacca cgatgctgt agcaatggca acaacgttgc gcaaactatt	3240
aactggcgaa ctacttactc tagcttcccg gcaacaatta atagactgga tggaggcgga	3300
taaagttgca ggaccacttc tgcgctcggc ccttcgggt ggctggttta ttgctgataa	3360
atctggagcc ggtgagcgtg ggtctcgcg tatcattgca gcaactgggc cagatggtaa	3420
gccctcccg atcgtagtta tctacacgac ggggagtcag gcaactatgg atgaacgaaa	3480
tagacagatc gctgagatag gtgcctcact gattaagcat tggtaactgt cagaccaagt	3540
ttactcatat taactttaga ttgatttaaa acttcatttt taatttaaaa ggatctaggt	3600
gaagatcctt tttgataatc tcatgaccaa aatcccttaa cgtgagtttt cgttccactg	3660
agcgtcagac ccgtagaaa agatcaaagg atcttcttga gatccttttt ttctgcgcgt	3720
aatctgctgc ttgcaaacaa aaaaaccacc gctaccagcg gtgggttggt tgccggatca	3780
agagctacca actctttttc cgaaggtaac tggcttcagc agagcgcaga taccaaatac	3840
tgttcttcta gtgtagccgt agttaggcca ccacttcaag aactctgtag caccgcctac	3900
atacctcgct ctgctaatac tgttaccagt ggctgctgcc agtggcgata agtcgtgtct	3960
taccgggttg gactcaagac gatagttacc ggataaggcg cagcggtcgg gctgaacggg	4020

gggttcgtgc	acacagccca	gcttgagcgc	aacgacctac	accgaactga	gataacctaca	4080
gcgtgagcta	tgagaaagcg	ccacgcttcc	cgaagggaga	aaggcggaca	ggtatccggt	4140
aagcggcagg	gtcggaaacag	gagagcgcac	gagggagcctt	ccagggggaa	acgcctggta	4200
tctttatagt	cctgtcgggt	ttcgccacct	ctgacttgag	cgctgatttt	tgtgatgctc	4260
gtcagggggg	cggagcctat	ggaaaaacgc	cagcaacgcg	gcctttttac	ggttcctggc	4320
cttttgctgg	ccttttgctc	acatggctcg	acagatcttc	aatattggcc	attagccata	4380
ttattcattg	gttatatagc	ataaatcaat	attggctatt	ggccattgca	tacgttgtat	4440
ctatatcata	atatgtacat	ttatatggc	tcatgtccaa	tatgaccgcc	atgttggcat	4500
tgattattga	ctagttatta	atagtaatca	attacgggggt	cattagttca	tagcccatat	4560
atggagttcc	gcgttacata	acttacggta	aatggccgcg	ctggctgacc	gcccacgcac	4620
ccccgccc	tgacgtcaat	aatgacgtat	gttcccatag	taacgccaat	agggactttc	4680
cattgacgtc	aatgggtgga	gtatttacgg	taaactgccc	acttggcagt	acatcaagtg	4740
tatcatatgc	caagtcgcc	ccctattgac	gtcaatgacg	gtaaatggcc	cgctggcat	4800
tatgccaggt	acatgacctt	acgggacttt	cctacttggc	agtacatcta	cgtattagtc	4860
atcgctatta	ccatgggtgat	gcggttttgg	cagtacacca	atgggcgtgg	atagcggttt	4920
gactcacggg	gatttccaag	tctccacccc	attgacgtca	atgggagttt	gttttggcac	4980
caaatcaac	gggactttcc	aaaatgtcgt	aacaactgcg	atcgcccgcc	ccgttgacgc	5040
aaatgggcgg	taggcgtgta	cggtgggagg	tctatataag	cagagctcgt	ttagtgaacc	5100
gtcagatcac	tagaagcttt	attgcggtag	tttatcacag	ttaaattgct	aaocgagtca	5160
gtgcttctga	cacaacagtc	tcgaacttaa	gctgcagtga	ctctcttaag	gtagccttgc	5220
agaagttgg	cgtgaggcac	tgggcaggta	agtatcaagg	ttacaagaca	ggtttaagga	5280
gaccaataga	aactgggctt	gtcgagacag	agaagactct	tgcgtttctg	ataggcacct	5340
attggctctta	ctgacatcca	ctttgccttt	ctctccacag	gtgtccactc	ccagttcaat	5400
tacagctctt	aaggctagag	tacttaatac	gactcactat	aggctagcct	cgagcgcgga	5460
gatgggggtg	cacgaatgtc	ctgcctggct	gtggcttctc	ctgtccctgc	tgtcgctccc	5520
tctgggcctc	ccagtcctgg	gcgccccacc	acgcctcatc	tgtgacagcc	gagtcctgga	5580
gaggtacctc	ttggaggcca	aggaggccga	gaatatcacg	acgggctgtg	ctgaacactg	5640
cagcttgaat	gagaatatca	ctgtcccaga	caccgacgtt	aatttctatg	cctggaagag	5700
gatggaggtc	gggcagcagg	ccgtagaagt	ctggcagggc	ctggccctgc	tgtcggaagc	5760
tgtcctgcgg	ggccaggccc	tgttgggtcaa	ctcttcccag	ccgtgggagc	ccctgcagct	5820
gcatgtggat	aaagccgtca	gtggccttcg	cagcctcacc	actctgcttc	gggctctgcg	5880

agcccagaag gaagccatct cccctccaga tgcggcctca gctgctccac tccgaacaat 5940
 cactgctgac actttccgca aactcttccg agtctactcc aatttcctcc ggggaaagct 6000
 gaagctgtac acaggggagg cctgcaggac aggggaccat catcaccatc accattgat 6059

<210> 211
 <211> 6059
 <212> DNA
 <213> Artificial

<220>
 <223> Description of Artificial Sequence: plasmid

<400> 211
 ctagagtcga cccgggcggc cgcttccctt tagtgagggt taatgcttcg agcagacatg 60
 ataagataca ttgatgagtt tggacaaacc acaactagaa tgcagtgaaa aaaatgcttt 120
 atttgtgaaa tttgtgatgc tattgcttta tttgtaacca ttataagctg caataaacia 180
 gttaacaaca acaattgcat tcattttatg tttcagggtc agggggagat gtgggaggtt 240
 ttttaaagca agtaaaacct ctacaaatgt ggtaaaatcc gataaggatc gatccgggct 300
 ggcgtaatag cgaagaggcc cgcaccgatc gcccttccca acagttgcgc agcctgaatg 360
 gcgaatggac gcgccttgta gcggcgcatc aagcgcggcg ggtgtggtgg ttacgcgcag 420
 cgtgaccgct acacttgcca gcgccttagc gcccgctcct ttcgctttct tcccttccct 480
 tctcgccacg ttcgcgggct ttccccgtca agctctaaat cgggggctcc ctttaggggt 540
 ccgatttagt gctttacggc acctcgacct caaaaaactt gattaggggt atgggttcacg 600
 tagtgggcca tcgcctgat agacgggttt tgcgcctttg acgttgaggt ccacgttctt 660
 taatagtgga ctcttggtcc aaactggaac aacactcaac cctatctcgg tctattcttt 720
 tgatttataa gggattttgc cgatttcggc ctatttggtta aaaaatgagc tgatttaaca 780
 aaaatttaac gcgaatttta acaaaatatt aacgcttaca atttcctgat gcggtatttt 840
 ctccctacgc atctgtgcgg tatttcacac cgcatacgcg gatctgcgca gcaccatggc 900
 ctgaaataac ctctgaaaga ggaacttggt taggtacctt ctgaggcgga aagaaccagc 960
 tgtggaatgt gtgtcagtta ggggtggtga agtccccagg ctcccagca ggcagaagta 1020
 tgcaaagcat gcatctcaat tagtcagcaa ccagggtgtg aaagtcccca ggctccccag 1080
 caggcagaag tatgcaaagc atgcatctca attagtcagc aaccatagtc ccgcccctaa 1140
 ctccgcccct cccgccccta actccgcccc gttccgcccc ttctccgccc catggctgac 1200
 taattttttt tatttatgca gaggccgagg ccgcctcggc ctctgagcta ttccagaagt 1260
 agtgaggagg ctttttttga ggcctaggct tttgcaaaaa gcttgattct tctgacacaa 1320

cagtctcgaa cttaaggcta gagccaccat gattgaacaa gatggattgc acgcaggttc 1380
 tccggccgct tgggtggaga ggctattcgg ctatgactgg gcacaacaga caatcggctg 1440
 ctctgatgcc gccgtgttcc ggctgtcagc gcagggggcg cgggttcttt ttgtcaagac 1500
 cgacctgtcc ggtgccctga atgaactgca ggacgaggca gcgcggctat cgtggctggc 1560
 cagcaggggc gttccttgcg cagctgtgct cgacgttgct actgaagcgg gaagggactg 1620
 gctgctattg ggcgaaagtgc cggggcagga tctcctgtca tctcaccttg ctctgccga 1680
 gaaagtatcc atcatggctg atgcaatgcg gcggctgcat acgcttgatc cggctacctg 1740
 cccattcgac caccaagcga aacatcgcat cgagcgagca cgtactcgga tggaaagccg 1800
 tcttgctgat caggatgatc tggacgaaga gcatcagggg ctcgcgccag ccgaactgtt 1860
 cgccaggctc aaggcgcgca tgcccagcgg cgaggatctc gtcgtgacct atggcgatgc 1920
 ctgcttgccg aatatcatgg tggaaaatgg ccgcttttct ggattcatcg actgtggccg 1980
 gctgggtgtg gcggaccgct atcaggacat agcgttggct acccgtgata ttgctgaaga 2040
 gcttggcggc gaatgggctg accgcttctc cgtgctttac ggtatcgccg ctcccgattc 2100
 gcagcgcac gccttctatc gccttcttga cgagttcttc tgagcgggac tctggggttc 2160
 gaaatgaccg accaagcgcg gcccaacctg coactacgat ggccgcaata aaatatcttt 2220
 attttcatta catctgtgtg ttgggttttt gtgtgaatcg atagcgataa ggatccgcgt 2280
 atgggtgact ctcaagtacaa tctgctctga tgccgcatag ttaagccagc cccgacacct 2340
 gccaacacct gctgacgcgc cctgacgggc ttgtctgctc ccggcatccg cttacagaca 2400
 agctgtgacc gtctccggga gctgcatgtg tcagaggttt tcaccgtcat caccgaaacg 2460
 cgcgagacga aagggcctcg tgatacgctt atttttatag gttaatgtca tgataataat 2520
 ggtttcttag acgtcagggtg gcacttttgc gggaaatgtg cgcggaacct ctatttgttt 2580
 atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct gataaatgct 2640
 tcaataatat tgaaaaagga agagtatgag tattcaacat ttccgtgtcg cccttattcc 2700
 cttttttgcg gcattttgcc ttctgtttt tgcctaccca gaaacgctgg tgaaagtaaa 2760
 agatgctgaa gatcagttgg gtgcacgagt ggggttacat gaactggatc tcaacagcgg 2820
 taagatcctt gagagttttc gcccgaaga acgttttcca atgatgagca cttttaagt 2880
 tctgctatgt ggcgcggtat tatcccgat tgacgcggg caagagcaac tcggctgccc 2940
 catacactat tctcagaatg acttggttga gtactacca gtcacagaaa agcatcttac 3000
 ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg ataacactgc 3060
 ggccaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt ttttgcacaa 3120
 catgggggat catgtaacct gccttgatcg ttgggaaccg gagctgaatg aagccatacc 3180

aaacgacgag	cgtgacacca	cgatgcctgt	agcaatggca	acaacgttgc	gcaaactatt	3240
aactggcgaa	ctacttactc	tagcttcccg	gcaacaatta	atagactgga	tggaggcgga	3300
taaagttgca	ggaccacttc	tgcgctcggc	ccttccggct	ggctggttta	ttgctgataa	3360
atctggagcc	ggtgagcgtg	ggtctcgccg	tatcattgca	gcactggggc	cagatggtaa	3420
gccctcccg	atcgtagtta	tctacacgac	ggggagtcag	gcaactatgg	atgaacgaaa	3480
tagacagatc	gctgagatag	gtgcctcact	gattaagcat	tggtaactgt	cagaccaagt	3540
ttactcatat	atactttaga	ttgatttaaa	acttcatttt	taatttaaaa	ggatctaggt	3600
gaagatccct	tttgataatc	tcatgaccaa	aatcccttaa	cgtgagtttt	cgttccactg	3660
agcgtcagac	cccgtagaaa	agatcaaagg	atcttcttga	gatccttttt	ttctgcgcgt	3720
aatctgctgc	ttgcaaacaa	aaaaaccacc	gctaccagcg	gtggtttggt	tgccggatca	3780
agagctacca	actctttttc	cgaaggtaac	tggcttcagc	agagcgcaga	taccaataac	3840
tgttcttcta	gtgtagccgt	agttaggcca	ccacttcaag	aactctgtag	caccgcctac	3900
atacctcgct	ctgctaatac	tgttaccagt	ggctgctgcc	agtggcgata	agtcgtgtct	3960
taccgggttg	gactcaagac	gatagttacc	ggataaggcg	cagcggtcgg	gctgaacggg	4020
gggttcgtgc	acacagccca	gcttggagcg	aacgacctac	accgaactga	gatacctaca	4080
gcgtgagcta	tgagaaagcg	ccacgcttcc	cgaagggaga	aaggcggaca	ggtatccggt	4140
aagcggcagg	gtcggaacag	gagagcgcac	gaggagcctt	ccagggggaa	acgcctggta	4200
tctttatagt	cctgtcgggt	ttcgccacct	ctgacttgag	cgtcgatttt	tgtgatgctc	4260
gtcagggggg	cggagcctat	ggaaaaacgc	cagcaacgcg	gccttttttac	ggttcctggc	4320
cttttgctgg	ccttttgctc	acatggctcg	acagatcttc	aatattggcc	attagccata	4380
ttattcattg	gttatatagc	ataaatcaat	attggctatt	ggccattgca	tacgttgtat	4440
ctatatcata	atatgtacat	ttatattggc	tcatgtccaa	tatgaccgcc	atgttggcat	4500
tgattattga	ctagttatta	atagtaatca	attacggggg	cattagttca	tagcccatat	4560
atggagttcc	gcgttacata	acttacggta	aatggcccg	ctggctgacc	gccaacgac	4620
ccccgcccat	tgacgtcaat	aatgacgtat	gttcccatag	taacgccaat	agggactttc	4680
cattgacgtc	aatgggtgga	gtatttacgg	taaactgccc	acttggcagt	acatcaagtg	4740
tatcatatgc	caagtccgcc	ccctattgac	gtcaatgacg	gtaaatggcc	cgcttggcat	4800
tatgcccgat	acatgacctt	acgggacttt	cctacttggc	agtacatcta	cgtattagtc	4860
atcgctatta	ccatgggtgat	gcggttttgg	cagtacacca	atgggcgtgg	atagcggttt	4920
gactcacggg	gatttccaag	tctccacccc	attgacgtca	atgggagttt	gttttggcac	4980

```

caaaatcaac gggactttcc aaaatgtcgt aacaactgcg atcgcccgcc ccgttgacgc 5040
aaatgggccc taggcgtgta cgggtgggagg tctatataag cagagctcgt ttagtgaacc 5100
gtcagatcac tagaagcttt attgcggtag tttatcacag ttaaattgct aacgcagtca 5160
gtgctttctga cacaacagtc tcgaacttaa gctgcagtga ctctcttaag gtagccttgc 5220
agaagttggt cgtgaggcac tgggcaggta agtatcaagg ttacaagaca ggtttaagga 5280
gaccaataga aactgggctt gtcgagacag agaagactct tgcgtttctg ataggcacct 5340
attggtctta ctgacatcca ctttgccttt ctctccacag gtgtccactc ccagttcaat 5400
tacagctctt aaggctagag tacttaatac gactcactat aggctagcct cgagcgcgga 5460
gatgggggtg cacgaatgtc ctgcctggct gtggcttctc ctgtccctgc tgtcgcctcc 5520
tctgggcctc ccagtcctgg gcgccccacc acgcctcatc tgtgacagcc gagtcctgga 5580
gaggtacctc ttggaggcca aggaggccga gaatatcacg acgggctgtg ctgaacactg 5640
cagcttgaat gagaatatca ctgtcccaga caccaaagtt aatttctatg cctggaagag 5700
gatggaggtc gggcagcagg ccgtagaagt ctggcagggc ctggccctgc tgtcgggaagc 5760
tgtcctgcgg ggccaggccc tgttgggtcaa ctcttcccag ccgtgggagc cctgcagct 5820
gcatgtggat aaagccgtcg agggccttcg cagcctcacc actctgcttc gggctctgcg 5880
agcccagaag gaagccatct cccctccaga tgcggcctca gctgctccac tccgaacaat 5940
cactgctgac actttccgca aactcttccg agtctactcc aatttctctc ggggaaagct 6000
gaagctgtac acaggggagg cctgcaggac aggggaccat catcaccatc accattgat 6059

```

<210> 212

<211> 6059

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: plasmid

<400> 212

```

ctagagtcga cccggggcgc cgcttccctt tagtgagggt taatgcttcg agcagacatg 60
ataagataca ttgatgagtt tggacaaacc acaactagaa tgcagtgaaa aaaatgcttt 120
atttgtgaaa tttgtgatgc tattgcttta tttgtaacca ttataagctg caataaaca 180
gttaacaaca acaattgcat tcattttatg tttcagggtc agggggagat gtgggaggtt 240
ttttaaagca agtaaaacct ctacaaatgt ggtaaaatcc gataaggatc gatccgggct 300
ggcgtaatag cgaagaggcc cgcaccgatc gcccttccca acagttgcgc agcctgaatg 360
gcgaatggac ggcgcctgta gcggcgcatc aagcgcggcg ggtgtggtgg ttacgcgcag 420
cgtgaccgct acacttgcca gcgccttagc gcccgctcct ttcgctttct tcccttcctt 480

```

tctcgccacg ttcgccggct ttccccgtca agctctaaat cgggggctcc ctttaggggtt 540
 ccgatttagt gctttacggc acctcgaccc caaaaaactt gattaggggtg atgggttcacg 600
 tagtggggcca tcgccctgat agacgggttt tcgccctttg acgttggagt ccacgttctt 660
 taatagtgga ctcttggtcc aaactggaac aacactcaac cctatctcgg totattcttt 720
 tgatttataa gggatttttg cgatttcggc ctattggtta aaaaatgagc tgatttaaca 780
 aaaatttaac gcgaatttta acaaaatatt aacgcttaca atttcctgat gcgggtatttt 840
 ctccctacgc atctgtgcgg tatttcacac cgcatacgcg gatctgcgca gcaccatggc 900
 ctgaaataac ctctgaaaga ggaacttggt taggtacctt ctgaggcgga aagaaccagc 960
 tgtggaatgt gtgtcagtta ggggtgtgga agtccccagg ctccccagca ggcagaagta 1020
 tgcaaagcat gcctctcaat tagtcagcaa ccagggtgtg aaagtcccca ggctccccag 1080
 caggcagaag tatgcaaagc atgcctctca attagtcagc aaccatagtc ccgcccctaa 1140
 ctccgcccct cccgccccta actccgccc gttccgccc ttctccgccc catggctgac 1200
 taattttttt tatttatgca gaggccgagg ccgcctcggc ctctgagcta ttccagaagt 1260
 agtgaggagg ctttttttga ggcctaggct tttgcaaaaa gcttgattct tctgacacaa 1320
 cagtctcgaa ctttaaggcta gagccaccat gattgaacaa gatggattgc acgcaggttc 1380
 tccggccgct tgggtggaga ggctattcgg ctatgactgg gcacaacaga caatcggctg 1440
 ctctgatgcc gccgtgttcc ggctgtcagc gcagggggcg ccggttcttt ttgtcaagac 1500
 cgacctgtcc ggtgccctga atgaactgca ggacgaggca gcgcggctat cgtggctggc 1560
 cacgaagggc gttccttgcg cagctgtgct cgacgttgct actgaagcgg gaagggactg 1620
 gctgctattg ggcgaagtgc cggggcagga tctcctgtca tctcaccttg ctccctgccga 1680
 gaaagtatcc atcatggctg atgcaatgcg gcggctgcat acgcttgatc cggctacctg 1740
 cccattcgac caccaagcga aacatcgcat cgagcgagca cgtactcgga tggaagccgg 1800
 tcttgtcgat caggatgatc tggacgaaga gcctcagggg ctccgcccag ccgaactgtt 1860
 cgccaggctc aaggcgcgca tgcccgacgg cgaggatctc gtcgtgacct atggcgatgc 1920
 ctgcttgccg aatatcatgg tggaaaatgg ccgcttttct ggattcatcg actgtggccg 1980
 gctgggtgtg gcggaccgct atcaggacat agcgttggtt acccgtgata ttgctgaaga 2040
 gcttggcggc gaatgggctg accgcttcct cgtgctttac ggtatcgccg ctcccgattc 2100
 gcagcgcata gccttctatc gccttcttga cgagttcttc tgagcgggac tctgggggtt 2160
 gaaatgaccg accaagcgac gcccaacctg ccatcacgat ggccgcaata aaatatcttt 2220
 attttcatta catctgtgtg ttgggttttt gtgtgaatcg atagcgataa ggatccgcgt 2280

atggtgcact ctcagtacaa totgctctga tgccgcatag ttaagccagc cccgacaccc 2340
 gccaacaccc gctgacgcgc cctgacgggc ttgtctgctc ccggcatccg cttacagaca 2400
 agctgtgacc gtctccggga gctgcatgtg tcagaggttt tcaccgtcat caccgaaacg 2460
 cgcgagacga aagggcctcg tgatacgccct atttttatag gttaatgtca tgataataat 2520
 ggtttcttag acgtcagggtg gcacttttcg gggaaatgtg cgcggaaccc ctatttgttt 2580
 atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct gataaatgct 2640
 tcaataatat tgaaaaagga agagtatgag tattoaacat ttccgtgtcg ccttatttcc 2700
 cttttttgcg gcatttttgcc ttctgtttt tgctcaccca gaaacgctgg tgaaagtaaa 2760
 agatgctgaa gatcagttgg gtgcacgagt gggttacatc gaactgggac tcaacagcgg 2820
 taagatcctt gagagttttc gcccgaaga acgttttcca atgatgagca cttttaaagt 2880
 tctgctatgt ggcgcggtat tatcccgat tgacgcggg caagagcaac tcgggtcgccg 2940
 catacactat tctcagaatg acttggttga gtactcacca gtcacagaaa agcatcttac 3000
 ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg ataactctgc 3060
 ggccaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt ttttgcaaaa 3120
 catgggggat catgtaactc gccttgatcg ttgggaaccg gagctgaatg aagccatacc 3180
 aaacgacgag cgtgacacca cgatgctctg agcaatggca acaacgcttc gcaaactatt 3240
 aactggcgaa ctacttactc tagcttcccg gcaacaatta atagactgga tggaggcgga 3300
 taaagttgca ggaccacttc tgctgctcggc ccttccggct ggctggttta ttgctgataa 3360
 atctggagcc ggtgagcgtg ggtctcgcgg tatcattgca gcactggggc cagatggtaa 3420
 gccctcccgat atcgtagtga tctacacgac ggggagtcag gcaactatgg atgaacgaaa 3480
 tagacagatc gctgagatag gtgcctcact gattaagcat tggtaactgt cagaccaagt 3540
 ttactcatat atactttaga ttgatttaaa acttcatttt taatttaaaa ggatctaggt 3600
 gaagatcctt tttgataatc tcatgaccaa aatcccttaa cgtgagtttt cgttccactg 3660
 agcgtcagac ccggtagaaa agatcaaagg atcttcttga gatccttttt ttctgcgcgt 3720
 aatctgctgc tggcaaacaa aaaaaccacc gctaccagcg gtggtttggt tgccggatca 3780
 agagctacca actctttttc cgaaggtaac tggcttcagc agagcgcaga taccaaatac 3840
 tgttcttcta gtgtagccgt agttaggcca ccacttcaag aactctgtag caccgcctac 3900
 atacctcgct ctgctaatac tgttaccagt ggctgctgcc agtggcgata agtcgtgtct 3960
 taccgggttg gactcaagac gatagttacc ggataaggcg cagcggtcgg gctgaacggg 4020
 gggttcgtgc acacagccca gcttgagcgg aacgacctac accgaactga gatactaca 4080
 gcgtgagcta tgagaaagcg ccacgcttcc cgaagggaga aaggcggaca ggtatccggt 4140

aagcggcagg gtcggaacag gagagcgac gagggagctt ccagggggaa acgcctggta 4200
tctttatagt cctgtcgggt ttgcacaact ctgacttgag cgtcgatttt tgtgatgctc 4260
gtcagggggg cggagcctat ggaaaaacgc cagcaacgcg gcctttttac ggttcctggc 4320
cttttgctgg ccttttgctc acatggctcg acagatcttc aatattggcc attagccata 4380
ttattcattg gttatatagc ataaatcaat attggctatt ggccattgca tacgttgat 4440
ctatatcata atatgtacat ttatattggc tcatgtccaa tatgaccgcc atgttggcat 4500
tgattattga ctagttatta atagtaatca attacggggg cattagttca tagccatata 4560
atggagttcc gcgttacata acttacggta aatggccgcg ctggctgacc gcccacacgac 4620
ccccgccat tgacgtcaat aatgacgtat gttcccatag taacgccat agggactttc 4680
cattgacgtc aatgggtgga gtatttacgg taaactgccc acttggcagt acatcaagt 4740
tatcatatgc caagtccgcc cctattgac gtcaatgacg gtaaatggcc cgctggcat 4800
tatgccagc acatgacctt acgggacttt cctacttgcc agtacatcta cgtattagtc 4860
atcgctatta ccattggtgat gcggttttgg cagtacacca atgggcgtgg atagcggttt 4920
gactcacggg gatttccaag tctccacccc attgacgtca atgggagttt gttttggcac 4980
caaatcaac gggactttcc aaaatgtcgt acaactgcg atcgccgcc ccgttgacgc 5040
aaatgggcgg taggcgtgta cgggtggagg tctataaag cagagctcgt ttagtgaacc 5100
gtcagatcac tagaagcttt attgcggtag tttatcacag ttaaattgct aacgcagtca 5160
gtgcttctga cacaacagtc tcgaactt aa gctgcagtga ctctcttaag gtagccttgc 5220
agaagttggc cgtgaggcac tgggcaggta agtatcaagg ttacaagaca ggtttaagga 5280
gaccaataga aactgggctt gtcgagacag agaagactct tgcgtttctg ataggcacct 5340
attggtctta ctgacatcca ctttgccttt ctctccacag gtgtccactc ccagttcaat 5400
tacagctctt aaggctagag taatttaatac gactcactat aggctagcct cgagcgcgga 5460
gatgggggtg cacgaatgtc ctgcctggct gtggcttctc ctgtccctgc tgcgctccc 5520
tctgggcctc ccagtcctgg gcgccccacc acgcctcatc tgtgacagcc ggtcctgga 5580
gaggtacctc ttggaggcca aggaggccga gaatatcacg acgggctgtg ctgaacactg 5640
cagcttgaat gagaatatca ctgtcccaga caccgacgtt aatttctatg cctggaagag 5700
gatggaggtc gggcagcagg ccgtagaagt ctggcagggc ctggccctgc tgtcggaagc 5760
tgtcctgcgg ggccaggccc tgttgggtcaa ctctcccag ccgtgggagc ccctgcagct 5820
gcatgtggat aaagccgtcg agggccttcg cagcctcacc actctgcttc gggctctgcg 5880
agcccagaag gaagccatct cccctccaga tgcggcctca gctgctccac tccgaacaat 5940

caatgctgac actttccgca aactcttccg agtctactcc aatttcctcc ggggaaagct 6000
gaagctgtac acaggggagg cctgcaggac aggggaccat catcaccatc accattgat 6059